

Differential Geometry of Curves and Surfaces

Homework 7

Due on November 17

1. Compute the first and second fundamental forms, the mean curvature and the Gauss curvature of the following surfaces:
 - (a) The cylinder $S_1 = \{(x, y, z) \in \mathbb{R}^3 : x^2 + y^2 = R^2\}$, using the parameterization $\mathbf{g}_1 : (0, 2\pi) \times \mathbb{R} \rightarrow S_1$ given by $\mathbf{g}_1(\varphi, z) = (R \cos \varphi, R \sin \varphi, z)$;
 - (b) The cone $S_2 = \{(x, y, z) \in \mathbb{R}^3 : x^2 + y^2 = z^2, z > 0\}$, using the parameterization $\mathbf{g}_2 : (0, 2\pi) \times \mathbb{R}^+ \rightarrow S_2$ given by $\mathbf{g}_2(\varphi, z) = (z \cos \varphi, z \sin \varphi, z)$;
 - (c) The catenoid $S_3 = \{(x, y, z) \in \mathbb{R}^3 : x^2 + y^2 = \cosh^2 z\}$, using the parameterization $\mathbf{g}_3 : (0, 2\pi) \times \mathbb{R} \rightarrow S_3$ given by $\mathbf{g}_3(\varphi, z) = (\cosh z \cos \varphi, \cosh z \sin \varphi, z)$;
 - (d) The sphere $S_4 = \{(x, y, z) \in \mathbb{R}^3 : x^2 + y^2 + z^2 = R^2\}$, using the parameterization $\mathbf{g}_4 : (0, \pi) \times (0, 2\pi) \rightarrow S_4$ given by $\mathbf{g}_4(\theta, \varphi) = (R \sin \theta \cos \varphi, R \sin \theta \sin \varphi, R \cos \theta)$.