

Geometric Mechanics

Homework 6

Due on November 4

1. Let M be an n -dimensional differentiable manifold and $\omega \in \Omega^1(M)$. Show that:

(a) If $X, Y \in \mathfrak{X}(M)$ then

$$d\omega(X, Y) = X \cdot \omega(Y) - Y \cdot \omega(X) - \omega([X, Y]).$$

(b) If ω does not vanish on M then the $(n - 1)$ -dimensional distribution determined by the kernel of ω is integrable if and only if

$$d\omega \wedge \omega = 0.$$

(Hint: Write $d\omega$ in a local basis of one-forms of the type $\{\theta^1, \dots, \theta^{n-1}, \omega\}$).

2. Recall that our model for an ice skate is given by the non-holonomic constraint Σ determined on $\mathbb{R}^2 \times S^1$ by the kernel of the 1-form $\omega = -\sin \theta dx + \cos \theta dy$.

(a) Show that the ice skate can access all points in the configuration space: given two points $p, q \in \mathbb{R}^2 \times S^1$, there exists a piecewise smooth curve $c : [0, 1] \rightarrow \mathbb{R}^2 \times S^1$, compatible with Σ , such that $c(0) = p$ and $c(1) = q$. Why does this show that Σ is non-integrable?

(b) Assuming that the kinetic energy of the skate is

$$K = \frac{M}{2} \left((v^x)^2 + (v^y)^2 \right) + \frac{I}{2} \left(v^\theta \right)^2$$

and that the reaction force is perfect, show that the skate moves with constant speed along straight lines or circles. What is the physical interpretation of the reaction force?