

The mathematics of waves and black holes

Master in Applied Mathematics and Computation

José Natário


Departamento de Matemática

Instituto Superior Técnico


jnatar@math.tecnico.ulisboa.pt

Waves








Linear waves



Sound waves




(NavSource Naval History)



Electromagnetic waves





Gravitational waves



Mathematical definition of a wave

Prototype: traveling wave with velocity c .

$$u(x, t) = f(x - ct), \quad c \in \mathbb{R}$$



Wave equation in one dimension

D'Alembert (1747) deduced the one-dimensional wave equation to study the vibration of a string:

$$\frac{1}{c^2} \frac{\partial^2 u}{\partial t^2} = \frac{\partial^2 u}{\partial x^2} \quad \left(c^2 = \frac{T}{\mu} \right)$$

Changing variables to $\xi = x - ct$, $\eta = x + ct$, the wave equation is written

$$\frac{\partial^2 u}{\partial \xi \partial \eta} = 0,$$

and can be solved by integrating twice:

$$u = f(\xi) + g(\eta) = f(x - ct) + g(x + ct)$$

Wave equation in three dimensions

The generalization to three dimensions is obvious:

$$\frac{1}{c^2} \frac{\partial^2 u}{\partial t^2} = \frac{\partial^2 u}{\partial x^2} + \frac{\partial^2 u}{\partial y^2} + \frac{\partial^2 u}{\partial z^2}$$

This is a linear partial differential equation, and gives a good description of (linear) sound waves, electromagnetic waves, and (linear) gravitational waves.

It admits a conserved energy:

$$E(t) = \int_{\mathbb{R}^3} \left[\frac{1}{c^2} \left(\frac{\partial u}{\partial t} \right)^2 + \left(\frac{\partial u}{\partial x} \right)^2 + \left(\frac{\partial u}{\partial y} \right)^2 + \left(\frac{\partial u}{\partial z} \right)^2 \right] dx dy dz$$

If u is a solution of the wave equation, then so are $\frac{\partial u}{\partial x}, \frac{\partial u}{\partial y}, \frac{\partial u}{\partial z}$. Therefore we also have the conserved generalized energies



$$E_x(t) = \int_{\mathbb{R}^3} \left[\frac{1}{c^2} \left(\frac{\partial^2 u}{\partial t \partial x} \right)^2 + \left(\frac{\partial^2 u}{\partial x^2} \right)^2 + \left(\frac{\partial^2 u}{\partial y \partial x} \right)^2 + \left(\frac{\partial^2 u}{\partial z \partial x} \right)^2 \right] dx dy dz$$



$$E_y(t) = \int_{\mathbb{R}^3} \left[\frac{1}{c^2} \left(\frac{\partial^2 u}{\partial t \partial y} \right)^2 + \left(\frac{\partial^2 u}{\partial x \partial y} \right)^2 + \left(\frac{\partial^2 u}{\partial y^2} \right)^2 + \left(\frac{\partial^2 u}{\partial z \partial y} \right)^2 \right] dx dy dz$$

$$E_z(t) = \int_{\mathbb{R}^3} \left[\frac{1}{c^2} \left(\frac{\partial^2 u}{\partial t \partial z} \right)^2 + \left(\frac{\partial^2 u}{\partial x \partial z} \right)^2 + \left(\frac{\partial^2 u}{\partial y \partial z} \right)^2 + \left(\frac{\partial^2 u}{\partial z^2} \right)^2 \right] dx dy dz$$


These can be used together with **Sobolev's inequality** to show that solutions with finite generalized energies are bounded (and even that they decay in time).

Black holes





Photonsphere



Wave equation on a black hole background

The spacetime around a spherical black hole of mass M is modeled by the Schwarzschild Lorentzian metric:

$$g = -V dt \otimes dt + V^{-1} dr \otimes dr + r^2 (d\theta \otimes d\theta + \sin^2 \theta d\varphi \otimes d\varphi)$$

where

$$V = 1 - \frac{2M}{r}$$

One can still write the wave equation by using the Levi-Civita connection of this metric:

$$d \star du = 0 \Leftrightarrow \nabla^\alpha \nabla_\alpha u = 0$$

If $M = 0$ this is the usual wave equation written in spherical coordinates.

One can use the divergence theorem to prove that again there is a conserved energy:

$$E(t) = \int_{\Sigma_t} T \left(\frac{\partial}{\partial t}, \frac{\partial}{\partial t} \right) \left(1 - \frac{2M}{r} \right)^{-1} r^2 \sin \theta dr d\theta d\varphi$$

where

$$T = du \otimes du - \frac{1}{2} g^{-1}(du, du) g$$

is the energy-momentum tensor associated to u . Again using the Sobolev inequality, one can prove boundedness (and even decay) of finite energy solutions. New challenges: the event horizon and photonsphere trapping.

Are they stable?

