

TORIC CONSTRUCTIONS OF MONOTONE LAGRANGIAN SUBMANIFOLDS IN $\mathbb{C}\mathbb{P}^2$ AND $\mathbb{C}\mathbb{P}^1 \times \mathbb{C}\mathbb{P}^1$

MIGUEL ABREU AND AGNÈS GADBLED

ABSTRACT. We extract from a toric model of the Chekanov-Schlenk exotic torus in $\mathbb{C}\mathbb{P}^2$ methods for constructing Lagrangian submanifolds in toric symplectic manifolds. These constructions allow for some control of monotonicity. We recover this way some known monotone Lagrangians in the toric symplectic manifolds $\mathbb{C}\mathbb{P}^2$ and $\mathbb{C}\mathbb{P}^1 \times \mathbb{C}\mathbb{P}^1$ as well as new examples.

1. INTRODUCTION

One can reconstruct $\mathbb{R}\mathbb{P}^2$, the real part of $\mathbb{C}\mathbb{P}^2$ with respect to the standard conjugation map, from its image under the standard moment map of $\mathbb{C}\mathbb{P}^2$. The real part projects under the moment map of $\mathbb{C}\mathbb{P}^2$ onto the entire moment polytope, each point in the interior of the polytope having four preimages, the points on the interior of the edges on the boundary of the triangle having two preimages, the points on the vertices having one preimage. By taking four copies of the moment polytope and gluing them along the edges according to the prescriptions of the torus action on $\mathbb{C}\mathbb{P}^2$ (see [1]), one recovers the real projective plane.

From the Lagrangian submanifold point of view, $L = \mathbb{R}\mathbb{P}^2$ is an important example of monotone Lagrangian submanifold in $\mathbb{C}\mathbb{P}^2$. Monotone means that there exists a positive constant K_L such that

$$\forall u \in H_2(\mathbb{C}\mathbb{P}^2, L), \int_u \omega = K_L \mu_L(u),$$

where $\mu_L : H_2(\mathbb{C}\mathbb{P}^2, L) \rightarrow \mathbb{Z}$ is the Maslov class of L .

The exotic torus of Chekanov and Schlenk (see [7]) is another important example of monotone Lagrangian submanifold of $\mathbb{C}\mathbb{P}^2$. The second author proved in [8] that this torus is Hamiltonian isotopic to a torus described by

MA was partially funded by FCT/Portugal through projects UID/MAT/04459/2013 and EXCL/MAT-GEO/0222/2012.

AG was partially supported by CMUP (UID/MAT/00144/2013), which is funded by FCT (Portugal) with national (MEC) and European structural funds through the programs FEDER, under the partnership agreement PT2020.

MA and AG were both partially funded through the project PTDC/MAT/117762/2010. The present work is part of the authors' activities within CAST, a Research Networking Program of the European Science Foundation.

Biran and Cornea in [3]. To do so, she Hamiltonian-isotoped both tori to a so-called modified Chekanov torus $\tilde{\Theta}_{\text{Ch}}$. This torus has a nice image under the moment map and can be reconstructed, as the real projective space, out of copies of this image and gluing patterns. The rules for gluing are of two types. The first are coming from the definition of the moment map and are the same as the ones used for the real part. The second are new and we have managed to interpret them as Lagrangian surgeries of two copies of the real part intersecting transversely at one isolated point and cleanly (in the sense of Pozniak [13]) along a circle.

The surgery for two Lagrangian submanifolds intersecting transversely at a point has been developed by Polterovich in [12] and we have modified it to keep a toric description of the result of the surgery. The surgery for two Lagrangian submanifolds intersecting along an isotropic submanifold not reduced to a point is new and we intend to develop it in full generality in a future work. We show in our case:

Theorem 1. *The Chekanov–Schlenk torus is Hamiltonian isotopic in $\mathbb{C}\mathbb{P}^2$ to a Lagrangian torus obtained from two copies of $\mathbb{R}\mathbb{P}^2$ by Lagrangian surgeries at a point and along an isotropic circle.*

With our method we can also recover Lagrangian embeddings of some surfaces in \mathbb{R}^4 that were constructed by Givental in [9] and that we then embed in $\mathbb{C}\mathbb{P}^2$ or $\mathbb{C}\mathbb{P}^1 \times \mathbb{C}\mathbb{P}^1$. The advantage of this construction is that we have a good control of the monotonicity condition. We knew so far only the monotone Lagrangian embeddings of tori in $\mathbb{C}\mathbb{P}^1 \times \mathbb{C}\mathbb{P}^1$ and of tori and real projective planes in $\mathbb{C}\mathbb{P}^2$. Our method enables us to prove

Theorem 2. *There exists a monotone Lagrangian embedding of the connected sum of a surface of genus 2 and a Klein bottle in $\mathbb{C}\mathbb{P}^2$ and a monotone Lagrangian embedding of the connected sum of a surface of genus 4 and a Klein bottle in the product $\mathbb{C}\mathbb{P}^1 \times \mathbb{C}\mathbb{P}^1$.*

Note that neither the Klein bottle (see [14]) nor the orientable surface of genus 2 (see [10]) can be embedded as Lagrangian submanifolds of $\mathbb{C}\mathbb{P}^2$.

The structure of this article is as follows: in Section 2 we study the gluing patterns for the modified Chekanov torus and we describe the surgeries we will use; in Section 3 we give our main construction and study the monotonicity of examples we can get with the surgery at a point; finally, in Section 4, we use the surgery for an intersection along a circle to describe two monotone Lagrangian embeddings.

Acknowledgements. *We thank Leonardo Macarini for useful conversations that, in particular, led to the identification of a mistake in an earlier draft of this paper. We also thank an anonymous referee for a very careful reading and detailed list of corrections.*

2. THE LOCAL MODELS

2.1. A toric model of the exotic torus of Chekanov and Schlenk in $\mathbb{C}\mathbb{P}^2$. There is a well-known monotone torus in $\mathbb{C}\mathbb{P}^2$ called the Clifford torus which can be described in homogeneous coordinates as

$$T_{\text{Cliff}} = \left\{ \left[e^{i\alpha} : e^{i\beta} : 1 \right] \mid \alpha, \beta \in [0, 2\pi] \right\}.$$

Its image under the moment map of $\mathbb{C}\mathbb{P}^2$ (corresponding to the normalization of the symplectic form we use in Section 3)

$$\begin{aligned} \mu : \quad \mathbb{C}\mathbb{P}^2 &\longrightarrow \mathbb{R}^2 \\ [z_0 : z_1 : z_2] &\longmapsto \left(3 \frac{|z_0|^2}{\sum |z_i|^2}, 3 \frac{|z_1|^2}{\sum |z_i|^2} \right), \end{aligned}$$

is the barycenter $(1, 1)$ of the image of $\mathbb{C}\mathbb{P}^2$, the triangle obtained as the convex hull of the points $(0, 0)$, $(3, 0)$, $(0, 3)$.

In 2004, Chekanov and Schlenk (see [7]) have studied the torus given in homogeneous coordinates of $\mathbb{C}\mathbb{P}^2$:

$$\Theta_{\text{CS}} = \left\{ \left[\frac{1}{\sqrt{2}} \gamma(s) e^{i\theta} : \frac{1}{\sqrt{2}} \gamma(s) e^{-i\theta} : \sqrt{\frac{3}{\pi} - |\gamma(s)|^2} \right] \mid \theta, s \in [0, 2\pi] \right\}$$

where $\gamma : [0, 2\pi] \rightarrow \mathbb{C}$ parametrizes a curve enclosing a domain of area 1 lying in the disk centered in the origin and of area $2 + \varepsilon$ of \mathbb{C} , in the half-disk of complex numbers of positive real part (see Figure 1). They have proved (see [7, 5, 6]) that Θ_{CS} is a monotone Lagrangian torus in $\mathbb{C}\mathbb{P}^2$, non-displaceable and non-Hamiltonian isotopic to the Clifford torus (therefore called exotic) in $\mathbb{C}\mathbb{P}^2$.

By its definition, the Chekanov–Schlenk torus projects under the moment map μ to a segment lying in the diagonal line of \mathbb{R}^2 . More precisely the image is

$$\left\{ (x, x) \in \mathbb{R}^2 \mid x \in \left[\frac{\pi}{2} \rho_{\min}^2, \frac{\pi}{2} \rho_{\max}^2 \right] \right\}$$

where ρ_{\min} is the minimum of $|\gamma(s)|$ and ρ_{\max} is the maximum.

There is a description of this exotic torus more adapted to the toric picture, that enables to reconstruct the torus from its moment map image as in the case of the real part of $\mathbb{C}\mathbb{P}^2$.

Such a description can be obtained by considering the modified Chekanov torus of [8]. This torus is a torus Hamiltonian isotopic to the exotic torus of Chekanov–Schlenk and is defined in homogeneous coordinates (with the normalizations of [7]) by

$$\tilde{\Theta}_{\text{Ch}} = \left\{ \left[\cos(\theta) \gamma(s) : \sin(\theta) \gamma(s) : \sqrt{\frac{3}{\pi} - |\gamma(s)|^2} \right] \mid \theta, s \in [0, 2\pi] \right\}.$$

The image of the torus under the moment map μ can be parametrised by

$$\mu(\tilde{\Theta}_{\text{Ch}}) = \left\{ (\pi \cos^2(\theta) |\gamma(s)|^2, \pi \sin^2(\theta) |\gamma(s)|^2) \mid \theta, s \in [0, 2\pi] \right\}.$$

It is a trapezoid sitting inside the polytope of $\mathbb{C}\mathbb{P}^2$ between the two parallel lines $x + y = \pi\rho_{\min}^2$ and $x + y = \pi\rho_{\max}^2$.

If the curve γ is such that $\gamma(0) = \rho_{\min}$, $\gamma(\pi) = \rho_{\max}$, γ is symmetric with respect to the real axis, and each point $|\gamma(s)| = \rho(s)$ has only one preimage $s \in (0, \pi)$ (see for example Figure 1), then for $s \neq 0, \pi$ and $\theta \neq 0, \pi$, the point

$$(1) \quad (\pi \cos^2(\theta)|\gamma(s)|^2, \pi \sin^2(\theta)|\gamma(s)|^2)$$

has 8 preimages in the torus.

Recall (see for example [2, 4]) that the image of the moment map for $\mathbb{C}\mathbb{P}^2$, or for a general (compact, connected) toric manifold (M, ω) is a convex polytope P such that the fiber of each point of P is an isotropic torus. Recall also that we have action-angle coordinates on the preimage \mathring{M} of the interior \mathring{P} which is the open dense set in M consisting of all the points where the action of the torus \mathbb{T}^n is free. One can describe this set as

$$\mathring{M} \cong \mathring{P} \times \mathbb{T}^n = \left\{ (x_1, \dots, x_n, e^{i\theta_1}, \dots, e^{i\theta_n}) \mid x \in \mathring{P}, \theta \in \mathbb{R}^n / 2\pi\mathbb{Z}^n \right\},$$

where (x, θ) are the action-angle coordinates for the symplectic form

$$\omega = \sum dx_j \wedge d\theta_j.$$

In the case of $\tilde{\Theta}_{\text{Ch}}$, we parametrise the curve γ above the real axis by

$$\gamma(s) = \rho(s)e^{it(s)},$$

such that $t(s) \in [0, t_{\max}]$, $t_{\max} < \frac{\pi}{2}$, $t(0) = 0$, $t(\pi) = 0$.




FIGURE 1. A suitable curve γ

Then for a fixed s in $(0, \pi)$ and a fixed θ in $(0, \frac{\pi}{2})$, the eight preimages of the point (1) in $\mu(\tilde{\Theta}_{\text{Ch}})$ are given in action-angle coordinates by:

$$A_{\epsilon,k,\ell} = \left(\pi \cos^2(\theta)|\gamma(s)|^2, \pi \sin^2(\theta)|\gamma(s)|^2, \epsilon t(s) + \frac{\pi}{2} + k\pi, \epsilon t(s) + \frac{\pi}{2} + \ell\pi \right)$$

with $\epsilon \in \{-1, 1\}$, $k, \ell \in \{0, 1\}$.

When s goes to 0 or π , $t(s)$ goes to 0 and the points in the torus fiber converge (moving along the diagonal direction) towards one of the four points

$$\left(\frac{\pi}{2} + k\pi, \frac{\pi}{2} + \ell\pi \right), k, \ell \in \{0, 1\},$$

see Figure 2.




FIGURE 2. The points are at the four corners of each of the dashed squares; when s goes to 0, the corners of the two squares are identified

This describes the gluing of the trapezoid along the segments $x+y = \pi\rho_{\min}^2$ and $x+y = \pi\rho_{\max}^2$.

2.2. The interpretation of the gluing along the segment $x+y = \pi\rho_{\min}^2$. The gluing along the segment $x+y = \pi\rho_{\min}^2$ can be described as the Lagrangian surgery defined in [12] of two Hamiltonian isotopic copies of the real part $\mathbb{R}\mathbb{P}^2$ intersecting transversally at the origin $[0 : 0 : 1]$ (see Section 4.2 for the details).

Let us describe the Lagrangian surgery we will use in the rest of this article which is a slight modification of [12]. Following Polterovich, one does the surgery of two transverse Lagrangians in a local chart around an intersection point. In this chart, one finds an almost-complex structure j such that locally $l_1 = jl_0$ and the local Lagrangian handle is the image of the sphere in l_0 times $[-T, T]$, for T large under the map $(\xi, t) \mapsto e^{-t}\xi + e^t j\xi$. One then connects the handle on its boundaries to the original Lagrangians by some smoothing.

Because we aim to control the monotonicity condition and keep the standard local chart in $\mathbb{C}\mathbb{P}^2$, we explicitly describe the Lagrangian surgery we will be using, without the use of an auxiliary almost-complex structure j in \mathbb{C}^2 .

The handle between two Lagrangian linear subspaces of \mathbb{C}^2 . Consider the linear \mathbb{C}^2 and the two Lagrangian linear subspaces

$$l_0 = \mathbb{R}^2$$

and

$$l_1 = \begin{pmatrix} e^{i\alpha} & 0 \\ 0 & e^{i\beta} \end{pmatrix} \mathbb{R}^2$$

the image of l_0 by the Hamiltonian diffeomorphism defined by the diagonal matrix $\text{diag}(e^{i\alpha}, e^{i\beta})$ for α and β not a multiple of π .

We define a handle h parametrised by:

$$h = \left\{ e^{-t} \begin{pmatrix} x_0 \\ x_1 \end{pmatrix} + e^t \begin{pmatrix} e^{i\alpha} x_0 \\ e^{i\beta} x_1 \end{pmatrix} \mid \begin{array}{l} t \in \mathbb{R} \\ x_0^2 + x_1^2 = 1 \end{array} \right\}.$$

It is asymptotic to l_0 when t goes to $-\infty$ and to l_1 when t goes to $+\infty$. One checks that it is a Lagrangian handle when $\sin(\alpha) = \sin(\beta)$.

Note that for the same reason, the handle

$$h' = \left\{ e^{-t} \begin{pmatrix} x_0 \\ x_1 \end{pmatrix} + e^t \begin{pmatrix} e^{i\alpha}(-x_0) \\ e^{i\beta}(-x_1) \end{pmatrix} \mid \begin{array}{l} t \in \mathbb{R} \\ x_0^2 + x_1^2 = 1 \end{array} \right\}$$

is also a Lagrangian submanifold asymptotic to l_0 and l_1 and corresponds to the first handle for the angles $(\alpha + \pi, \beta + \pi)$.

Note that when $\sin(\alpha) = -\sin(\beta)$, one can also define two Lagrangian handles parametrized by

$$h = \left\{ e^{-t} \begin{pmatrix} x_0 \\ x_1 \end{pmatrix} + e^t \begin{pmatrix} e^{i\alpha} x_0 \\ e^{i\beta} (-x_1) \end{pmatrix} \mid \begin{array}{l} t \in \mathbb{R} \\ x_0^2 + x_1^2 = 1 \end{array} \right\}$$

and

$$h' = \left\{ e^{-t} \begin{pmatrix} x_0 \\ x_1 \end{pmatrix} + e^t \begin{pmatrix} e^{i\alpha} (-x_0) \\ e^{i\beta} x_1 \end{pmatrix} \mid \begin{array}{l} t \in \mathbb{R} \\ x_0^2 + x_1^2 = 1 \end{array} \right\}.$$

The smoothing. For some (large) T , denote by h_T the image of the handle for t between $-T$ and T :

$$h_T = \left\{ e^{-t} \begin{pmatrix} x_0 \\ x_1 \end{pmatrix} + e^t \begin{pmatrix} e^{i\alpha} x_0 \\ e^{i\beta} x_1 \end{pmatrix} \mid \begin{array}{l} t \in [-T, T] \\ x_0^2 + x_1^2 = 1 \end{array} \right\}.$$

Fix a parameter T large. We smooth the handle at the ends of h_T as in [12]. Notice that the original surgery of Polterovich corresponds to the case when $\alpha = \beta = \frac{\pi}{2}$ and we can obtain any of our handles from Polterovich's surgery by applying the linear transformation of $\mathbb{C}^2 = \mathbb{R}^4$ with matrix

$$\begin{pmatrix} 1 & \cos(\alpha) & 0 & 0 \\ 0 & \sin(\alpha) & 0 & 0 \\ 0 & 0 & 1 & \cos(\beta) \\ 0 & 0 & 0 & \sin(\beta) \end{pmatrix}.$$

Hence we can smooth the handle at the boundary of h_T when $\alpha = \beta = \frac{\pi}{2}$ as in [12] and then take the image of this smoothing by the linear map above to get a smoothing in our case.

Note that this surgery lies inside a big ball of radius R , and outside it, the Lagrangian submanifold obtained is the union of l_0 and l_1 . As the linear Lagrangians are homogenous with respect to homotheties centered at the origin of \mathbb{C}^2 , one can use a conformal transformation to make this Lagrangian surgery happen in a ball B_0 of small radius. Equivalently, one can take the handle to be, not the image of the unit sphere in l_0 , but of a smaller one

$$x_0^2 + x_1^2 = \varepsilon_1^2$$

and the smoothing happening outside a ball of radius ε_2 such that the Lagrangian submanifold after surgery identifies with the union of l_0 and l_1 outside a ball of radius ε for a parameter $\varepsilon > \varepsilon_2 > \varepsilon_1 > 0$ small.

Controlling the area. Let us study the restriction of this surgery (before conformal transformation) to one factor \mathbb{C} of \mathbb{C}^2 , for instance the first. The trace of the surgery along this coordinate is given by:

$$h_T^1 = \{ e^{-t} + e^t e^{i\alpha} \mid t \in [-T, T] \}$$

for some large T , followed by some small smoothing between its ends and the original l_0 and l_1 , together with the symmetric curve about the origin.

Let us compute the area between the original l_0 and l_1 and one of the arcs of the surgery, namely the area in grey in Figure 3.




FIGURE 3. The intersection of the handle with the first \mathbb{C} -factor of \mathbb{C}^2

As before, we can deduce this area from the computation of the area when $\alpha = \frac{\pi}{2}$. In this case, h_T^1 is the arc of hyperbola in the plane given by the equation $xy = 1$ so that the area in grey is equal to $2T + 1$. This area is equivalent to $2T$ when T goes to $+\infty$. Moreover, when T is large, the smoothing between the original linear Lagrangians and the arc of hyperbola is very small so that the area is still equivalent to $2T$.

The other cases can be obtained from the $\alpha = \frac{\pi}{2}$ situation by applying the linear transformation of the plane $\mathbb{C} = \mathbb{R}^2$ given by the matrix

$$\begin{pmatrix} 1 & \cos(\alpha) \\ 0 & \sin(\alpha) \end{pmatrix},$$

so that the area considered above is equivalent to $2\sin(\alpha)T$ when T goes to ∞ .

In particular, when $\sin(\alpha) = \sin(\beta)$, the condition which ensures the surgery to be Lagrangian, the areas between the original Lagrangians and the handle along each coordinate are equivalently the same. For T large enough, we can (and will) do the smoothing so that the areas along each \mathbb{C} -factor are equal.

Now, given the conformality property of the surgery, we can make this area as small as we want and equal to some $a(\varepsilon)$ (small) if the surgery is done inside the ball of radius ε .

In a general symplectic manifold. Let L_0 and L_1 be two Lagrangian submanifolds of a symplectic manifold W intersecting transversally at a point x_0 . One can take a Darboux chart U_0 around x_0 symplectomorphic to a ball B_0 endowed with the standard symplectic form of \mathbb{C}^2 such that under the Darboux map, the two Lagrangians are the intersection of Lagrangian linear subspaces of \mathbb{C}^2 with the ball. One is then in the linear situation from above

and can perform the surgery in the ball as described provided the sines of angles between the restriction of the linear Lagrangians to each factor of \mathbb{C}^2 are the same (up to sign).

2.3. The interpretation of the gluing along the segment $x + y = \pi\rho_{\max}^2$. As we shall see in Section 4.2, the gluing along the segment $x + y = \pi\rho_{\max}^2$ can be interpreted as the Lagrangian surgery along a circle of two Hamiltonian isotopic copies of the real part $\mathbb{R}\mathbb{P}^2$ intersecting along a circle in the $\mathbb{C}\mathbb{P}^1$ at infinity

$$\{[z_0 : z_1 : z_2] \mid z_2 = 0\}.$$

This circle is isotropic and, as in the case of the surgery at a point, the surgery along an isotropic submanifold is a local process that we will now describe.

The neighbourhood of an isotropic manifold. Let P be a symplectic manifold of dimension $2n$. Let N be an isotropic submanifold of dimension k . In [15], Weinstein noticed that the tangent bundle of P along N is isomorphic as symplectic vector bundle over N to

$$(TN \oplus TN^*) \oplus SN(N, P),$$

where $SN(N, P) = TN^\perp/TN$ is called the symplectic normal bundle of N in P .

Conversely, one can embed any manifold which is the base of a symplectic vector bundle as an isotropic submanifold of a symplectic manifold such that the tangent bundle looks like this:

Theorem 2.1 (The existence theorem, Weinstein [16]). *Let N be a manifold of dimension k and $E \rightarrow N$ a symplectic vector bundle with fibre dimension $2(n - k)$, $k \leq n$. Then N can be embedded as an isotropic submanifold of a symplectic manifold $P(E)$ of dimension $2n$ such that the tangent bundle of $P(E)$ along N is isomorphic as symplectic vector bundle to the sum $(TN \oplus TN^*) \oplus E$.*

This space $P(E)$ is the Whitney sum $P(E) = T^*N \oplus E$ as Weinstein explains in [15]. The symplectic structure on $P(E)$ is not canonical and is described in [16].

And we have a uniqueness result:

Theorem 2.2 (Weinstein [15]). *The isotropic manifold theorem: Let N be a manifold of dimension k . Then the extensions of N to a $2n$ -dimensional symplectic manifold in which N is isotropic are classified, up to local symplectomorphism about N , by the isomorphism classes of $2(n - k)$ -dimensional symplectic vector bundles over N .*

This means that if N is an isotropic submanifold of a symplectic manifold P , then a neighbourhood of N in P is symplectomorphic to a neighbourhood of the embedding of N in $P(E)$ for $E = SN(N, P)$.

We will extend the surgery at an intersection point of two Lagrangian submanifolds to the surgery of two Lagrangian submanifolds intersecting cleanly (in the sense of Pozniak [13]) along an isotropic submanifold. In his thesis, Pozniak proves

Theorem 2.3 (Pozniak [13]). *If two Lagrangian submanifolds L_0 and L_1 of a symplectic manifold P intersect cleanly along N , that is if $N = L_0 \cap L_1$ and for each $x \in N$, $T_x N = T_x L_0 \cap T_x L_1$, then there exists a vector bundle $L \rightarrow N$ such that a neighbourhood of N in P is symplectomorphic to a neighbourhood of N in T^*L , L_0 being mapped to the zero section of T^*L and L_1 to the conormal of N in T^*L .*

In this setting, identifying L_0 and L_1 with their image in T^*L , one can see that $E = SN(N, T^*L)$ is isomorphic to the Whitney sum of the vector bundles $L \rightarrow N$ and $L^* \rightarrow N$ and that in the Whitney sum $P(E) = T^*N \oplus E$, the Lagrangian L_0 is mapped to the zero section in the T^*N -summand and to $L \oplus \{0\}$ in the E -summand and the Lagrangian L_1 is mapped to the zero section in the T^*N -summand and to $\{0\} \oplus L^*$ in the E -summand, so that the intersection of L_0 and L_1 is the sum of the zero-section of T^*N and the transverse intersection in each fibre of E of $L_x \oplus \{0\}$ with $\{0\} \oplus L_x^*$.

The surgery we will construct in this neighbourhood will also fiber over N , be equal to the zero-section in the T^*N -summand and will resolve the intersection in each fiber of E , so that it is enough to define it in the symplectic normal bundle E .

The bundle surgery. In this paper, the constructions are done only in real dimension 4 with the clean intersection of two Lagrangians along an isotropic circle. The symplectic normal bundle $E = SN(N, T^*L)$ is then a rank 2 symplectic vector bundle. There exists only one rank 2 symplectic vector bundle over the circle, the trivial bundle $\mathbb{S}^1 \times \mathbb{C}$. However, the rank 1 Lagrangian subbundle $L \rightarrow N$ can be the trivial line bundle or the non-orientable line bundle over the circle. So in real dimension 4 one will be in one of the following case:

- either E is $\mathbb{S}^1 \times \mathbb{C}$ and $L \rightarrow N$ is $\mathbb{S}^1 \times \mathbb{R}$, the trivial real subbundle;
- or E can be described as $[0, 1] \times \mathbb{C}$ identifying the fiber at 0 and the fiber at 1 by multiplication by -1 and $L \rightarrow N$ is the associated subbundle $[0, 1] \times \mathbb{R}$ with the same identification.

Now if the restriction of the Lagrangian L_1 to E is the associated subbundle with fiber $e^{i\alpha}\mathbb{R}$ as it will be the case in our examples, one can perform a surgery in dimension 1 parametrized in each fiber as

$$\{e^{-t}x + e^t e^{i\alpha}x \mid t \in [-T, T], x \in \mathbb{R}, x^2 = \varepsilon_1^2\}$$

or

$$\{e^{-t}x + e^t e^{i\alpha}(-x) \mid t \in [-T, T], x \in \mathbb{R}, x^2 = \varepsilon_1^2\},$$

followed by a smoothing at the end.

Now note that in this situation, the restrictions of L_0 and L_1 are invariant by multiplication by -1 , so the handles and the smoothing can be made invariant as well. Therefore, in both cases, the trivial and the non-trivial symplectic bundle over \mathbb{S}^1 , the change of trivialisation preserves the construction so that the handle can be defined globally and fibers over \mathbb{S}^1 .

3. CONSTRUCTION OF NEW MONOTONE LAGRANGIAN SUBMANIFOLDS USING THE SURGERY AT A POINT

3.1. A non-orientable monotone Lagrangian in $\mathbb{C}\mathbb{P}^2$. In the following sections, we explain how to get via a Lagrangian surgery on two copies of $\mathbb{R}\mathbb{P}^2$ a monotone Lagrangian connected sum of a Klein bottle and an orientable surface of genus two in $\mathbb{C}\mathbb{P}^2$.

3.1.1. The construction. We will take two copies of $\mathbb{R}\mathbb{P}^2$ in $\mathbb{C}\mathbb{P}^2$ that intersect in three points exactly, the three points of $\mathbb{C}\mathbb{P}^2$ projecting on the three corners of the image of the moment map. Let us consider:

$$L_0 = \{[x_0 : x_1 : x_2] \in \mathbb{C}\mathbb{P}^2 \mid x_0, x_1, x_2 \in \mathbb{R}\}$$

and

$$L_1 = \{[e^{i\frac{\pi}{3}}x_0 : e^{-i\frac{\pi}{3}}x_1 : x_2] \in \mathbb{C}\mathbb{P}^2 \mid x_0, x_1, x_2 \in \mathbb{R}\}.$$

These are two copies of $\mathbb{R}\mathbb{P}^2$ in $\mathbb{C}\mathbb{P}^2$, L_1 being obtained from L_0 by a Hamiltonian isotopy. Indeed, L_1 is the image of L_0 under the map given by the action of the following diagonal matrix on the two first coordinates of the homogeneous coordinates:

$$A = \begin{pmatrix} e^{i\frac{\pi}{3}} & 0 \\ 0 & e^{-i\frac{\pi}{3}} \end{pmatrix}.$$

It is the time-one map of the transformation given by

$$A_t = \begin{pmatrix} e^{it\frac{\pi}{3}} & 0 \\ 0 & e^{-it\frac{\pi}{3}} \end{pmatrix}$$

$A_t \in SU(2)$ and it defines a Hamiltonian diffeomorphism Φ_t of $\mathbb{C}\mathbb{P}^2$.

One can check that these two copies of $\mathbb{R}\mathbb{P}^2$ intersect in the three points $[0 : 0 : 1]$, $[0 : 1 : 0]$, $[1 : 0 : 0]$.

We want to perform a Lagrangian surgery at each intersection point as described in Section 2.2. Let us give the choices of handles we make for the construction.

At $[0 : 0 : 1]$, the local chart is

$$[z_0 : z_1 : z_2] \mapsto \left(\frac{z_0}{z_2}, \frac{z_1}{z_2} \right),$$

so that locally, L_0 is the real plane

$$l_0 = \{(x_0, x_1) \mid x_0, x_1 \in \mathbb{R}\}$$

and L_1 is

$$l_1 = \{(e^{i\frac{\pi}{3}}x_0, e^{-i\frac{\pi}{3}}x_1) \mid x_0, x_1 \in \mathbb{R}\}.$$

We are in the case when $\sin(\frac{\pi}{3}) = -\sin(-\frac{\pi}{3})$, so that we need the modified version of the handle to do the Lagrangian surgery. We will use the one defined by the smoothing of:

$$\left\{ e^{-t} \begin{pmatrix} x_0 \\ x_1 \end{pmatrix} + e^t \begin{pmatrix} e^{i\frac{\pi}{3}}x_0 \\ e^{-i\frac{\pi}{3}}(-x_1) \end{pmatrix} \mid \begin{array}{l} t \in \mathbb{R} \\ x_0^2 + x_1^2 = \varepsilon_1^2 \end{array} \right\}.$$

At $[0 : 1 : 0]$, the local chart is

$$[z_0 : z_1 : z_2] \mapsto \left(\frac{z_0}{z_1}, \frac{z_2}{z_1} \right),$$

so that locally, L_0 is the real plane

$$l_0 = \{(x_0, x_2) \mid x_0, x_2 \in \mathbb{R}\}$$

and L_1 is

$$l_1 = \{(e^{i\frac{2\pi}{3}}x_0, e^{i\frac{\pi}{3}}x_2) \mid x_0, x_2 \in \mathbb{R}\}.$$

As $\sin(2\pi/3) = \sin(\pi/3)$, we can use the first description of the handle to define the Lagrangian surgery:

$$\left\{ e^{-t} \begin{pmatrix} x_0 \\ x_2 \end{pmatrix} + e^t \begin{pmatrix} e^{i\frac{2\pi}{3}}x_0 \\ e^{i\frac{\pi}{3}}x_2 \end{pmatrix} \mid \begin{array}{l} t \in \mathbb{R} \\ x_0^2 + x_2^2 = \varepsilon_1^2 \end{array} \right\}.$$

At $[1 : 0 : 0]$, the local chart is

$$[z_0 : z_1 : z_2] \mapsto \left(\frac{z_1}{z_0}, \frac{z_2}{z_0} \right),$$

so that locally, L_0 is the real plane

$$l_0 = \{(x_1, x_2) \mid x_1, x_2 \in \mathbb{R}\}$$

and L_1 is

$$l_1 = \{(e^{-i\frac{2\pi}{3}}x_1, e^{-i\frac{\pi}{3}}x_2) \mid x_1, x_2 \in \mathbb{R}\}.$$

As $\sin(-2\pi/3) = \sin(-\pi/3)$, we can also use the first description of the handle to do the Lagrangian surgery. But for the monotonicity condition to be satisfied in Section 3.1.2, we will use instead the smoothing of the h' -handle:

$$\left\{ e^{-t} \begin{pmatrix} x_1 \\ x_2 \end{pmatrix} + e^t \begin{pmatrix} e^{-i\frac{2\pi}{3}}(-x_1) \\ e^{-i\frac{\pi}{3}}(-x_2) \end{pmatrix} \mid \begin{array}{l} t \in \mathbb{R} \\ x_1^2 + x_2^2 = \varepsilon_1^2 \end{array} \right\}.$$

After these surgeries, the projection of the Lagrangian L we constructed will be contained in the polytope obtained from the polytope of $\mathbb{C}\mathbb{P}^2$ by cutting the three vertices as in Figure 4.

The choice of these copies of $\mathbb{R}\mathbb{P}^2$ and these surgeries is motivated by the monotonicity condition we aim to prove for this construction in the next section. Let us describe the restriction of the surgery along the coordinate- $\mathbb{C}\mathbb{P}^1$ s, that is the projective lines which are the preimages of the edges on




FIGURE 4. The image of L under the moment map is contained in and smoothly approximates the shaded polytope.

the boundary of the moment polytope and can be defined by the vanishing of one of the homogeneous coordinates. We will describe the case of

$$\{[z_0 : z_1 : z_2] \mid z_0 = 0\},$$

the other coordinate- $\mathbb{C}\mathbb{P}^1$ s being similar.

Along the sphere $z_0 = 0$, L_0 and L_1 are two circles intersecting transversally at the north and the south pole. Locally in the chart \mathbb{C} at $[0 : 0 : 1]$, we have L_0 on the real axis and L_1 on the axis $e^{-i\frac{\pi}{3}}\mathbb{R}$. In this chart, the intersection of the chosen surgery with $\{z_0 = 0\}$ consists in two curves (see Figure 5-left): inside a ball centered at the origin and of area ε they lie in two opposite "quadrants" defined by these two axes, that is in the quadrants making an angle of $2\frac{\pi}{3}$.




FIGURE 5. The surgery along $z_0 = 0$ in the chart at $[0 : 0 : 1]$ (on the left) and in the chart at $[0 : 1 : 0]$ (right)

Locally at $[0 : 1 : 0]$ we have a similar picture but with curves in the quadrants making an angle of $\frac{\pi}{3}$ (see Figure 5-right).

Away from the small neighbourhoods where we do the surgery, namely on the part where we glue the two charts, the restriction of L is the restriction of L_0 and L_1 to this complex projective line. One sees then that the restriction of L to this $\mathbb{C}\mathbb{P}^1$ is one circle joining L_0 and L_1 through the two handles constructed at each intersection point and looking like the seam of a tennis ball (see Figure 6).




FIGURE 6. The intersection of the surgery with $\{z_0 = 0\}$

If we had chosen the first handle we described in Section 2.2 at $[0 : 0 : 1]$, the restriction of L to $z_0 = 0$ would have been the union of two circles.

Actually, the choice of handles we made is such that in each of the other coordinate- $\mathbb{C}\mathbb{P}^1$ s (namely $z_1 = 0$ and $z_2 = 0$), the restriction of L is one circle joining L_0 and L_1 through the two handles constructed at each intersection point at the poles of $\mathbb{C}\mathbb{P}^1$. Indeed, in the case of two circles in the intersection with a coordinate- $\mathbb{C}\mathbb{P}^1$, one cannot expect to satisfy the monotonicity condition. Note however that as the two intersecting Lagrangians L_0 and L_1 are not orientable, it follows from [12, Proposition 2] that the topology of the Lagrangian we obtain after surgeries does not depend on the choice of handles.

3.1.2. *Monotonicity.* Let us normalize the symplectic form on $\mathbb{C}\mathbb{P}^2$ such that the area of a projective line is 3:

$$\int_{\mathbb{C}\mathbb{P}^1} \omega = 3.$$

With this normalization, $\mathbb{C}\mathbb{P}^2$ is monotone with monotonicity constant 1:

$$\forall v \in H_2(\mathbb{C}\mathbb{P}^2), \int_v \omega = c_1(T\mathbb{C}\mathbb{P}^2)(v)$$

and any monotone Lagrangian submanifold L will have monotonicity constant equal to half the monotonicity constant of $\mathbb{C}\mathbb{P}^2$ (see [11]), namely $\frac{1}{2}$:

$$\forall u \in H_2(\mathbb{C}\mathbb{P}^2, L), \int_u \omega = \frac{1}{2} \mu_L(u),$$

where μ_L is the Maslov class of L .

Theorem 3.1. *The construction of Section 3.1.1 produces a monotone Lagrangian embedding of the connected sum of a Klein bottle and a compact orientable surface of genus 2 in $\mathbb{C}\mathbb{P}^2$.*

Proof. Topologically, the surgery at a point between two copies of the real projective plane gives the connected sum of these two spaces, namely a Klein bottle. Then attaching a 2-dimensional handle corresponds to a connected sum with a torus, so that the Lagrangian submanifold constructed in 3.1.1 is diffeomorphic to the connected sum

$$L \cong \mathbb{R}\mathbb{P}^2 \# \mathbb{R}\mathbb{P}^2 \# \mathbb{T}^2 \# \mathbb{T}^2 \cong K \# \Sigma_2$$

where K is a Klein bottle and Σ_2 is a compact orientable surface of genus 2.

We know that $H_2(\mathbb{C}\mathbb{P}^2, L) \cong H_2(\mathbb{C}\mathbb{P}^2) \oplus H_1(L)$ as $H_2(L) = 0$. Let us examine the monotonicity condition on each factor of this direct sum.

On the factor $H_2(\mathbb{C}\mathbb{P}^2)$, we already have the monotonicity condition from the one on $\mathbb{C}\mathbb{P}^2$ so that we only need to verify the monotonicity condition on the disks representing generators of $H_1(L)$. This means also that for a given generator of $H_1(L)$, it is enough to satisfy the monotonicity condition for one choice of disk with boundary this generator, as the symplectic invariants for another disc with the same boundary will differ by the invariants coming from a sphere (the sphere obtained by gluing the two disks along their boundary) where the condition is already verified.

Now $H_1(L) = \mathbb{Z}^5 \oplus \mathbb{Z}/2$ can be generated by the two loops generating the first homology group of each copy of $\mathbb{R}\mathbb{P}^2$, the three loops inside each handle generating the homology of the handle and three loops "between" the handles (see Figure 7).

Any loop sitting in one of the original copies of $\mathbb{R}\mathbb{P}^2$, L_0 or L_1 , satisfies the monotonicity condition because L_0 and L_1 are monotone.

As a representative of a generator of the homology of the handle, one can take the circle on one of the extremities of the handle lying on one of the copies of the Lagrangian $\mathbb{R}\mathbb{P}^2$, say for example L_0 . More explicitly, one may take the image of the circle $\{(x_0, x_1), x_0^2 + x_1^2 = \varepsilon^2\}$, for $\varepsilon > \varepsilon_1$ small, in the chart around an intersection point of L_0 and L_1 . We take a disc in $\mathbb{C}\mathbb{P}^2$ with boundary this circle in L and compute the symplectic invariants of that disc. One can for example choose the disc in L_0 which was cut out from L_0 to build L . But as the disc is Lagrangian, the two invariants, area and Maslov class, vanish on this disc.

One is left with checking the monotonicity condition on the circles between handles. One can prove that the monotonicity condition is satisfied




FIGURE 7. Three loops between the handles

on the three circles drawing the tennis ball seam on the coordinate- \mathbb{CP}^1 s we described at the end of the previous section and the discs they bound on these \mathbb{CP}^1 . Unfortunately these circles are not in our set of generators for $H_1(L)$, since their homology class is 2 times the loop between the corresponding handles depicted in Figure 7. But we can use for the generators loops which partially follow these seams. Let us describe a loop γ we can choose between the handles created at $[0 : 0 : 1]$ and $[0 : 1 : 0]$ and a disk it bounds. Two other loops between handles can be constructed in a similar way.


The loop γ is almost entirely lying in the coordinate \mathbb{CP}^1 of homogeneous equation $z_0 = 0$. See Figure 8.

It is based at a point where L coincides with L_0 , for example the point a of local coordinates $(z_0, z_1) = (0, \varepsilon)$ in the local chart at $[0 : 0 : 1]$. From this point, follow the handle at $[0 : 0 : 1]$ along the path parametrized by

$$\{(0, e^{-t}\varepsilon_1 + e^t e^{-i\frac{\pi}{3}}(-\varepsilon_1))\}$$

(we include here the smoothing by considering we can locally take the parametrization of the handle for t varying from $-\infty$ to $+\infty$) till the point b of local coordinates $(z_0, z_2) = (0, e^{-i\frac{\pi}{3}}\varepsilon)$. Then follow $L_1 \cap \{z_0 = 0\}$ "up" towards $[0 : 1 : 0]$ till the point c of local coordinates $(z_0, z_2) = (0, e^{i\frac{\pi}{3}}(-\varepsilon))$ in the local chart at $[0 : 1 : 0]$. Next, γ goes back to L_0 through the handle at $[0 : 1 : 0]$, following "backwards" the path parametrized by

$$(0, e^{-t}(-\varepsilon_1) + e^t e^{i\frac{\pi}{3}}(-\varepsilon_1))$$


 FIGURE 8. The loop γ

in local coordinates in the chart at $[0 : 1 : 0]$ till it reaches the point d of local coordinates $(0, -\varepsilon)$ in the chart at $[0 : 0 : 1]$. The path then follows $L_0 \cap \{z_0 = 0\}$ "down" to $[0 : 0 : 1]$ till the point e of local coordinates $(0, -\varepsilon)$ in the chart at $[0 : 0 : 1]$. Now we close the loop γ with a path contained in $L_0 \cap L$ but leaving the coordinate- $\mathbb{C}\mathbb{P}^1 \{z_0 = 0\}$ by following the half circle parametrized in the chart at $[0 : 0 : 1]$ by

$$\{(-\varepsilon \sin(t), -\varepsilon \cos(t)) \mid t \in [0; \pi]\}.$$

This loop encloses a disk u in $\mathbb{C}\mathbb{P}^2$ which can be described as the union of the portion of sphere $\{z_0 = 0\}$ lying between L_0 and L_1 in the sector making a $\frac{\pi}{3}$ -angle and delimited by γ in the "north", the portion of the same sphere in the $\frac{2\pi}{3}$ -sector between γ and the segment in L_0 (but not L) of coordinates in the chart at $[0 : 0 : 1]$

$$[e; a] = \{(0, z_2) \mid z_2 \in [-\varepsilon, \varepsilon]\},$$

and the half disk in L_0 enclosed by this segment $[e; a]$ and γ (the part of the disk u in $\{z_0 = 0\}$) and the half disk are glued along the segment $[e; a]$.

This disk u and the similar ones we can build between the other handles together with the disks considered before generate $H_2(\mathbb{C}\mathbb{P}^2, L)$, so that the monotonicity of L will follow from the next two lemmas which compute the area and the Maslov class of u . \square

Lemma 3.2. *The disk u has area $\frac{1}{2}$.*

Proof. We will compute the area of u by adding the area of the three portions we described above.

As we noticed in Section 2.2, we can make the Lagrangian surgery such that the areas between the restrictions to each \mathbb{C} -factor of \mathbb{C}^2 of the original Lagrangians and the handles are small and equal. We do the surgeries as

in Section 2.2 so that these areas are equal to $a(\varepsilon)$ small at each of the intersection points.

The area of the first portion is then the area of the $\frac{\pi}{3}$ -sector, namely one sixth of the total area of the sphere, minus the area lost at the handle, namely $a(\varepsilon)$. The area in the other sector of the coordinate projective line is the area gained through the handle, that is $a(\varepsilon)$. The contribution of the portions of the disk in the coordinate sphere is thus $\frac{1}{2}$. The contribution of the half-disk in L_0 is zero as this half-disk lies totally in a Lagrangian submanifold. \square

Lemma 3.3. *The disk u has Maslov class 1.*

Proof. The first crucial remark is that the disk u is lying entirely in the chart at $[0 : 0 : 1]$, so that the tangent bundle of $\mathbb{C}\mathbb{P}^2$ is already trivialized along the disk when we work in this chart. Now, to compute the Maslov class of this disk, we will write the loop in the Lagrangian Grassmannian we have along the boundary γ as the action of a loop $A(t)$ of matrices in $U(2)$ on the reference linear Lagrangian space \mathbb{R}^2 of \mathbb{C}^2 . The Maslov class $\mu(u)$ is then the degree of the square of the determinant of A seen as a map from \mathbb{S}^1 to \mathbb{S}^1 .

To describe this action, we will decompose the action along the different portions of the loop we considered above. The loop γ is the concatenation of the paths γ_1 from a to b , γ_2 from b to c , ... and, γ_5 from e to a . On each of these paths, we will decompose the action of $U(2)$ so that the Maslov class of u can be written as the product of these different paths of matrices.

At the point a , we are in $L_0 \cap L$, with L_0 a linear Lagrangian in the chart, so that the submanifold identifies with its tangent space, namely \mathbb{R}^2 , our reference linear Lagrangian subspace.

Between the points b and c , we stay on L_1 , the tangent space is identically equal to the linear Lagrangian subspace $l_1 = \{(e^{i\frac{\pi}{3}}x_0, e^{-i\frac{\pi}{3}}x_1) \mid x_0, x_1 \in \mathbb{R}\}$ so that $A(t)$ is the identity along γ_2 and this portion has no contribution to the degree.

Similarly, along γ_4 and γ_5 , we stay on the same linear Lagrangian (either l_1 or l_0) so that the matrix $A(t)$ is again the identity along these portions of γ .

We are left to compute the contributions of the handles to the Maslov class.

Computing the contribution of the handle at $[0 : 0 : 1]$ along γ_1 is straightforward because we have the parametrization of the handle explicitly written in this chart, namely

$$\left\{ e^{-t} \begin{pmatrix} x_0 \\ x_1 \end{pmatrix} + e^t \begin{pmatrix} e^{i\frac{\pi}{3}}x_0 \\ e^{-i\frac{\pi}{3}}(-x_1) \end{pmatrix} \middle| \begin{array}{l} t \in \mathbb{R} \\ x_0^2 + x_1^2 = \varepsilon_1^2 \end{array} \right\}.$$

Then the tangent spaces to that handle along the points in $\{z_0 = 0\}$ can be parametrized by

$$\left\{ e^{-t} \begin{pmatrix} X_0 \\ -\varepsilon_1 T \end{pmatrix} + e^t \begin{pmatrix} e^{i\frac{\pi}{3}}X_0 \\ e^{-i\frac{\pi}{3}}(-\varepsilon_1)T \end{pmatrix} \middle| t, T, X_0 \in \mathbb{R} \right\}.$$

For t going to $-\infty$, the tangent space is asymptotic to \mathbb{R}^2 for which we can take the canonical basis $\{(1, 0), (0, 1)\}$. Through the handle, the vectors $(X_0, -\varepsilon_1 T) = (1, 0)$ and $(X_0, -\varepsilon_1 T) = (0, 1)$ are mapped to $(X_0, -\varepsilon_1 T) = (e^{i\frac{\pi}{3}}, 0)$ and $(X_0, -\varepsilon_1 T) = (0, e^{-i\frac{\pi}{3}})$, a basis of l_1 through a path of matrices homotopic to

$$A_1(s) = \begin{pmatrix} e^{-is\frac{\pi}{3}} & \\ 0 & e^{is\frac{\pi}{3}} \end{pmatrix}$$

for s going from $s = 0$ to $s = 1$. The determinant of A_1 being identically equal to 1, this part of γ will not contribute to the degree.

The contribution of the handle at $[0 : 1 : 0]$ can also be computed thanks to the explicit parametrization of the handle, but we have first to write it in the chart at $[0 : 0 : 1]$ for our computation. In that chart, the handle is now parametrized by

$$\left\{ \left(\frac{e^{-t}x_0 + e^t e^{i\frac{2\pi}{3}}x_0}{e^{-t}x_2 + e^t e^{i\frac{\pi}{3}}x_2}, \frac{1}{e^{-t}x_2 + e^t e^{i\frac{\pi}{3}}x_2} \right) \middle| \begin{array}{l} t \in \mathbb{R} \\ x_0^2 + x_2^2 = \varepsilon_1^2 \end{array} \right\},$$

so that the tangent spaces are described by

$$\left\{ \left(\frac{e^{-t}X_0 + e^t e^{i\frac{2\pi}{3}}X_0}{e^{-t}x_2 + e^t e^{i\frac{\pi}{3}}x_2}, -\frac{e^{-t}x_2T + e^t e^{i\frac{\pi}{3}}x_2T}{(e^{-t}x_2 + e^t e^{i\frac{\pi}{3}}x_2)^2} \right) \middle| \begin{array}{l} t, T, X_0 \in \mathbb{R} \\ x_2 = -\varepsilon_1 \end{array} \right\}.$$

When t tends to $-\infty$, the handle is indeed asymptotic to l_0 and when t tends to $+\infty$, the handle is asymptotic to $l_1 = \{(e^{i\frac{\pi}{3}}x_0, e^{-i\frac{\pi}{3}}x_1) \mid x_0, x_1 \in \mathbb{R}\}$. The canonical basis $\{(1, 0), (0, 1)\}$ is mapped through the handle to $\{(e^{i\frac{\pi}{3}}, 0), (0, -e^{-i\frac{\pi}{3}})\}$. The action on the first coordinate can be homotopic to the path in $U(1)$ $s \in [0, 1] \mapsto e^{is\frac{\pi}{3}}$ or to $s \in [0, 1] \mapsto e^{-is\frac{5\pi}{3}}$. But for $t = 0$ at the middle of the handle, one can check via the formula that the image of the vector $(1, 0)$ is positively proportional to the vector $(e^{i\frac{\pi}{6}}, 0)$ so that the path is

$$s \in [0, 1] \mapsto e^{is\frac{\pi}{3}}.$$

For the second coordinate, in a similar manner one can act either by a path homotopic to $s \in [0, 1] \mapsto e^{is\frac{2\pi}{3}}$ or to $s \in [0, 1] \mapsto e^{-is\frac{4\pi}{3}}$. For $t = 0$ at the middle of the handle, one can check via the formula that the image of the vector $(0, 1)$ is positively proportional to the vector $(0, -e^{i\frac{\pi}{3}})$ so that the path is

$$s \in [0, 1] \mapsto e^{-is\frac{4\pi}{3}}.$$

The contribution of the handle along $\{z_0 = 0\}$ from l_0 to l_1 is thus homotopic to

$$s \in [0, 1] \mapsto \begin{pmatrix} e^{is\frac{\pi}{3}} & \\ 0 & e^{-is\frac{4\pi}{3}} \end{pmatrix}.$$

But along γ_3 we move from L_1 to L_0 so that the matrix of the action of $U(2)$ along this portion of the boundary is homotopic to

$$s \in [0, 1] \mapsto \begin{pmatrix} e^{-is\frac{\pi}{3}} & \\ 0 & e^{+4is\frac{\pi}{3}} \end{pmatrix},$$

whose determinant squared is equal to

$$\begin{array}{ccc} [0, 1] & \longrightarrow & \mathbb{S}^1 \\ s & \longmapsto & e^{2is\pi} \end{array}$$

which is of degree 1. In conclusion, the Maslov class of u being the sum of all the contributions of the portions of the loop γ is equal to 1. \square

3.2. A monotone $K\#\Sigma_4$ in $\mathbb{CP}^1 \times \mathbb{CP}^1$. In this section, we explain the construction of a monotone Lagrangian embedding of the connected sum of a Klein bottle and a surface of genus 4 in the product $\mathbb{CP}^1 \times \mathbb{CP}^1$.

Let us normalize the symplectic form on $\mathbb{CP}^1 \times \mathbb{CP}^1$ such that the area of a projective line is 2:

$$\int_{\mathbb{CP}^1} \omega = 2.$$

With this normalisation, $\mathbb{CP}^1 \times \mathbb{CP}^1$ is monotone with monotonicity constant 1:

$$\forall v \in H_2(\mathbb{CP}^1 \times \mathbb{CP}^1), \int_v \omega = c_1(T(\mathbb{CP}^1 \times \mathbb{CP}^1))(v),$$

and any monotone Lagrangian submanifold L will have a monotonicity constant equal to $\frac{1}{2}$:

$$\forall u \in H_2(\mathbb{CP}^1 \times \mathbb{CP}^1, L), \int_u \omega = \frac{1}{2} \mu_L(u).$$

To construct a monotone $K\#\Sigma_4$ in $\mathbb{CP}^1 \times \mathbb{CP}^1$, we take two Hamiltonian isotopic copies of the real part of $\mathbb{CP}^1 \times \mathbb{CP}^1$, i.e. two Lagrangian tori, that intersect in four points and perform a suitable Lagrangian surgery at these four points.

The image of the resulting Lagrangian L under the moment map will be contained in the original moment polytope of the ambient symplectic manifold chopped at its four corners (see Figure 9).

In view of the monotonicity condition, we have here two possible choices for the copies of the real part and corresponding surgeries.

One choice is to take one copy of the real part to be

$$L_0 = \{([x_0 : x_1], [u_0 : u_1]) \in \mathbb{CP}^1 \times \mathbb{CP}^1 \mid x_0, x_1, u_0, u_1 \in \mathbb{R}\}$$

and the second one its "rotation" by i :

$$L_1 = \{([e^{i\frac{\pi}{2}}x_0 : x_1], [e^{i\frac{\pi}{2}}u_0 : u_1]) \in \mathbb{CP}^1 \times \mathbb{CP}^1 \mid x_0, x_1, u_0, u_1 \in \mathbb{R}\}.$$

They intersect each other in the four points

$$([0 : 1], [0 : 1]), ([1 : 0], [0 : 1]), ([0 : 1], [1 : 0]), ([1 : 0], [1 : 0]),$$

the preimage of the four corners under the standard moment map. Then choose the following handles at the intersection points:




FIGURE 9. The image of L under the moment map is contained in and smoothly approximates the shaded polytope.

- at $([0 : 1], [0 : 1])$, $l_0 = \{(x_0, u_0) \mid x_0, u_0 \in \mathbb{R}\}$, $l_1 = \{(ix_0, iu_0) \mid x_0, u_0 \in \mathbb{R}\}$, and we insert the handle:

$$\left\{ e^{-t} \begin{pmatrix} x_0 \\ u_0 \end{pmatrix} + e^t \begin{pmatrix} ix_0 \\ iu_0 \end{pmatrix} \mid \begin{array}{l} t \in \mathbb{R} \\ x_0^2 + u_0^2 = \varepsilon_1 \end{array} \right\},$$

- at $([1 : 0], [0 : 1])$, $l_0 = \{(x_1, u_0) \mid x_1, u_0 \in \mathbb{R}\}$, $l_1 = \{(-ix_1, iu_0) \mid x_1, u_0 \in \mathbb{R}\}$, and we insert the handle:

$$\left\{ e^{-t} \begin{pmatrix} x_1 \\ u_0 \end{pmatrix} + e^t \begin{pmatrix} -i(-x_1) \\ iu_0 \end{pmatrix} \mid \begin{array}{l} t \in \mathbb{R} \\ x_1^2 + u_0^2 = \varepsilon_1 \end{array} \right\},$$

- at $([0 : 1], [1 : 0])$, $l_0 = \{(x_0, u_1) \mid x_0, u_1 \in \mathbb{R}\}$, $l_1 = \{(ix_0, -iu_1) \mid x_0, u_1 \in \mathbb{R}\}$, and we insert the handle:

$$\left\{ e^{-t} \begin{pmatrix} x_0 \\ u_1 \end{pmatrix} + e^t \begin{pmatrix} ix_0 \\ -i(-u_1) \end{pmatrix} \mid \begin{array}{l} t \in \mathbb{R} \\ x_0^2 + u_1^2 = \varepsilon_1 \end{array} \right\},$$

- at $([1 : 0], [1 : 0])$, $l_0 = \{(x_1, u_1) \mid x_1, u_1 \in \mathbb{R}\}$, $l_1 = \{(-ix_1, -iu_1) \mid x_1, u_1 \in \mathbb{R}\}$, and we insert the handle:

$$\left\{ e^{-t} \begin{pmatrix} x_1 \\ u_1 \end{pmatrix} + e^t \begin{pmatrix} -i(-x_1) \\ -i(-u_1) \end{pmatrix} \mid \begin{array}{l} t \in \mathbb{R} \\ x_1^2 + u_1^2 = \varepsilon_1 \end{array} \right\},$$

so that the intersection of the Lagrangian obtained by surgery with any of the $\mathbb{C}\mathbb{P}^1$ which are preimages of the edges on the boundary of the moment polytope is one circle of the shape of the tennis ball seam as before.

As in the previous section, one cannot take this circle to check the monotonicity of the surface as it goes around the handle twice, but we can construct similar loops going only once around the handle and check that they satisfy the monotonicity condition.

Alternatively, one can choose the surgeries such that the intersection of the Lagrangian after surgery with each coordinate- $\mathbb{C}\mathbb{P}^1$ is a union of two circles. To ensure monotonicity in this case, we need each circle to enclose an area slightly bigger than the one we get with the choice of L_1 above.

We will take the same Lagrangian L_0 and for L_1 the following Hamiltonian isotopic copy:

$$L_1 = \left\{ \left([e^{i(\frac{\pi}{2}+\delta)}x_0 : x_1], [e^{i(\frac{\pi}{2}+\delta)}u_0 : u_1] \right) \in \mathbb{C}\mathbb{P}^1 \times \mathbb{C}\mathbb{P}^1 \mid x_0, x_1, u_0, u_1 \in \mathbb{R} \right\},$$

for a small positive parameter δ that will be fixed later. They intersect each another again in the four corners of the moment polytope of $\mathbb{C}\mathbb{P}^1 \times \mathbb{C}\mathbb{P}^1$.

In each chart around the intersection points, we are making the following choices:

- at $([0 : 1], [0 : 1])$, L_0 and L_1 are the linear subspaces $l_0 = \{(x_0, u_0) \mid x_0, u_0 \in \mathbb{R}\}$, $l_1 = \{(e^{i(\frac{\pi}{2}+\delta)}x_0, e^{i(\frac{\pi}{2}+\delta)}u_0) \mid x_0, u_0 \in \mathbb{R}\}$, and we insert the handle:

$$\left\{ e^{-t} \begin{pmatrix} x_0 \\ u_0 \end{pmatrix} + e^t \begin{pmatrix} e^{i(\frac{\pi}{2}+\delta)}x_0 \\ e^{i(\frac{\pi}{2}+\delta)}u_0 \end{pmatrix} \mid \begin{array}{l} t \in \mathbb{R} \\ x_0^2 + u_0^2 = \varepsilon_1 \end{array} \right\};$$

- at $([1 : 0], [0 : 1])$, L_0 and L_1 are the linear subspaces $l_0 = \{(x_1, u_0) \mid x_1, u_0 \in \mathbb{R}\}$, $l_1 = \{(e^{-i(\frac{\pi}{2}+\delta)}x_1, e^{i(\frac{\pi}{2}+\delta)}u_0) \mid x_1, u_0 \in \mathbb{R}\}$, and we insert the handle:

$$\left\{ e^{-t} \begin{pmatrix} x_1 \\ u_0 \end{pmatrix} + e^t \begin{pmatrix} e^{-i(\frac{\pi}{2}+\delta)}x_1 \\ e^{i(\frac{\pi}{2}+\delta)}(-u_0) \end{pmatrix} \mid \begin{array}{l} t \in \mathbb{R} \\ x_1^2 + u_0^2 = \varepsilon_1 \end{array} \right\};$$

- at $([0 : 1], [1 : 0])$, L_0 and L_1 are the linear subspaces $l_0 = \{(x_0, u_1) \mid x_0, u_1 \in \mathbb{R}\}$, $l_1 = \{(e^{i(\frac{\pi}{2}+\delta)}x_0, e^{-i(\frac{\pi}{2}+\delta)}u_1) \mid x_0, u_1 \in \mathbb{R}\}$, and we insert the handle:

$$\left\{ e^{-t} \begin{pmatrix} x_0 \\ u_1 \end{pmatrix} + e^t \begin{pmatrix} e^{i(\frac{\pi}{2}+\delta)}(-x_0) \\ e^{-i(\frac{\pi}{2}+\delta)}u_1 \end{pmatrix} \mid \begin{array}{l} t \in \mathbb{R} \\ x_0^2 + u_1^2 = \varepsilon_1 \end{array} \right\};$$

- at $([1 : 0], [1 : 0])$, L_0 and L_1 are the linear subspaces $l_0 = \{(x_1, u_1) \mid x_1, u_1 \in \mathbb{R}\}$, $l_1 = \{(e^{-i(\frac{\pi}{2}+\delta)}x_1, e^{-i(\frac{\pi}{2}+\delta)}u_1) \mid x_1, u_1 \in \mathbb{R}\}$, and we insert the handle:

$$\left\{ e^{-t} \begin{pmatrix} x_1 \\ u_1 \end{pmatrix} + e^t \begin{pmatrix} e^{-i(\frac{\pi}{2}+\delta)}(-x_1) \\ e^{-i(\frac{\pi}{2}+\delta)}(-u_1) \end{pmatrix} \mid \begin{array}{l} t \in \mathbb{R} \\ x_1^2 + u_1^2 = \varepsilon_1 \end{array} \right\}.$$

Theorem 3.4. *The construction produces a monotone Lagrangian embedding of a compact surface $K \# \Sigma_4$ in $\mathbb{C}\mathbb{P}^1 \times \mathbb{C}\mathbb{P}^1$ for an appropriate choice of δ .*

Proof. Note that even though the two Lagrangian submanifolds L_0 and L_1 are oriented (they are tori), one can check that given an orientation of the two tori, two of these handles do not preserve the orientation (this cannot be avoided, it is related to the fact that the signs of the intersection points

cancel in pairs for any choice of orientation). Therefore, we get through these four surgeries a non-orientable Lagrangian which is the connected sum of the two initial tori with one torus and two Klein bottles. It is diffeomorphic to

$$L \cong K \# \Sigma_4.$$

Following the remarks from Section 2.2, we can do the surgery in each corner of the moment map such that the areas between the handle and the initial Lagrangians are small and equal along each $\mathbb{C}\mathbb{P}^1$ preimage of the boundary of the moment polytope. With the choice of handles we made above, the intersection of L with each coordinate- $\mathbb{C}\mathbb{P}^1$ consists of two circles lying in the sectors of the coordinate sphere making an angle $\frac{\pi}{2} + \delta$.

We will pick one of these circles and the disk u it encloses in one of the $\frac{\pi}{2} + \delta$ -sectors. The area of this disk is equal to the difference of the area of one sector and $2a(\varepsilon)$:

$$\frac{1}{2} + \frac{\delta}{\pi} - 2a(\varepsilon).$$

We can now fix δ (i.e. take $\delta = 2\pi a(\varepsilon)$) such that the area of u is equal to $\frac{1}{2}$.

We now compute that the Maslov class of u is equal to 1. Let us take u in the projective line of homogeneous equation $z_0 = 0$ in the homogeneous coordinates $([z_0 : z_1], [w_0 : w_1])$ of $\mathbb{C}\mathbb{P}^1 \times \mathbb{C}\mathbb{P}^1$. In a similar way as in Section 3.1.2, we trivialize the tangent bundle of $\mathbb{C}\mathbb{P}^1 \times \mathbb{C}\mathbb{P}^1$ over u by considering u in the chart of $([0 : 1], [0 : 1])$. As before, one can decompose the loop along its boundary in four paths: two paths lying on the restriction of the initial Lagrangians to this chart and two paths inside the handles. For the first type of paths, as the restrictions of L_0 and L_1 are linear in the chart, they will not contribute to the Maslov class. The handle at $([0 : 1], [0 : 1])$ is contributing with a path from L_0 to L_1 homotopic to

$$s \in [0, 1] \mapsto \begin{pmatrix} e^{is(\frac{\pi}{2} + \delta)} & \\ 0 & e^{-is(\frac{\pi}{2} - \delta)} \end{pmatrix}.$$

The handle at $([1 : 0], [0 : 1])$ is contributing with a path from L_0 to L_1 homotopic to

$$s \in [0, 1] \mapsto \begin{pmatrix} e^{-is(\frac{\pi}{2} - \delta)} & \\ 0 & e^{is(\frac{3\pi}{2} + \delta)} \end{pmatrix},$$

so that the Maslov class of u is the degree of

$$s \in [0, 1] \mapsto e^{-is4\delta} e^{is(2\pi + 4\delta)}$$

that is equal to 1.

This is enough for checking the monotonicity of this Lagrangian $K \# \Sigma_4$ in $\mathbb{C}\mathbb{P}^1 \times \mathbb{C}\mathbb{P}^1$ as the relative homology group $H_2(\mathbb{C}\mathbb{P}^1 \times \mathbb{C}\mathbb{P}^1, K \# \Sigma_4)$ is generated by disks with boundary either on L_0 or L_1 (which satisfy the monotonicity condition as L_0 and L_1 are monotone) and the disks considered above in the $\mathbb{C}\mathbb{P}^1$'s. \square

4. CONSTRUCTION OF MONOTONE LAGRANGIAN SUBMANIFOLDS USING
THE LOCAL MODEL ALONG AN ISOTROPIC CIRCLE

4.1. Case of two copies of a torus intersecting along two circles in $\mathbb{C}\mathbb{P}^1 \times \mathbb{C}\mathbb{P}^1$. Let $P = \mathbb{C}\mathbb{P}^1 \times \mathbb{C}\mathbb{P}^1$ with the product symplectic form normalized as before.

Let us consider the two following Hamiltonian isotopic copies of the real part of P :

$$L_0 = \{([x_0 : x_1], [u_0 : u_1]) \in \mathbb{C}\mathbb{P}^1 \times \mathbb{C}\mathbb{P}^1 \mid x_0, x_1, u_0, u_1 \in \mathbb{R}\}$$

and

$$L_1 = \{([e^{i\frac{\pi}{2}}x_0 : x_1], [u_0 : u_1]) \in \mathbb{C}\mathbb{P}^1 \times \mathbb{C}\mathbb{P}^1 \mid x_0, x_1, u_0, u_1 \in \mathbb{R}\}.$$

The two Lagrangians L_0 and L_1 intersect exactly along two isotropic circles:

$$L_0 \cap L_1 = \{[0 : 1]\} \times \mathbb{R}\mathbb{P}^1 \cup \{[1 : 0]\} \times \mathbb{R}\mathbb{P}^1.$$

Let us study the neighbourhood of $N = \{[0 : 1]\} \times \mathbb{R}\mathbb{P}^1$ in P . The circle N can be covered by the following two charts of $\mathbb{C}\mathbb{P}^1 \times \mathbb{C}\mathbb{P}^1$:

$$\begin{aligned} \phi_0 : U_0 = \{[z_0 : z_1] \mid z_1 \neq 0\} \times \{[w_0 : w_1] \mid w_0 \neq 0\} &\longrightarrow \mathbb{C} \oplus \mathbb{C} \\ ([z_0 : z_1], [w_0 : w_1]) &\longmapsto \left(\frac{z_0}{z_1}, \frac{w_0}{w_1} \right) \end{aligned}$$

and

$$\begin{aligned} \phi_1 : U_1 = \{[z_0 : z_1] \mid z_1 \neq 0\} \times \{[w_0 : w_1] \mid w_1 \neq 0\} &\longrightarrow \mathbb{C} \oplus \mathbb{C} \\ ([z_0 : z_1], [w_0 : w_1]) &\longmapsto \left(\frac{z_0}{z_1}, \frac{w_0}{w_1} \right) \end{aligned}$$

so that $T(\mathbb{C}\mathbb{P}^1 \times \mathbb{C}\mathbb{P}^1)|_N$ can be trivialised along $N_0 = N \cap U_0$ and $N_1 = N \cap U_1$ as

$$T(\mathbb{C}\mathbb{P}^1 \times \mathbb{C}\mathbb{P}^1)|_{N_j} \cong N_j \times (\mathbb{C} \oplus \mathbb{C}),$$

where the first summand is just $\mathbb{C} = T_{[0:1]}\mathbb{C}\mathbb{P}^1$. In these trivialisations,

$$TN|_{N_j} \cong N_j \times (\{0\} \oplus \mathbb{R})$$

and we have

$$TN^\perp|_{N_j} \cong N_j \times (\mathbb{C} \oplus \mathbb{R})$$

so that

$$SN(N, \mathbb{C}\mathbb{P}^1 \times \mathbb{C}\mathbb{P}^1)|_{N_j} = TN^\perp/TN|_{N_j} \cong N_j \times \mathbb{C},$$

and the change of trivialisation from $N_0 \times \mathbb{C}$ to $N_1 \times \mathbb{C}$ in a fibre of a point $([0 : 1], [c : d])$ of $N_0 \cap N_1$ is the identity as the fibre at each point is $\mathbb{C} = T_{[0:1]}\mathbb{C}\mathbb{P}^1$. We are in the case where the symplectic normal bundle is the trivial complex line bundle over N .

Now, in the trivialisations, the restriction of the initial Lagrangians to $SN(N, \mathbb{C}\mathbb{P}^1 \times \mathbb{C}\mathbb{P}^1)|_{N_j}$ are:

- for L_0 : $N_j \times \mathbb{R}$,
- for L_1 : $N_j \times e^{i\frac{\pi}{2}}\mathbb{R}$,

with the identity as change of trivialisation so that the restrictions of L_0 and L_1 are globally products $N \times \mathbb{R}$ and $N \times e^{i\frac{\pi}{2}}\mathbb{R}$ respectively.

As we saw in Section 2.3, this means that the fibrewise surgery with fixed parameter ε globalizes to a bundle surgery over N .

The same construction can be done along the other intersection circle $N' = \{[1 : 0]\} \times \mathbb{R}\mathbb{P}^1$.

Let us describe now the choice of surgeries we make to produce a monotone Lagrangian. In the symplectic normal bundle of N , for any fiber \mathbb{C} of a point $p \in N$ we choose the surgery

$$\left\{ e^{-t}x_0 + e^t e^{i\frac{\pi}{2}}x_0 \mid x_0 \in \mathbb{R}, x_0^2 = \varepsilon_1^2 \right\}.$$

For the other intersection circle N' , the symplectic normal bundle is again trivial and the restriction of L_0 and L_1 are $N' \times \mathbb{R}$ and $N' \times e^{-i\frac{\pi}{2}}\mathbb{R}$ respectively, and we choose the following handle :

$$\left\{ e^{-t}x_1 + e^t e^{i\frac{\pi}{2}}(-x_1) \mid x_1 \in \mathbb{R}, x_1^2 = \varepsilon_1^2 \right\}.$$

The Lagrangian constructed via these surgeries fibers over N (and N') as both handles and the Lagrangians L_0 and L_1 do, the projection being the restriction of the projection of the product onto its second factor:

$$pr_2 : \mathbb{C}\mathbb{P}^1 \times \mathbb{C}\mathbb{P}^1 \longrightarrow \mathbb{C}\mathbb{P}^1.$$

It actually lies in the $\mathbb{C}\mathbb{P}^1$ -bundle over N , given by restricting pr_2 to $\mathbb{C}\mathbb{P}^1 \times \mathbb{R}\mathbb{P}^1$. In a fiber $\mathbb{C}\mathbb{P}^1$ of this fibration, with our choice of surgery, the restriction of L is one circle in the tennis ball seam shape cutting the fiber into two disks of equal area 1. As through this fibration we see that the Lagrangian is just a product of N and the tennis ball seam, the Maslov class of any of these disks of area 1 is the Maslov class of the tennis ball seam in the fiber $\mathbb{C}\mathbb{P}^1$, that is 2.

This shows that the Lagrangian we constructed is a torus and that it is monotone.

Unfortunately, this torus is not new, it is Hamiltonian isotopic to the real part of $\mathbb{C}\mathbb{P}^1 \times \mathbb{C}\mathbb{P}^1$. For the isotopy we could just take the extension of the exact Lagrangian isotopy that isotopes in each fiber of pr_2 the tennis ball seam on the real line $\mathbb{R}\mathbb{P}^1$. We have proved:

Theorem 4.1. *The Lagrangian bundle surgery construction above produces a monotone Lagrangian embedding of a torus L in $\mathbb{C}\mathbb{P}^1 \times \mathbb{C}\mathbb{P}^1$ which is Hamiltonian isotopic to the real part of $\mathbb{C}\mathbb{P}^1 \times \mathbb{C}\mathbb{P}^1$ and projects under the moment map to a smooth interior approximation of the shaded polytope in Figure 10.*

4.2. Recovering the Chekanov–Schlenk torus : Case of two copies of $\mathbb{R}\mathbb{P}^2$ intersecting in a point and along a circle in $\mathbb{C}\mathbb{P}^2$. We detail here how our method produces a torus Hamiltonian isotopic to the model torus we started with, i.e. the exotic torus of Chekanov and Schlenk.




FIGURE 10. The image of L under the moment map is contained in and smoothly approximates the shaded polytope.

Theorem 4.2. *With a Lagrangian surgery at a point and a Lagrangian surgery along a circle of two copies of $\mathbb{R}\mathbb{P}^2$ one can construct a monotone Lagrangian embedding of the torus in $\mathbb{C}\mathbb{P}^2$ that projects through the moment map to a smooth interior approximation of the shaded polytope in Figure 11.*




FIGURE 11. The image of the torus under the moment map is contained in and smoothly approximates the shaded polytope.

Proof. Let us consider the following two Hamiltonian isotopic copies of $\mathbb{R}\mathbb{P}^2$ in $\mathbb{C}\mathbb{P}^2$:

$$L_0 = \{[x_0 : x_1 : x_2] \in \mathbb{C}\mathbb{P}^2 \mid x_0, x_1, x_2 \in \mathbb{R}\}$$

and

$$L_1 = \{[e^{i\alpha}x_0 : e^{i\alpha}x_1 : x_2] \in \mathbb{C}\mathbb{P}^2 \mid x_0, x_1, x_2 \in \mathbb{R}\},$$

for some $\alpha \in (0, \pi)$. The two Lagrangians L_0 and L_1 intersect exactly at the point $[0 : 0 : 1]$ and along the isotropic circle

$$N = \{[x_0 : x_1 : 0] \mid (x_0, x_1) \in \mathbb{R}^2 \setminus \{(0, 0)\}\}.$$

We will make a Lagrangian bundle surgery along the isotropic circle and a simple Lagrangian surgery at the point $[0 : 0 : 1]$.

Let us first understand the neighbourhood of N in $\mathbb{C}\mathbb{P}^2$. The circle N can be covered by the following two charts of $\mathbb{C}\mathbb{P}^2$:

$$\begin{aligned} \phi_0 : U_0 = \{[z_0 : z_1 : z_2] \mid z_0 \neq 0\} &\longrightarrow \mathbb{C} \oplus \mathbb{C} \\ [z_0 : z_1 : z_2] &\longmapsto \begin{pmatrix} z_1 \\ z_0, z_2 \end{pmatrix} \end{aligned}$$

and

$$\begin{aligned} \phi_1 : U_1 = \{[z_0 : z_1 : z_2] \mid z_1 \neq 0\} &\longrightarrow \mathbb{C} \oplus \mathbb{C} \\ [z_0 : z_1 : z_2] &\longmapsto \begin{pmatrix} z_0 \\ z_1, z_2 \end{pmatrix} \end{aligned}$$

so that $T\mathbb{C}\mathbb{P}^2|_N$ can be trivialised along $N_0 = N \cap U_0$ and $N_1 = N \cap U_1$ as

$$T\mathbb{C}\mathbb{P}^2|_{N_j} \cong N_j \times (\mathbb{C} \oplus \mathbb{C}).$$

In these trivialisations,

$$TN|_{N_j} \cong N_j \times (\mathbb{R} \oplus \{0\})$$

and we have

$$TN^\perp|_{N_j} \cong N_j \times (\mathbb{R} \oplus \mathbb{C})$$

so that

$$SN(N, \mathbb{C}\mathbb{P}^2)|_{N_j} = TN^\perp/TN|_{N_j} \cong N_j \times \mathbb{C},$$

and the change of trivialisation from $N_0 \times \mathbb{C}$ to $N_1 \times \mathbb{C}$ in a fibre over a point $[a : b : 0]$ of $N_0 \cap N_1$ is $([a : b : 0], Z) \mapsto ([a : b : 0], \frac{a}{b}Z)$. As the intersection $N_0 \cap N_1$ retracts onto $\{[1 : 1 : 0], [-1 : 1 : 0]\}$, we have only two changes of trivialisation to consider: in $[1 : 1 : 0]$ it is the identity, and in $[-1 : 1 : 0]$ it is minus the identity.

Now, in the trivialisations, the trace of the initial Lagrangians in $SN(N, \mathbb{C}\mathbb{P}^2)|_{N_j}$ are:

- for L_0 : $N_j \times \mathbb{R}$,
- for L_1 : $N_j \times e^{-i\alpha}\mathbb{R}$,

with the same change of trivialization as before.

One can then make a Lagrangian surgery in each fibre \mathbb{C} , and it globalizes to a Lagrangian subbundle over N in the symplectic normal bundle.

We next do a surgery at the transverse intersection point $[0 : 0 : 1]$, so that we get an embedded Lagrangian submanifold out of the surgeries on L_0 and L_1 .

We show that for some choices of handles and of α , this Lagrangian submanifold is monotone. Take $\alpha = \frac{2\pi}{3} + \delta$, for $\delta > 0$ a small parameter to be

determined later. We choose the bundle surgery that in each fiber \mathbb{C} over a point of N it is parametrized by

$$\left\{ e^{-t}x_2 + e^t e^{-i\alpha}x_2 \left| \begin{array}{l} t \in \mathbb{R} \\ x_2 \in \mathbb{R}, x_2^2 = \varepsilon_1^2 \end{array} \right. \right\},$$

followed by a symmetric smoothing.

In the chart at the transverse intersection point $[0 : 0 : 1]$, the two Lagrangians are the linear Lagrangian subspaces $l_0 = \mathbb{R}^2$ and $l_1 = \{(e^{i\alpha}x_0, e^{i\alpha}x_1) | x_0, x_1 \in \mathbb{R}\}$ and we choose the handle

$$\left\{ e^{-t} \begin{pmatrix} x_0 \\ x_1 \end{pmatrix} + e^t \begin{pmatrix} e^{i\alpha}x_0 \\ e^{i\alpha}x_1 \end{pmatrix} \left| \begin{array}{l} t \in \mathbb{R} \\ x_0^2 + x_1^2 = \varepsilon_1^2 \end{array} \right. \right\}.$$

After surgery, the restriction to the coordinate- $\mathbb{C}\mathbb{P}^1$ s of homogeneous equations $z_0 = 0$ (resp. $z_1 = 0$) of the Lagrangian we constructed is the union of two circles enclosing disks of area

$$1 + \frac{\delta}{2\pi} - 2a(\varepsilon).$$

One checks with the same method as before that the Maslov class of these disks is equal to 2. Then we choose δ such that the area of each of these disks is 1. This is enough to check monotonicity since this Lagrangian is a circle subbundle over N in the normal bundle of $\{z_2 = 0\}$, and hence a torus. Let us denote it by Θ_{surg} . We have checked the monotonicity condition on just one generator of the relative second homology group, the monotonicity on a second generator is immediate as it can be represented by a disk with boundary on L_0 or L_1 for which this condition is satisfied. \square

Theorem 4.3. *The Lagrangian torus Θ_{surg} is Hamiltonian isotopic to the modified Chekanov torus in $\mathbb{C}\mathbb{P}^2$ and consequently also to the Chekanov-Schlenk exotic torus.*

Proof. We use the strategy of [8] and prove that the torus obtained by surgery is invariant under the same Hamiltonian action (called ρ_{Ch} in [8]) as the modified Chekanov torus.

The modified Chekanov torus is invariant under the Hamiltonian circle action ρ_{Ch} defined on $\mathbb{C}\mathbb{P}^2$ by applying the following matrix to the homogeneous coordinates:

$$\begin{pmatrix} \cos(\theta) & -\sin(\theta) & 0 \\ \sin(\theta) & \cos(\theta) & 0 \\ 0 & 0 & 1 \end{pmatrix},$$

where $\theta \in \mathbb{R}/2\pi\mathbb{Z}$.

This circle action preserves L_0, L_1, N and $[0 : 0 : 1]$.

In the chart around $[0 : 0 : 1]$, it restricts on a ball in \mathbb{C}^2 to the action defined by the rotation matrix

$$\begin{pmatrix} \cos(\theta) & -\sin(\theta) \\ \sin(\theta) & \cos(\theta) \end{pmatrix}.$$

In particular,

- it preserves the handle of the local surgery at the point $[0 : 0 : 1]$,
- one can ask the smoothing to be invariant under the action, so that the entire surgery is preserved,
- the action commutes with the homotheties defining the conformal transformation.

Moreover, L_0 is the orbit under this action restricted to the projective line of homogeneous equation $z_0 = 0$ and even to $\{[0 : x_1 : x_2] \in \mathbb{C}\mathbb{P}^2 \mid x_1, x_2 \in \mathbb{R}, x_1 \geq 0\}$. Similarly, L_1 is the orbit under this action of $\{[0 : e^{i\alpha}x_1 : x_2] \in \mathbb{C}\mathbb{P}^2 \mid x_1, x_2 \in \mathbb{R}, x_1 \geq 0\}$, and the handle is the orbit of one of its branches intersected with $z_0 = 0$, for instance of $\{[0 : e^{-t}x_1 + e^te^{i\alpha}x_1 : x_2] \in \mathbb{C}\mathbb{P}^2 \mid x_1, x_2 \in \mathbb{R}, x_1 = +\varepsilon_1\}$.

One can describe the handle in homogeneous coordinates as

$$\{[x_0 : x_1 : e^{-t}x_2 + e^te^{-i\alpha}x_2] \in \mathbb{C}\mathbb{P}^2 \mid x_0, x_1 \in \mathbb{R}, x_2^2 = \varepsilon_1^2\}$$

and see that it is preserved by the circle action ρ_{Ch} . In fact, it is the orbit of one of its branches in $z_0 = 0$, for instance

$$\{[0 : x_1 : e^{-t}x_2 + e^te^{-i\alpha}x_2] \in \mathbb{C}\mathbb{P}^2 \mid x_0, x_1 \in \mathbb{R}, x_2 = +\varepsilon_1\}.$$

We can also take a smoothing that is preserved by the circle action and the orbit under the circle action of the smoothing in $z_0 = 0$.

This shows that the torus Θ_{surg} is the orbit under the circle action ρ_{Ch} of one of the two circles that constitute its intersection with $z_0 = 0$. Actually, as the third homogeneous coordinate in the handle never vanishes, the torus Θ_{surg} lies in the complement of $z_2 = 0$, that is the ball of capacity 3 around $[0 : 0 : 1]$, and is also in this ball the orbit under the circle action of the circle lying in its intersection with the half plane of equations $z_0 = 0$ and $\Re(z_0) \geq 0$.

As this circle is by construction of area 1, one can isotope it inside this half-plane to the curve γ used to define $\tilde{\Theta}_{\text{Ch}}$. This isotopy composed with the Hamiltonian circle action gives an exact Lagrangian isotopy between the two tori that can be extended to a Hamiltonian isotopy of $\mathbb{C}\mathbb{P}^2$ as in [8]. \square

REFERENCES

- [1] Miguel Abreu and Leonardo Macarini. Remarks on Lagrangian intersections in toric manifolds. *Trans. Amer. Math. Soc.*, 365(7):3851–3875, 2013.
- [2] Michèle Audin. *Torus actions on symplectic manifolds*, volume 93 of *Progress in Mathematics*. Birkhäuser Verlag, Basel, revised edition, 2004.
- [3] Paul Biran and Octav Cornea. Rigidity and uniruling for Lagrangian submanifolds. *Geom. Topol.*, 13(5):2881–2989, 2009.
- [4] Ana Cannas da Silva. *Lectures on symplectic geometry*, volume 1764 of *Lecture Notes in Mathematics*. Springer-Verlag, Berlin, 2001.
- [5] Yuri Chekanov and Felix Schlenk. Twist tori I: Construction and classification. *in preparation*.
- [6] Yuri Chekanov and Felix Schlenk. Twist tori II: Non-displaceability. *in preparation*.
- [7] Yuri Chekanov and Felix Schlenk. Notes on monotone Lagrangian twist tori. *Electron. Res. Announc. Math. Sci.*, 17:104–121, 2010.

- [8] Agnès Gadbled. On exotic monotone Lagrangian tori in $\mathbb{C}\mathbb{P}^2$ and $\mathbb{S}^2 \times \mathbb{S}^2$. *J. Symplectic Geom.*, 11(3):343–361, 2013.
- [9] A. B. Givental. Lagrangian imbeddings of surfaces and the open Whitney umbrella. *Funktsional. Anal. i Prilozhen.*, 20(3):35–41, 96, 1986.
- [10] Hông-Vân Lê. A minimizing deformation of Legendrian submanifolds in the standard sphere. *Differential Geom. Appl.*, 21(3):297–316, 2004.
- [11] Yong-Geun Oh. Floer cohomology of Lagrangian intersections and pseudo-holomorphic disks. I. *Comm. Pure Appl. Math.*, 46(7):949–993, 1993.
- [12] Leonid Polterovich. The surgery of Lagrange submanifolds. *Geom. Funct. Anal.*, 1(2):198–210, 1991.
- [13] Marcin Poźniak. Floer homology, Novikov rings and clean intersections. In *Northern California Symplectic Geometry Seminar*, volume 196 of *Amer. Math. Soc. Transl. Ser. 2*, pages 119–181. Amer. Math. Soc., Providence, RI, 1999.
- [14] V. V. Shevchishin. Lagrangian embeddings of the Klein bottle and the combinatorial properties of mapping class groups. *Izv. Ross. Akad. Nauk Ser. Mat.*, 73(4):153–224, 2009.
- [15] Alan Weinstein. *Lectures on symplectic manifolds*. American Mathematical Society, Providence, R.I., 1977. Expository lectures from the CBMS Regional Conference held at the University of North Carolina, March 8–12, 1976, Regional Conference Series in Mathematics, No. 29.
- [16] Alan Weinstein. Neighborhood classification of isotropic embeddings. *J. Differential Geom.*, 16(1):125–128, 1981.

(MA) CENTER FOR MATHEMATICAL ANALYSIS, GEOMETRY AND DYNAMICAL SYSTEMS, INSTITUTO SUPERIOR TÉCNICO, UNIVERSIDADE DE LISBOA, AV. ROVISCO PAIS, 1049-001 LISBOA, PORTUGAL

E-mail address: `mabreu@math.tecnico.ulisboa.pt`

(AG) CENTRO DE MATEMÁTICA DA UNIVERSIDADE DO PORTO, DEPARTAMENTO DE MATEMÁTICA DA FCUP, RUA DO CAMPO ALEGRE, 687, 4169-007 PORTO, PORTUGAL

E-mail address: `agnes.gadbled@fc.up.pt`