

**Exercise 1** – Open string mode expansion

Consider an open string  $X(\tau, \sigma)$  with  $\sigma \in [0, \pi]$ .

a) Define  $\hat{X}(\tau, \sigma)$  by

$$\hat{X}(\tau, \sigma) = \begin{cases} X(\tau, \sigma) & , \quad 0 \leq \sigma \leq \pi \\ X(\tau, -\sigma) & , \quad -\pi \leq \sigma < 0 \end{cases}$$

Show that the extension of  $\hat{X}(\tau, \sigma)$  to  $\sigma \in \mathbb{R}$  as a smooth periodic function with period  $2\pi$  requires Neumann-Neumann (NN) boundary conditions for  $X(\tau, \sigma)$  at  $\sigma = 0, \pi$ . Use this to obtain the mode expansion for the NN open string: consider first the mode expansion of  $\partial_{\pm} \hat{X}$ , then integrate this to obtain  $X(\tau, \sigma)$  subject to NN boundary conditions.

b) Redo the procedure for Dirichlet-Dirichlet (DD) boundary conditions, by suitably defining  $\hat{X}(\tau, \sigma)$  so as to incorporate DD boundary conditions for  $X(\tau, \sigma)$ . Then, integrate the solution for  $\partial_{\pm} \hat{X}$  to obtain  $X(\tau, \sigma)$  subject to DD boundary conditions.

c) Similarly, find the mode expansion for the DN open string and for the ND open string.

**Exercise 2** – Virasoro algebra

Let  $L_n$  denote the normal ordered operators arising in light-cone quantization,

$$L_n = \frac{1}{2} \sum_{m=-\infty}^{\infty} : \alpha_{n-m}^i \alpha_m^i : \quad , \quad i = 1, \dots, D-2 \quad , \quad n \in \mathbb{Z} . \quad (1)$$

a) Using the commutator relation  $[A, BC] = [A, B]C + B[A, C]$  show that

$$[\alpha_m^i, L_n] = m \alpha_{m+n}^i . \quad (2)$$

b) Using (2), show that the  $L_n$  satisfied the following, centrally extended algebra, called Virasoro algebra,

$$[L_m, L_n] = (m - n) L_{m+n} + \frac{c}{12} m (m^2 - 1) \delta_{m+n, 0} \quad , \quad (3)$$

where  $[c, L_n] = 0$ . Verify that the central charge  $c$  arises because of normal ordering, and determine its value.

**Exercise 3** – Analytic continuation of the zeta function  $\zeta(s)$ 

Consider the gamma function  $\Gamma(s) = \int_0^{\infty} dt e^{-t} t^{s-1}$ ,  $s \in \mathbb{C}$ . It is absolutely convergent for  $\Re(s) > 0$ . Let  $t \rightarrow nt$  in this integral, and use the resulting equation to prove that

$$\Gamma(s) \zeta(s) = \int_0^{\infty} dt \frac{t^{s-1}}{e^t - 1} \quad , \quad \Re(s) > 1 \quad , \quad (4)$$

where  $\zeta(s) = \sum_{n=1}^{\infty} n^{-s}$ ,  $\Re(s) > 1$ . Show that for  $\Re(s) > 1$

$$\begin{aligned} \Gamma(s) \zeta(s) = & \int_0^1 dt t^{s-1} \left( \frac{1}{e^t - 1} - \frac{1}{t} + \frac{1}{2} - \frac{t}{12} \right) + \frac{1}{s-1} - \frac{1}{2s} + \frac{1}{12(s+1)} \\ & + \int_1^{\infty} dt \frac{t^{s-1}}{e^t - 1} . \end{aligned} \tag{5}$$

Explain why the first integral on the right-hand side above is well behaved for  $\Re(s) > -2$ .

The right-hand side defines the analytic continuation of the left-hand side to  $\Re(s) > -2$ . Using that  $\Gamma(s)$  has a simple pole at  $s = -1$  with residue  $-1$ , show that  $\zeta(-1) = -1/12$ , a celebrated result used in light-cone quantization.