

Exercise 1 – The modular group $SL(2, \mathbb{Z})$

The special linear group $SL(2, \mathbb{R})$ acts on the complex upper half plane $\mathcal{H} = \{\tau \in \mathbb{C} | \text{Im } \tau > 0\}$ by fractional linear transformations

$$\tau \mapsto M \cdot \tau = \frac{a\tau + b}{c\tau + d} \quad , \quad M = \begin{pmatrix} a & b \\ c & d \end{pmatrix} \in SL(2, \mathbb{R}) . \quad (1)$$

a) Verify:

1. $M \cdot \tau \in \mathcal{H} \quad \forall M \in SL(2, \mathbb{R}) , \tau \in \mathcal{H}$
2. $(M_1 M_2) \cdot \tau = M_1 \cdot (M_2 \cdot \tau) \quad \forall M_1, M_2 \in SL(2, \mathbb{R}) , \tau \in \mathcal{H}$

b) The set $\bar{\mathcal{F}} = \{\tau \in \mathbb{C} | |\tau| \geq 1, |\text{Re } \tau| \leq \frac{1}{2}\}$ denotes the closure of the fundamental domain for the modular group $SL(2, \mathbb{Z})$. Show that any point $\tau \in \mathcal{H}$ can be mapped into $\bar{\mathcal{F}}$ by an application of T- and S-transformations, where

$$T \cdot \tau = \tau + 1 \quad , \quad S \cdot \tau = -1/\tau . \quad (2)$$

Exercise 2 – Eisenstein series of weight $k > 2$

Let $k \in \mathbb{Z}$. A modular form of weight k is a function $f : \mathcal{H} \rightarrow \mathbb{C}$ that is holomorphic on $\mathcal{H} \cup \{\infty\}$, and that satisfies

$$f(M \cdot \tau) = (c\tau + d)^k f(\tau) \quad \forall \tau \in \mathcal{H} , \forall M \in SL(2, \mathbb{Z}) . \quad (3)$$

Choosing $M = T$, we have $f(\tau + 1) = f(\tau) \quad \forall \tau \in \mathcal{H}$, and hence f is a periodic function of period 1. Introducing $q = e^{2\pi i \tau}$, it has the Fourier development, valid in the open complex unit disc,

$$f(\tau) = \sum_{n=0}^{\infty} a_n q^n . \quad (4)$$

Consider the following series for even $k \geq 4$ ($k \in \mathbb{N}$),

$$G_k(\tau) = \frac{(k-1)!}{2(2\pi i)^k} \sum_{m,n \in \mathbb{Z} \text{ with } (m,n) \neq (0,0)} \frac{1}{(m\tau + n)^k} \quad , \quad \tau \in \mathcal{H} . \quad (5)$$

This series, called Eisenstein series, is absolutely and locally uniformly convergent for $k > 2$.

a) Show that $G_k(\tau)$ is a modular form of weight k .

b) Show that the Fourier expansion of the Eisenstein series is

$$G_k(\tau) = \frac{(k-1)! \zeta(k)}{(2\pi i)^k} + \sum_{n=1}^{\infty} q^n \sigma_{k-1}(n) , \quad (6)$$

where $\zeta(k) = \sum_{n=1}^{\infty} n^{-k}$ and where $\sigma_{k-1}(n)$ denotes the sum of the $(k-1)$ st power of the positive divisors of n , i.e. $\sigma_{k-1}(n) = \sum_{r|n} r^{k-1}$. *Hint:* Use $\frac{\pi}{\tan(\pi\tau)} = \sum_{m \in \mathbb{Z}} \frac{1}{\tau+m}$, $\tau \in \mathbb{C} \setminus \mathbb{Z}$.

Exercise 3 – The Eisenstein series of weight 2

The Eisenstein series of weight 2 is defined by

$$G_2(\tau) = \frac{\zeta(2)}{(2\pi i)^2} + \sum_{n=1}^{\infty} q^n \sigma_1(n) . \quad (7)$$

a) Show that $G_2(\tau)$ equals

$$G_2(\tau) = -\frac{1}{4\pi^2} \left[\frac{1}{2} \sum_{n \neq 0} \frac{1}{n^2} + \frac{1}{2} \sum_{m \neq 0} \sum_{n \in \mathbb{Z}} \frac{1}{(m\tau + n)^2} \right] , \quad (8)$$

where one first carries out the summation over n , and then over m . Note that this series is not absolutely convergent, and hence one cannot interchange the order of summation to obtain $G_2(-1/\tau) = \tau^2 G_2(\tau)$.

b) Define $G_{2,\epsilon}(\tau)$ by

$$G_{2,\epsilon}(\tau) = -\frac{1}{4\pi^2} \left[\frac{1}{2} \sum_{m,n \in \mathbb{Z} \text{ with } (m,n) \neq (0,0)} \frac{1}{(m\tau + n)^2 |m\tau + n|^{2\epsilon}} \right] , \quad (9)$$

which converges absolutely. How does it transform under modular transformations $\tau \mapsto M \cdot \tau$ with $M \in SL(2, \mathbb{Z})$?

c) It can be shown that

$$G_2^*(\tau) \equiv \lim_{\epsilon \rightarrow 0} G_{2,\epsilon}(\tau) = G_2(\tau) + \frac{1}{8\pi \operatorname{Im} \tau} . \quad (10)$$

Using this, deduce the transformation behaviour

$$G_2(M \cdot \tau) = (c\tau + d)^2 G_2(\tau) + \frac{i}{4\pi} c(c\tau + d) . \quad (11)$$

c) Show that the Dedekind η function

$$\eta(\tau) = q^{1/24} \prod_{n=1}^{\infty} (1 - q^n) \quad (12)$$

satisfies

$$\frac{d}{d\tau} \ln \eta(\tau) = -2\pi i G_2(\tau) . \quad (13)$$

d) Using the above, deduce the transformation law

$$\eta(-1/\tau) = \sqrt{-i\tau} \eta(\tau) . \quad (14)$$