

DYNAMICAL IMPLICATIONS OF CONVEXITY BEYOND DYNAMICAL CONVEXITY

MIGUEL ABREU AND LEONARDO MACARINI

ABSTRACT. We establish sharp dynamical implications of convexity on symmetric spheres that do not follow from dynamical convexity. It allows us to show the existence of elliptic and non-hyperbolic periodic orbits and to furnish new examples of dynamically convex contact forms, in any dimension, that are not equivalent to convex ones via contactomorphisms that preserve the symmetry. Moreover, these examples are C^1 -stable in the sense that they are actually not equivalent to convex ones via contactomorphisms that are C^1 -close to those preserving the symmetry. We also show the multiplicity of symmetric non-hyperbolic and symmetric (not necessarily non-hyperbolic) closed Reeb orbits under suitable pinching conditions.

CONTENTS

1. Introduction and main results	2
1.1. Introduction	2
1.2. Main results	4
1.3. Organization of the paper	10
1.4. Conventions	11
1.5. Acknowledgements	11
2. Equivariant symplectic homology and index theory	11
2.1. Equivariant symplectic homology	11
2.2. Lusternik-Schnirelmann theory	13
2.3. Grading	14
2.4. Index computations and non-vanishing of the first Chern class in lens spaces	16
2.5. Bott's function	20
3. Proof of Theorem 1.4	22
3.1. Idea of the proof	22
3.2. Proof of the theorem	22
4. Proof of Corollary 1.10	27
5. Proof of Theorem 1.14	27
5.1. Proof of Assertion 1	29
5.2. Proof of Lemmas 5.4 and 5.5	30
5.3. Proof of Assertion 2	35
5.4. Proof of Lemmas 5.12 and 5.13	37
6. Proof of Theorem 1.6	39

2010 *Mathematics Subject Classification.* 53D40, 37J10, 37J55.

Key words and phrases. Closed orbits, Conley-Zehnder index, Reeb flows, dynamical convexity, equivariant symplectic homology.

The authors were partially supported by FCT/Portugal through projects UID/MAT/04459/2020 and PTDC/MAT-PUR/29447/2017.

6.1. Proof of Assertion 1	39
6.2. Proof of Assertion 2	40
6.3. Proof of Assertion 3	40
7. Proof of Theorem 1.18	40
8. Proof of Theorem 1.20	41
9. Proof of Theorem 1.21	43
References	44

1. INTRODUCTION AND MAIN RESULTS

1.1. Introduction. In this work we focus on the dynamics of Reeb flows on convex hypersurfaces with symmetry in the standard symplectic vector space. In this setting, we explore dynamical implications of convexity, the existence of elliptic and non-hyperbolic closed orbits, the relation between convexity and dynamical convexity and the multiplicity problem for symmetric non-hyperbolic and symmetric (not necessarily non-hyperbolic) closed Reeb orbits.

To elaborate, consider Reeb flows on the standard contact sphere $(S^{2n+1}, \xi_{\text{std}})$. Two of the most fundamental problems in Hamiltonian dynamics are the multiplicity question for simple (i.e., non-iterated) closed Reeb orbits and the existence of elliptic closed Reeb orbits. Two long standing conjectures establish that there are at least $n + 1$ simple closed orbits and at least one elliptic closed orbit for *any* contact form β on $(S^{2n+1}, \xi_{\text{std}})$. The first conjecture was proved for $n = 1$ by Cristofaro-Gardiner and Hutchings [11] (in the more general setting of Reeb flows in dimension three) and independently by Ginzburg, Hein, Hryniewicz and Macarini [16]; see also [24] where an alternate proof was given using a result from [16]. In higher dimensions, the question is widely open without additional assumptions on β , such as convexity or certain index requirements or non-degeneracy of closed Reeb orbits. The second conjecture is completely open in any dimension (bigger than one) without further hypotheses.

In order to explain the convexity assumption, note that there is a natural bijection between contact forms β on $(S^{2n+1}, \xi_{\text{std}})$ and starshaped hypersurfaces Σ_β in \mathbb{R}^{2n+2} in the following way. Consider the restriction of the Liouville form λ to the unit sphere $S^{2n+1} \subset \mathbb{R}^{2n+2}$. A contact form β supporting the (cooriented) standard contact structure ξ_{std} is a 1-form given by $f\lambda|_{S^{2n+1}}$, where $f : S^{2n+1} \rightarrow \mathbb{R}$ is a positive function. The bijection is given by

$$\beta = f\lambda|_{S^{2n+1}} \longleftrightarrow \Sigma_\beta = \{\sqrt{f(x)}x; x \in S^{2n+1}\}$$

and it satisfies the property that if Σ_β is the energy level of a homogeneous of degree two Hamiltonian $H_\beta : \mathbb{R}^{2n+2} \rightarrow \mathbb{R}$ then the Hamiltonian flow of H_β on Σ_β is equivalent to the Reeb flow of β . We say that β is *convex* (resp. *strictly convex*) if Σ_β bounds a convex (resp. strictly convex) domain.

Let us denote by \mathcal{P} (resp. \mathcal{P}_e) the set of simple (resp. simple elliptic) closed Reeb orbits for β . When β is strictly convex, a remarkable result due to Long and Zhu [29] asserts that $\#\mathcal{P} \geq [(n + 1)/2] + 1$. This result was improved when n is even by Wang [33], furnishing the lower bound $\#\mathcal{P} \geq [(n + 1)/2] + 1$. No lower bound for $\#\mathcal{P}_e$ is known under only the hypothesis that β is strictly convex.

Under the assumption that β is strictly convex *and invariant by the antipodal map*, Liu, Long and Zhu [25] showed that $\#\mathcal{P} \geq n + 1$ and Dell'Antonio, D'Onofrio and Ekeland [13]

proved that $\#\mathcal{P}_e \geq 1$. Thus, the aforementioned conjectures are true for this important class of contact forms.

The convexity requirement is used in several ways in these results. However, it is not natural from the point of view of contact topology since it is not a condition invariant under contactomorphisms. An alternative notion, introduced by Hofer, Wysocki and Zehnder [20], is *dynamical convexity*. A contact form β on S^{2n+1} is called dynamically convex if every closed Reeb orbit γ of β has Conley-Zehnder index $\mu(\gamma)$ greater than or equal to $n + 2$. Clearly dynamical convexity is invariant under contactomorphisms and it is not hard to see that convexity implies dynamical convexity. When β is dynamically convex, Ginzburg and Gürel [18] and, independently, Duan and Liu [14], proved that $\#\mathcal{P} \geq [(n + 1)/2] + 1$, showing that the lower bound established by Long, Zhu and Wang in [29, 33] in the convex case holds for dynamically convex hypersurfaces.

When β is dynamically convex and invariant by the antipodal map, Abreu and Macarini proved in [2] that $\mathcal{P}_e \geq 1$, extending the aforementioned result of Dell'Antonio, D'Onofrio and Ekeland [13] to dynamically convex contact forms. The extension of the results of Liu, Long and Zhu [25] to the dynamically convex scenario is more subtle and carried out in the following way. In [19], Ginzburg and Macarini introduced the notion of *strong dynamical convexity*: a symmetric contact form β on S^{2n+1} is strongly dynamically convex if it is dynamically convex and its degenerate symmetric periodic orbits satisfy a technical additional assumption involving the normal forms of the eigenvalue one; see [19] for details. They proved that if β is strongly dynamically convex then $\#\mathcal{P} \geq n + 1$. Moreover, they showed that if β is convex and invariant by the antipodal map then it is strongly dynamically convex. Using this, they were able to construct the first examples of antipodally symmetric dynamically convex contact forms (in dimension at least five) that are not equivalent to convex ones via contactomorphisms that commute with the antipodal map. The main point in the construction of these examples was to realize that, under the presence of symmetries, convexity implies more than dynamical convexity.

In this paper, continuing the study in [19], we explore further implications of convexity for symmetric contact forms on the sphere. More precisely, we consider contact forms α on lens spaces $L_p^{2n+1}(\ell_0, \dots, \ell_n)$ endowed with the induced contact structure ξ . Given a homotopy class $a \in \pi_1(L_p^{2n+1}(\ell_0, \dots, \ell_n))$ we will associate to it three rational numbers k_a , h_a and \tilde{h}_a related to the positive equivariant symplectic homology of $L_p^{2n+1}(\ell_0, \dots, \ell_n)$ and that can be easily computed from the weights ℓ_0, \dots, ℓ_n . (Note here that since, in general, $c_1(\xi) \neq 0$, the grading of the positive equivariant symplectic homology is fractional.) We say that α is convex (resp. strictly convex) if so is its lift $\beta = \pi^*\alpha$, where $\pi : S^{2n+1} \rightarrow L_p^{2n+1}(\ell_0, \dots, \ell_n)$ is the quotient projection. Given a convex (resp. strictly convex) contact form on $L_p^{2n+1}(\ell_0, \dots, \ell_n)$ and a closed orbit γ with non-trivial homotopy class a , we show in Theorem 1.4 that its index satisfies $\mu(\gamma) \geq k_a$ and the following holds: γ must be non-hyperbolic if $\mu(\gamma) < h_a$ (resp. $\mu(\gamma) < \tilde{h}_a$) and γ has to be elliptic if $\mu(\gamma) = k_a$ and a satisfies a suitable positivity assumption.

In Theorem 1.6 we show that this result is sharp: we must have a closed orbit γ with homotopy class a such that $\mu(\gamma) = k_a$, there are examples of strictly convex contact forms carrying a hyperbolic closed orbit γ with homotopy class a such that $\mu(\gamma) = h_a = \tilde{h}_a$ and examples of strictly convex contact forms with a non-elliptic closed orbit γ with positive homotopy class a such that $\mu(\gamma) = k_a + 1$. Using these results, Corollaries 1.8, 1.9 and 1.10 show that convex contact forms on $L_p^{2n+1}(\ell_0, \dots, \ell_n)$ must carry elliptic and non-hyperbolic closed

orbits in certain homotopy classes and furnish sufficient conditions to ensure the existence of such homotopy classes.

We say that α is dynamically convex if so is its lift β to S^{2n+1} ; in other words, α is dynamically convex if every *contractible* closed orbit γ of α satisfies $\mu(\gamma) \geq n + 2$. We show in Theorem 1.14 that the dynamical implications of Theorem 1.4 do not follow from dynamical convexity at all. It allows us to furnish new examples of symmetric dynamically convex contact forms on S^{2n+1} (in any dimension) that are not equivalent to convex contact forms via contactomorphisms that preserve the symmetry. Actually, these examples are not equivalent to convex contact forms via contactomorphisms C^1 -close to those that preserve the symmetry (Theorem 1.18).

Finally, let us briefly describe our results on the multiplicity of closed orbits. Given a symmetric contact form on the sphere, a refinement of the multiplicity question on periodic orbits is to give a lower bound for the number of simple *symmetric* closed Reeb orbits. It is equivalent to the multiplicity question of closed orbits of contact forms on lens spaces whose homotopy classes are generators of the fundamental group. Using Theorem 1.4 and Floer homology techniques, we obtain results on the multiplicity of *symmetric non-hyperbolic* closed Reeb orbits for suitable contact forms. More precisely, we show in Theorem 1.20 the existence of $\lfloor (n+1)/2 \rfloor$ simple symmetric non-hyperbolic closed Reeb orbits for contact forms on certain lens spaces satisfying suitable pinching conditions. At last, using a lower bound for the period of closed orbits of convex contact forms due to Croke and Weinstein [12] (which is also used in the proof of Theorem 1.20) and our Floer homology tools, we establish in Theorem 1.21 the existence of $n+1$ symmetric (not necessarily non-hyperbolic) closed orbits for convex contact forms satisfying weaker pinching conditions, generalizing a result due to Ekeland and Lasry [15].

1.2. Main results. Given an integer $p > 0$, consider the \mathbb{Z}_p -action on S^{2n+1} , regarded as a subset of $\mathbb{C}^{n+1} \setminus \{0\}$, generated by the map

$$\psi(z_0, \dots, z_n) = \left(e^{\frac{2\pi\sqrt{-1}\ell_0}{p}} z_0, e^{\frac{2\pi\sqrt{-1}\ell_1}{p}} z_1, \dots, e^{\frac{2\pi\sqrt{-1}\ell_n}{p}} z_n \right), \quad (1.1)$$

where ℓ_0, \dots, ℓ_n are integers called the *weights* of the action. Such an action is free when the weights are coprime with p and in that case we have a lens space obtained as the quotient of S^{2n+1} by the action of \mathbb{Z}_p . We denote such a lens space by $L_p^{2n+1}(\ell_0, \ell_1, \dots, \ell_n)$.

Note that different sequences of weights might produce diffeomorphic lens spaces, for instance by permuting the weights, adding a multiple of p to some weights, multiplying every weight by some k coprime with p and changing the sign of some weights. Moreover these are the only possibilities leading to diffeomorphic lens spaces. Clearly ψ preserves ξ_{std} and therefore $L_p^{2n+1}(\ell_0, \dots, \ell_n)$ carries the induced contact structure ξ . Permuting the weights, adding a multiple of p to some weights, multiplying every weight by some k coprime with p and changing the sign of *all* the weights lead to contactomorphic lens spaces. (As showed in [3], changing the sign of some weights can lead to lens spaces that are diffeomorphic but not contactomorphic.) *Throughout this work, we will choose the weights ℓ_0, \dots, ℓ_n such that $\ell_0 = 1$ and $-p/2 < \ell_i \leq p/2$ for every i .* These conditions determine the weights uniquely up to permutation in the last n weights.

Given a homotopy class $a \in \pi_1(L_p^{2n+1}(\ell_0, \dots, \ell_n))$, we will associate to it three key rational numbers in this work. Firstly, we have the lowest degree with non-trivial positive equivariant

symplectic homology associated to a , that is,

$$k_a := \min\{k \in \mathbb{Q}; \text{HC}_k^a(L_p^{2n+1}(\ell_0, \dots, \ell_n)) \neq 0\},$$

see Section 2 for the definition of equivariant symplectic homology and, in particular, its (fractional) grading. This number can be computed in terms of the weights in the following way. Let $j_a \in \{1, \dots, p\}$ be such that the deck transformation corresponding to a is ψ^{j_a} . Let $\ell_0^a, \ell_1^a, \dots, \ell_n^a$ be the corresponding *homotopy weights*, that is the unique integers satisfying $-p/2 < \ell_i^a \leq p/2$ for every i and

$$\psi^{j_a}(z_0, \dots, z_n) = \left(e^{\frac{2\pi\sqrt{-1}\ell_0^a}{p}} z_0, e^{\frac{2\pi\sqrt{-1}\ell_1^a}{p}} z_1, \dots, e^{\frac{2\pi\sqrt{-1}\ell_n^a}{p}} z_n \right).$$

Consider the number of positive/negative homotopy weights counted with multiplicity:

$$w_+^a = \#\{i; \ell_i^a > 0\} \quad \text{and} \quad w_-^a = \#\{i; \ell_i^a < 0\}.$$

Then one can show that

$$k_a = w_-^a - w_+^a + \frac{2\sum_i \ell_i^a}{p} + 1, \quad (1.2)$$

see Proposition 2.6.

Example 1.1. Consider the following illustrative examples:

- (1) Let a be a non-trivial homotopy class of $L_p^{2n+1}(1, \dots, 1)$. If $j_a \leq p/2$ then $\ell_i^a = j_a > 0$ for every i and therefore

$$k_a = -(n+1) + \frac{2j_a(n+1)}{p} + 1 = \frac{2j_a(n+1)}{p} - n.$$

If $j_a > p/2$ then $\ell_i^a = j_a - p < 0$ for every i and therefore

$$k_a = n+1 + \frac{2(j_a-p)(n+1)}{p} + 1 = \frac{2j_a(n+1)}{p} - n.$$

In particular, we have that $k_a \neq k_b$ whenever $a \neq b$.

- (2) Let a be a non-trivial homotopy class of $L_p^{2n+1}(1, -1, \dots, 1, -1)$ with $p > 2$ and n odd. (Note that $L_p^{2n+1}(1, -1, \dots, 1, -1) = L_p^{2n+1}(1, \dots, 1)$ when $p = 2$.) If $a^2 \neq 0$ we have that $w_-^a = w_+^a = (n+1)/2$ and $\sum_i \ell_i^a = 0$ and therefore $k_a = 1$. If $a^2 = 0$ then $w_+^a = n+1$, $w_-^a = 0$ and $\frac{2\sum_i \ell_i^a}{p} = n+1$ implying that $k_a = 1$ as well. Hence, $k_a = 1$ for every a .

Now, we will consider two rational numbers related to k_a and the *multiplicity* of the weights. More precisely, let $\bar{\ell}_1^a, \dots, \bar{\ell}_k^a$ be the absolute values of the homotopy weights $\ell_0^a, \dots, \ell_n^a$. Order $\bar{\ell}_i^a$ such that $\bar{\ell}_1^a < \bar{\ell}_2^a < \dots < \bar{\ell}_k^a$. Given $i \in \{1, \dots, k\}$ we define

$$\mu_i^a = \#\{j; \ell_j^a = \bar{\ell}_i^a \text{ and } \ell_j^a \neq p/2\}$$

and

$$\nu_i^a = \#\{j; \ell_j^a = -\bar{\ell}_i^a \text{ or we have that } \bar{\ell}_i^a = p/2 \text{ and } \ell_j^a = \bar{\ell}_i^a = p/2\}.$$

(Note that if $\bar{\ell}_i^a = p/2$ then $i = k$.) We also define

$$\tilde{\mu}_i^a = \#\{j; \ell_j^a = \bar{\ell}_i^a\} \quad \text{and} \quad \tilde{\nu}_i^a = \#\{j; \ell_j^a = -\bar{\ell}_i^a\}.$$

Set $\mu_0^a = \nu_0^a = \tilde{\nu}_0^a = 0$. Then one considers the numbers

$$h_a = \max \left\{ k_a - 1 + \sum_{i=0}^j \mu_i^a - \sum_{i=0}^j \nu_i^a; j \in \{0, \dots, k\} \right\}$$

and

$$\tilde{h}_a = \max \left\{ k_a - 1 + \sum_{i=1}^j \tilde{\mu}_i^a - \sum_{i=0}^{j-1} \tilde{\nu}_i^a; j \in \{1, \dots, k\} \right\}.$$

Note that $h_a \leq \tilde{h}_a$ since $\mu_i^a \leq \tilde{\mu}_i^a$ and $\nu_i^a \geq \tilde{\nu}_i^a$ for every i . Moreover, $h_a = \tilde{h}_a$ whenever p is odd.

Example 1.2. Let us compute these numbers in the previously considered examples:

- (1) Let a be a non-trivial homotopy class of $L_p^{2n+1}(1, \dots, 1)$. If $j_a < p/2$ then $\ell_i^a = j_a > 0$ for every i and therefore

$$h_a = \tilde{h}_a = k_a - 1 + n + 1 = \frac{2j_a(n+1)}{p}.$$

If $j_a = p/2$ then $\ell_i^a = p/2$ for every i and consequently

$$h_a = k_a - 1 = 0 \quad \text{and} \quad \tilde{h}_a = k_a + n = n + 1.$$

If $j_a > p/2$ then $\ell_i^a = j_a - p < 0$ for every i and therefore

$$h_a = \tilde{h}_a = k_a - 1 = \frac{2j_a(n+1)}{p} - (n+1).$$

- (2) Let a be a non-trivial homotopy class of $L_p^{2n+1}(1, -1, \dots, 1, -1)$ with $p > 2$ and n odd. As we saw in Example 1.1, $k_a = 1$ for every a . Note that we have only one absolute value $\bar{\ell}_1^a$, i.e., $k = 1$. Consider first the case that $a^2 \neq 0$. Then we have that $h_a = k_a - 1 = 0$ because $\mu_1^a = \nu_1^a = (n+1)/2$. Since $k = 1$ and $\tilde{\mu}_1^a = (n+1)/2$, we have that $\tilde{h}_a = (n+1)/2$. Now, consider the case where $a^2 = 0$. Then $\mu_1^a = 0$ and $\nu_1^a = n+1$ and therefore $h_a = k_a - 1 = 0$. On the other hand, $\tilde{\mu}_1^a = n+1$ and $\tilde{\nu}_1^a = 0$ and consequently $\tilde{h}_a = n+1$.

Remark 1.3. Note that the identity $\tilde{h}_a = k_a + n$ (resp $h_a = k_a + n$) actually holds for every non-trivial homotopy class a in $L_p^{2n+1}(\ell_0, \dots, \ell_n)$ such that $\ell_i^a > 0$ (resp. $\ell_i^a > 0$ and $\ell_i^a \neq p/2$) for every i .

Recall that a periodic orbit is called *hyperbolic* if every eigenvalue of its linearized Poincaré map has modulus different from one. On the other hand, it is called *elliptic* if every eigenvalue of its linearized Poincaré map has modulus one. Our first result establishes new dynamical implications of the (not necessarily strict) convexity of contact forms on the standard contact sphere under the presence of symmetries.

Theorem 1.4. *Let α be a convex (resp. strictly convex) contact form on a lens space $L_p^{2n+1}(\ell_0, \dots, \ell_n)$ and γ a closed Reeb orbit of α with non-trivial homotopy class a . Then the following assertions hold:*

- (1) $\mu(\gamma) \geq k_a$;
- (2) if $\mu(\gamma) < h_a$ (resp. $\mu(\gamma) < \tilde{h}_a$) then γ is non-hyperbolic;
- (3) if $\ell_i^a > 0$ and $\ell_i^a \neq p/2$ (resp. $\ell_i^a > 0$) for every i and $\mu(\gamma) = k_a$ then γ is elliptic.

Remark 1.5. We have that $k_0 = n + 2$. Therefore, when $a = 0$ the inequality $\mu(\gamma) \geq k_a$ means precisely dynamical convexity. Thus, the first assertion is a generalization of dynamical convexity for periodic orbits with non-trivial homotopy class. In fact, a notion of dynamical convexity for general contact manifolds was introduced in [2] and the first assertion states that α is a -dynamically convex in the terminology of [2]. (Note that to achieve the inequality $\mu(\gamma) \geq k_0$ for contractible orbits we need strict convexity while the lower bound $\mu(\gamma) \geq k_a$ for non-contractible orbits holds without assuming *strict* convexity.) The second and third assertions have no counterparts for contractible orbits.

Theorem 1.6. *Theorem 1.4 is sharp. More precisely, we have the following:*

- (1) *Given any convex contact form α on $L_p^{2n+1}(\ell_1, \dots, \ell_n)$ and a homotopy class a we must have at least one periodic orbit γ with homotopy class a such that $\mu(\gamma) = k_a$.*
- (2) *Given any integers $n \geq 1$ and $p \geq 2$ there exists a strictly convex contact form α on $L_p^{2n+1}(1, \dots, 1)$ and a hyperbolic closed Reeb orbit γ of α with non-trivial homotopy class a satisfying $\mu(\gamma) = h_a = \tilde{h}_a$.*
- (3) *There exists a strictly convex contact form α on $L_4^3(1, 1)$ and a hyperbolic closed Reeb orbit γ of α with non-trivial homotopy class a such that $\mu(\gamma) = k_a + 1$ and a satisfies $\ell_i^a > 0$ and $\ell_i^a \neq p/2$ for every i .*

Remark 1.7. By Proposition 2.2, the contact structure ξ on $L_4^3(1, 1)$ satisfies the following. The smallest positive integer N such that $Nc_1(\xi) = 0$ is equal to 2. Therefore, by (2.7) we have that the indexes of the periodic orbits are integers. Hence, $\mu(\gamma) > k_a$ if and only if $\mu(\gamma) \geq k_a + 1$.

Theorems 1.4 and 1.6 have the following straightforward consequences on the existence of symmetric elliptic and non-hyperbolic closed orbits of contact forms on $(S^{2n+1}, \xi_{\text{std}})$ which improves previous results due to Arnaud [5] and Liu, Wang and Zhang [26]. Let β be a contact form on S^{2n+1} invariant under the \mathbb{Z}_p -action generated by (1.1). A closed orbit γ of β is called symmetric if $\psi(\gamma(\mathbb{R})) = \gamma(\mathbb{R})$. Note that, the simple symmetric periodic orbits of β are in bijection with the simple periodic orbits of α whose homotopy classes are generators of $\pi_1(L_p^{2n+1}(\ell_0, \dots, \ell_n))$, where α is the contact form on $L_p^{2n+1}(\ell_0, \dots, \ell_n)$ whose lift to S^{2n+1} is β .

Corollary 1.8. *Let α be a convex (resp. strictly convex) contact form on $L_p^{2n+1}(\ell_0, \dots, \ell_n)$ with $p \geq 2$. Assume that $\ell_i > 0$ and $\ell_i \neq p/2$ (resp. $\ell_i > 0$) for every i . Then α carries at least one elliptic closed orbit whose homotopy class is a generator of $\pi_1(L_p^{2n+1}(\ell_0, \dots, \ell_n))$.*

Corollary 1.9. *Let α be a convex (resp. strictly convex) contact form on $L_p^{2n+1}(\ell_0, \dots, \ell_n)$ with $p \geq 2$. Assume that $\#\{i; \ell_i > 0 \text{ and } \ell_i \neq p/2\} - \#\{i; \ell_i < 0 \text{ or } \ell_i = p/2\} \geq 2$ (resp. $\#\{i; \ell_i > 0\} - \#\{i; \ell_i < 0\} \geq 2$) for every i . Then α carries at least one non-hyperbolic closed orbit whose homotopy class is a generator of $\pi_1(L_p^{2n+1}(\ell_0, \dots, \ell_n))$.*

Indeed, consider the generator a of $\pi_1(L_p^{2n+1}(\ell_0, \dots, \ell_n))$ whose homotopy weights are equal to ℓ_0, \dots, ℓ_n . We have from the first assertion of Theorem 1.6 that α must carry a closed orbit γ with homotopy class a such that $\mu(\gamma) = k_a$. Then, under the hypotheses of Corollary 1.8, by the third assertion of Theorem 1.4, we conclude that γ has to be elliptic. On the other hand, under the hypotheses of Corollary 1.9, we have that $h_a \geq k_a + 1$ (resp. $\tilde{h}_a \geq k_a + 1$) and therefore, by the second assertion of Theorem 1.4, we conclude that γ has to be non-hyperbolic.

Heuristically, we can look at Theorem 1.4 using the following analogy with Floer homology. Given a closed symplectic aspherical manifold M^{2n} we have that $\mathrm{HF}_*(M) \cong \mathrm{H}_{*+n}(M)$. Consider a C^2 -small non-degenerate autonomous Hamiltonian $H : M \rightarrow \mathbb{R}$. Then the 1-periodic orbits of H correspond to the critical points of H . Given a critical point p , let γ_p be the corresponding 1-periodic orbit of H and note that $\mu(\gamma_p) = \mu_{\mathrm{Morse}}(p) - n$, where $\mu_{\mathrm{Morse}}(p)$ is the Morse index of p . It is easy to see that if $\mu(\gamma_p) = -n$ then it is elliptic and if $\mu(\gamma_p) < 0$ then it must be non-hyperbolic (actually, it must be non-hyperbolic whenever $\mu(\gamma_p) \neq 0$). Thus, if the index of γ_p is the lowest one then γ_p must be elliptic and if the index of γ_p is lower than the “middle degree” then it has to be non-hyperbolic.

We say that a homotopy class $a \in \pi_1(L_p^{2n+1}(\ell_0, \dots, \ell_n))$ is *positive* (resp. *strictly positive*) if the corresponding homotopy weights satisfy $\ell_i^a > 0$ for every i (resp. $\ell_i^a > 0$ and $\ell_i^a \neq p/2$ for every i). (Equivalently, a is positive (resp. strictly positive) if the imaginary part of $e^{2\pi\sqrt{-1}\ell_i^a/p}$ is non-negative (resp. strictly positive) for every i .) Note that if a is positive (resp. strictly positive) then $\tilde{h}_a = k_a + n$ (resp. $h_a = k_a + n$) so that the previous heuristic analogy is more enlightening for these homotopy classes. A natural question is the existence of positive or strictly positive homotopy classes. Theorems 1.4 and 1.6 yield the following corollary, proved in Section 4, which addresses this question.

Corollary 1.10. *Let α be a convex (resp. strictly convex) contact form on a lens space and a be a strictly positive (resp. positive) homotopy class. Then α carries an elliptic periodic orbit with homotopy class a . In particular, the following assertions hold:*

- (1) *For a 3-dimensional lens space $L(p, q) = L_p^3(1, q)$, if $q \neq -1$ then $L(q, p)$ carries a positive homotopy class a . Therefore, every strictly convex contact form on such lens space has an elliptic closed orbit with homotopy class a .*
- (2) *For $L(p, q)$, if $q = -1$ then $L(q, p)$ carries a positive homotopy class a if p is even and it does not carry a positive homotopy class if p is odd. Therefore, every strictly convex contact form on such lens space has an elliptic closed orbit with homotopy class a whenever p is even.*
- (3) *For a higher dimensional lens space $L_p^{2n+1}(\ell_0, \dots, \ell_n)$, if the weights assume only two values $\ell_0 = 1$ and $\ell_1 = q$ with $q \neq -1$ then $L_p^{2n+1}(\ell_0, \dots, \ell_n)$ carries a positive homotopy class a . If $q = -1$ and p is even then $L_p^{2n+1}(\ell_0, \dots, \ell_n)$ carries a positive homotopy class a as well. Therefore, every strictly convex contact form on such lens space has an elliptic closed orbit with homotopy class a .*

Furthermore, if γ is a closed orbit of α with strictly positive (resp. positive) homotopy class a such that $\mu(\gamma) < k_a + n$ then γ must be non-hyperbolic.

Remark 1.11. The same reasoning shows the following. Let α be a convex (resp. strictly convex) contact form on a lens space and a be a homotopy class such that

$$\#\{i; \ell_i^a > 0 \text{ and } \ell_i^a \neq p/2\} - \#\{i; \ell_i^a < 0 \text{ or } \ell_i^a = p/2\} \geq 2 \quad (\text{resp. } \#\{i; \ell_i^a > 0\} - \#\{i; \ell_i^a < 0\} \geq 2).$$

Then α carries a non-hyperbolic closed orbit with homotopy class a . Note that, when $n = 1$, this condition is equivalent to a being strictly positive (resp. positive) (which is consistent with the fact that in dimension three a periodic orbit is elliptic if and only if it is non-hyperbolic).

Remark 1.12. The existence of an elliptic closed orbit when p is even in the corollary above also follows from the results in [2, 13] since in this case the lift of the contact form to S^{2n+1} is invariant by the antipodal map.

Another consequence of Theorem 1.4 is the following immediate corollary.

Corollary 1.13. *Let α be a strictly convex contact form on $\mathbb{R}P^{2n+1}$. Then every closed Reeb orbit γ of α satisfying $\mu(\gamma) < n + 1$ is non-hyperbolic.*

If $n = 1$ this result readily follows from the dynamical convexity of α and the multiplicative property of the index for hyperbolic periodic orbits: if γ is hyperbolic then $\mu(\gamma^k) = k\mu(\gamma)$ for every k ; hence, if $\mu(\gamma) \leq 1$ then $\mu(\gamma^2) \leq 2$, contradicting the dynamical convexity (note that the contact structure on $\mathbb{R}P^3$ has vanishing first Chern class and therefore the index is an integer so that $\mu(\gamma) < 2$ if and only if $\mu(\gamma) \leq 1$). However, there is no reason why this is true in higher dimensions. Indeed, the next result shows that the assumption that α is convex in Theorem 1.4 cannot be relaxed to the condition that α is dynamically convex at all.

Theorem 1.14. *The following assertions hold:*

- (1) *Consider integers $n \geq 1$ and $p \geq 3$. If $n = 1$ (resp. $n = 2$), assume furthermore that $p \geq 5$ (resp. $p \geq 4$). Then there exists a dynamically convex contact form α on $L_p^{2n+1}(1, \dots, 1)$ carrying a closed Reeb orbit with non-trivial homotopy class a such that $\mu(\gamma) < k_a$.*
- (2) *There exists a dynamically convex contact form α on $L_{11}^5(1, 1, 1)$ and a hyperbolic closed Reeb orbit γ of α with non-trivial homotopy class a such that $\ell_i^a > 0$ for every i and $\mu(\gamma) = k_a < h_a$.*

Remark 1.15. Note that the second assertion shows that both the second and third assertions of Theorem 1.4 do not hold assuming that α is dynamically convex. The first Chern class of the contact structure on $L_{11}^5(1, 1, 1)$ does not vanish. However, we can construct examples with vanishing first Chern class for which the second assertion of Theorem 1.4 does not hold; see Remarks 5.10, 5.14, 5.19, 5.20 and 5.21.

Remark 1.16. The closed orbit γ in the first assertion is non-degenerate. Therefore, every contact form $\tilde{\alpha}$ C^2 -close to α carries a closed Reeb orbit $\tilde{\gamma}$ with non-trivial homotopy class a such that $\mu(\tilde{\gamma}) < k_a$. Clearly, a similar stability statement also holds for the contact form in the second assertion: every contact form $\tilde{\alpha}$ C^2 -close to α carries a hyperbolic closed Reeb orbit $\tilde{\gamma}$ with non-trivial homotopy class a such that $\ell_i^a > 0$ for every i and $\mu(\tilde{\gamma}) = k_a < h_a$. It shows a contrast between these examples and those obtained in [19] where the degeneracy of the periodic orbits plays a crucial role.

Remark 1.17. The contact form α in both assertions can be chosen arbitrarily C^1 -close to a convex contact form; see Remarks 5.3 and 5.11.

Note that, by the invariance of equivariant symplectic homology, if $\phi : L_p^{2n+1}(\ell_0, \dots, \ell_n) \leftarrow$ is a contactomorphism then $k_a = k_{\phi_*a}$. In particular, if $k_a \neq k_b$ whenever $a \neq b$ then ϕ acts trivially on $\pi_1(L_p^{2n+1}(\ell_0, \dots, \ell_n))$. Hence, in this case, all the assertions in Theorem 1.4 are invariant by ϕ . Therefore, since this property holds for $L_p^{2n+1}(1, \dots, 1)$ (see Example 1.1), Theorem 1.14 furnish new examples of dynamically convex contact forms β on spheres that are not contactomorphic to convex contact forms via contactomorphisms that commute with the corresponding symmetry.

The fact that we do not assume *strict* convexity implies more than this: let $S \subset \text{Cont}(S^{2n+1})$ be the subset of contactomorphisms that commute with the corresponding \mathbb{Z}_p -action. Then there exists a C^2 -neighborhood U of S such that β is not equivalent to a convex contact form via any contactomorphism $\varphi \in U$. Indeed, if there exists a sequence $\varphi_i \xrightarrow{C^2} \bar{\varphi} \in S$ such that $\varphi_i^*\beta$ is convex then so is $\bar{\varphi}^*\beta$, furnishing a contradiction. However, we actually have this property for a C^1 -neighborhood of S as the following theorem shows.

Theorem 1.18. *Let α be one of the contact forms furnished by Theorem 1.14 and consider its lift β to S^{2n+1} . Let $S \subset \text{Cont}(S^{2n+1})$ be the subset of contactomorphisms that commute with the corresponding \mathbb{Z}_p -action. Then there exists a C^1 -neighborhood U of S such that β is not equivalent to a convex contact form via any contactomorphism $\varphi \in U$.*

Remark 1.19. Very recently, Chaidez and Edtmair [10] showed examples of dynamically convex contact forms on S^3 that are not equivalent to a convex contact form via any contactomorphism. However, their methods work only in dimension three.

Another application of Theorem 1.4, obtained using Floer homology techniques, is the following result on the multiplicity of symmetric non-hyperbolic periodic orbits on convex spheres. Before we state it, we need to introduce some definitions. Let α be a contact form on $L_p^{2n+1}(\ell_0, \dots, \ell_n)$ and β its lift to S^{2n+1} . Let $H_\beta : \mathbb{R}^{2n+2} \rightarrow \mathbb{R}$ be the unique homogeneous of degree two Hamiltonian such that $\Sigma_\beta = H_\beta^{-1}(1)$. Note that β is convex if and only if so is H_β (that is, the Hessian of H_β is positive semi-definite at every point). Given real numbers $0 < r \leq R$ we say that α is (r, R) -pinched if $R^{-2}\|x\|^2 \leq H_\beta(x) \leq r^{-2}\|x\|^2$ for every $x \in \Sigma_\beta$. We say that α is (r, R) -H-pinched if $R^{-2}\|v\|^2 \leq \text{Hess } H_\beta(x)(v, v) \leq r^{-2}\|v\|^2$ for every $x \in \Sigma_\beta$ and $v \in \mathbb{R}^{2n+2}$. (Here ‘‘H’’ stands for the Hessian.) Using the homogeneity of H_β , it is easy to see that if β is (r, R) -H-pinched then it is $(\sqrt{2}r, \sqrt{2}R)$ -pinched.

Theorem 1.20. *Let $n \geq 1$ and $p \geq 2$ be integers and $0 < r \leq R$ be real numbers such that $\frac{R}{r} < \sqrt{p+1}$. Let α be an (r, R) -H-pinched contact form on $L_p^{2n+1}(1, \dots, 1)$. Denote by a the generator of $\pi_1(L_p^{2n+1}(1, \dots, 1))$ such that $\ell_i^a = 1$ for every i . Suppose that the periodic orbits of α with homotopy class a are isolated. Then α carries at least $\lfloor \frac{n+1}{2} \rfloor$ geometrically distinct non-hyperbolic closed Reeb orbits with homotopy class a .*

Finally, we prove the following result on the multiplicity of symmetric (not necessarily non-hyperbolic) closed orbits for contact forms satisfying a weaker pinching condition. It improves results from [4, 22] and does not use Theorem 1.4; the convexity of the contact form is used to obtain a lower bound for the period of closed orbits of convex contact forms due to Croke and Weinstein [12] (which is also used in the proof of Theorem 1.20).

Theorem 1.21. *Let $n \geq 1$ and $p \geq 1$ be integers and $0 < r \leq R$ be real numbers such that $\frac{R}{r} < \sqrt{p+1}$. Let α be a convex (r, R) -pinched contact form on $L_p^{2n+1}(1, \dots, 1)$. Denote by a the generator of $\pi_1(L_p^{2n+1}(1, \dots, 1))$ such that $\ell_i^a = 1$ for every i . Then α carries at least $n+1$ geometrically distinct closed Reeb orbits with homotopy class a .*

Note that we allow $p = 1$ in the previous theorem. This case corresponds to a classical result due to Ekeland and Lasry [15].

1.3. Organization of the paper. The rest of the paper is organized as follows. The background on equivariant symplectic homology and Lusternik-Schnirelmann theory in Floer homology necessary for this work is presented in Sections 2.1 and 2.2 respectively. Its (fractional) grading is discussed in Section 2.3. Section 2.4 contains index computations relevant for this work and shows how to deal with the non-vanishing of the first Chern class in lens spaces. The tools from index theory are introduced in Section 2.5. Theorem 1.4 is proved in Section 3 and its Corollary 1.10 is proved in Section 4. Theorem 1.6, that establishes the sharpness of Theorem 1.4, is proved in Section 6. Its proof appears after the proof of Theorem 1.14, presented in Section 5, since it uses several ingredients from that. Finally, Sections 8 and 9 are devoted to the proof of our multiplicity results, namely, Theorems 1.20 and 1.21 respectively.

1.4. Conventions. Throughout this work, we will use the convention that the natural numbers are given by the positive integers. Given a symplectic manifold (M, ω) and a Hamiltonian $H_t : M \rightarrow \mathbb{R}$, we take Hamilton's equation to be $i_{X_{H_t}} \omega = -dH_t$. A compatible almost complex structure J is defined by the condition that $\omega(\cdot, J\cdot)$ is a Riemannian metric. Throughout this work, the Conley-Zehnder index μ is normalized so that when Q is a small positive definite quadratic form the path $\Gamma : [0, 1] \rightarrow \mathrm{Sp}(2n)$ generated by Q and given by $\Gamma(t) = \exp(tJQ)$ has $\mu(\Gamma) = n$. We also take the canonical symplectic form on \mathbb{R}^{2n} to be $\sum dq_i \wedge dp_i$. For degenerate paths, the Conley-Zehnder index μ is defined as the lower semi-continuous extension of the Conley-Zehnder index from the paths with non-degenerate endpoint. More precisely,

$$\mu(\Gamma) = \liminf_{\tilde{\Gamma} \rightarrow \Gamma} \mu(\tilde{\Gamma}),$$

where $\tilde{\Gamma}$ is a small perturbation of Γ with non-degenerate endpoint. These conventions are consistent with the ones used in [19].

1.5. Acknowledgements. We are grateful to Viktor Ginzburg and Umberto Hryniewicz for useful comments on a preliminary version of this paper.

2. EQUIVARIANT SYMPLECTIC HOMOLOGY AND INDEX THEORY

2.1. Equivariant symplectic homology. Let (M^{2n+1}, ξ) be a contact manifold endowed with a strong symplectic filling given by a Liouville domain W such that $c_1(TW)|_{H_2(W, \mathbb{R})} = 0$. The positive equivariant symplectic homology $\mathrm{SH}_*^{S^1, +}(W)$ is a symplectic invariant introduced by Viterbo [32] and developed by Bourgeois and Oancea [6, 7, 8, 9].

It turns out that the positive equivariant symplectic homology can be obtained as the homology of a chain complex $\mathrm{CC}_*(\alpha)$ with rational coefficients generated by the good closed Reeb orbits of a non-degenerate contact form α on M . This complex is filtered by the action and graded by the Conley-Zehnder index; see [18, Proposition 3.3] and Section 2.3 for a discussion concerning the grading and good orbits. The differential in the complex $\mathrm{CC}_*(\alpha)$, but not its homology, depends on several auxiliary choices, and the nature of the differential is not essential for our purposes. The complex $\mathrm{CC}_*(\alpha)$ is functorial in α in the sense that a symplectic cobordism equipped with a suitable extra structure gives rise to a map of complexes. For the sake of brevity and to emphasize the obvious analogy with contact homology, we denote the homology of $\mathrm{CC}_*(\alpha)$ by $\mathrm{HC}_*(M)$ rather than $\mathrm{SH}_*^{S^1, +}(W)$. Furthermore, once we fix a free homotopy class of loops in W , the part of $\mathrm{CC}_*(\alpha)$ generated by closed Reeb orbits in that class is a subcomplex. As a consequence, the entire complex $\mathrm{CC}_*(\alpha)$ breaks down into a direct sum of such subcomplexes indexed by free homotopy classes of loops in W .

A remarkable observation by Bourgeois and Oancea in [9, Section 4.1.2] is that under suitable additional assumptions on the indices of closed Reeb orbits the positive equivariant symplectic homology is defined even when M does not have a symplectic filling and therefore is a contact invariant. To be more precise, we assume that $c_1(\xi)|_{H_2(M, \mathbb{R})} = 0$ and that M admits a non-degenerate contact form α such that all of its closed *contractible* Reeb orbits have Conley-Zehnder index strictly greater than $3 - n$. Furthermore, under this assumption once again the positive equivariant symplectic homology of M can be described as the homology of a complex $\mathrm{CC}_*(\alpha)$ generated by good closed Reeb orbits of α , graded by the Conley-Zehnder index and filtered by the action. The complex breaks down into the direct sum of subcomplexes indexed by free homotopy classes of loops in M . We will use the notation $\mathrm{HC}_*^a(M)$ to denote

the homology of the complex generated by the orbits with free homotopy class a . Given a non-degenerate contact form α and numbers $0 < T_1 < T_2 \leq \infty$ we denote by $\mathrm{HC}_*^{a,(T_1,T_2)}(\alpha)$ the equivariant symplectic homology of α with free homotopy class a and action window (T_1, T_2) . When α is degenerate and both T_1 and T_2 are not in the action spectrum of α we define $\mathrm{HC}_*^{a,(T_1,T_2)}(\alpha)$ as $\mathrm{HC}_*^{a,(T_1,T_2)}(\bar{\alpha})$ for some small non-degenerate perturbation $\bar{\alpha}$ of α . (Recall that the action spectrum of α is given by $\mathcal{A}(\alpha) = \{\int_\gamma \alpha; \gamma \text{ is a closed Reeb orbit of } \alpha\}$.) Given $T \in (0, \infty]$ we denote by $\mathrm{HC}_*^{a,T}(\alpha)$ the filtered equivariant symplectic homology $\mathrm{HC}_*^{a,(\epsilon,T)}(\alpha)$ for some $\epsilon > 0$ sufficiently small such that $\epsilon < \min\{T; T \in \mathcal{A}(\alpha)\}$.

The functoriality of $\mathrm{CC}_*(\alpha)$ respects the action and homotopy filtrations and therefore the following holds. Given contact forms α_0, α_1 and α_2 , we say that $\alpha_0 < \alpha_1 < \alpha_2$ if $\alpha_1 = f_1\alpha_0$ and $\alpha_2 = f_2\alpha_0$ for functions $f_i : M \rightarrow \mathbb{R}$ ($i = 1, 2$) such that $f_1(x) > 1$ and $f_1(x) < f_2(x)$ for every $x \in M$. Then, given $0 < T_1 < T_2$ not in the action spectrum of α_i ($i = 0, 1, 2$), we have continuation maps $\phi_{\alpha_i, \alpha_j} : \mathrm{HC}_*^{a,(T_1,T_2)}(\alpha_i) \rightarrow \mathrm{HC}_*^{a,(T_1,T_2)}(\alpha_j)$ (with $i, j \in \{0, 1, 2\}$ such that $i > j$) that fit into the commutative diagram

$$\begin{array}{ccc} \mathrm{HC}_*^{a,(T_1,T_2)}(\alpha_2) & \xrightarrow{\phi_{\alpha_2, \alpha_0}} & \mathrm{HC}_*^{a,(T_1,T_2)}(\alpha_0) \\ & \searrow \phi_{\alpha_2, \alpha_1} & \nearrow \phi_{\alpha_1, \alpha_0} \\ & \mathrm{HC}_*^{a,(T_1,T_2)}(\alpha_1) & \end{array} \quad (2.1)$$

These continuation maps have the following property that will be useful in this work. Let $\alpha_t, t \in [0, 1]$, be a smooth family of contact forms such that $\alpha_t < \alpha_{t'}$ whenever $t < t'$. Suppose that there exists $T \in \mathbb{R}$ such that $T \notin \cup_{t \in [0,1]} \mathcal{A}^a(\alpha_t)$, where

$$\mathcal{A}^a(\alpha_t) = \left\{ \int_\gamma \alpha_t; \gamma \text{ is a closed Reeb orbit of } \alpha_t \text{ with homotopy class } a \right\}.$$

Then

$$\phi_{\alpha_1, \alpha_0} : \mathrm{HC}_*^{a,T}(\alpha_1) \rightarrow \mathrm{HC}_*^{a,T}(\alpha_0)$$

is an isomorphism. Hence, by the diagram (2.1), we conclude that if a contact form α satisfies $\alpha_0 < \alpha < \alpha_1$ and $T \notin \mathcal{A}^a(\alpha)$ then the corresponding continuation map

$$\phi_{\alpha_1, \alpha} : \mathrm{HC}_*^{a,T}(\alpha_1) \rightarrow \mathrm{HC}_*^{a,T}(\alpha)$$

is injective.

Now, suppose that (M, ξ) is given by a lens space $L_p^{2n+1}(\ell_0, \dots, \ell_n)$ with the contact structure ξ induced by $(S^{2n+1}, \xi_{\mathrm{std}})$. Then $c_1(\xi)|_{H_2(M, \mathbb{R})} = 0$, because $H^2(M, \mathbb{Z})$ is torsion, and every contractible closed orbit of a dynamically convex contact form α on $L_p^{2n+1}(\ell_0, \dots, \ell_n)$ has Conley-Zehnder index strictly greater than $3-n$ (recall that a contact form on $L_p^{2n+1}(\ell_0, \dots, \ell_n)$ is dynamically convex if every contractible closed orbit has index bigger than or equal to $n+2$). The existence of dynamically convex contact forms follows from the fact that every convex contact form is dynamically convex. Thus, $\mathrm{HC}_*^a(L_p^{2n+1}(\ell_0, \dots, \ell_n))$ is a contact invariant that can be obtained as the homology of a chain complex $\mathrm{CC}_*^a(\alpha)$ with rational coefficients generated by the good closed Reeb orbits of α with homotopy class a (note that $\pi_1(L_p^{2n+1}(\ell_0, \dots, \ell_n))$ is abelian). Moreover, $\mathrm{HC}_*^a(L_p^{2n+1}(\ell_0, \dots, \ell_n))$ have continuation maps that satisfy the diagram (2.1).

Finally, let us briefly recall the definition of equivariant local symplectic homology. Given an isolated (possibly degenerate) closed orbit γ of α we have its equivariant local symplectic homology $\mathrm{HC}_*(\gamma)$ [18, 21]. It is supported in $[\mu(\gamma), \mu(\gamma) + \nu(\gamma)]$ (i.e. $\mathrm{HC}_k(\gamma) = 0$ for every $k \notin [\mu(\gamma), \mu(\gamma) + \nu(\gamma)]$), where $\nu(\gamma)$ is the nullity of γ , i.e., the geometric multiplicity of the eigenvalue 1 of the linearized Poincaré map. If α carries finitely many simple closed orbits with free homotopy class a then if $\mathrm{HC}_k^a(M) \neq 0$ there exists a closed orbit γ with free homotopy class a such that $\mathrm{HC}_k(\gamma) \neq 0$.

2.2. Lusternik-Schnirelmann theory. In what follows, we will explain briefly the results from Lusternik-Schnirelmann theory in Floer homology necessary for this work. We refer to [18] for details.

Let $M = L_p^{2n+1}(\ell_0, \dots, \ell_n)$ and $a \in \pi_1(M)$. Let α be a contact form on M such that every closed orbit with homotopy class a is isolated and $T \in (0, \infty] \setminus \mathcal{A}(\alpha)$. Given a non-trivial element $w \in \mathrm{HC}_k^{a,T}(\alpha)$ we have a spectral invariant given by

$$c_w(\alpha) = \inf\{T' \in (0, T) \setminus \mathcal{A}(\alpha); w \in \mathrm{Im}(i^{a,T'})\}$$

where $i^{a,T'} : \mathrm{HC}_*^{a,T'}(\alpha) \rightarrow \mathrm{HC}_*^{a,T}(\alpha)$ is the map induced in the homology by the inclusion of the complexes. It turns out that there exists a periodic orbit γ with action $c_w(\alpha)$ and free homotopy class a such that $\mathrm{HC}_k(\gamma) \neq 0$; c.f. [18, Corollary 3.9].

There is a shift operator $D : \mathrm{HC}_*^a(\alpha) \rightarrow \mathrm{HC}_{*-2}^a(\alpha)$ introduced in [8] which respects the action filtration and satisfies the property

$$c_w(\alpha) > c_{D(w)}(\alpha) \tag{2.2}$$

see [18, Theorem 1.1]. The shift operator and the spectral invariants are functorial with respect to the continuation maps in the sense that given contact forms $\alpha_0 < \alpha_1$ we have that D commutes with $\phi_{\alpha_1, \alpha_0} : \mathrm{HC}_*^{a,T}(\alpha_1) \rightarrow \mathrm{HC}_*^{a,T}(\alpha_0)$ and $c_{\phi_{\alpha_1, \alpha_0}(w)}(\alpha_0) \leq c_w(\alpha_1)$ for every $w \in \mathrm{HC}_*^{a,T}(\alpha_1)$, see [18, Proposition 3.1].

Suppose now that M admits a contact form α_0 whose Reeb flow generates a free circle action such that a is the homotopy class of the simple orbits of α_0 . Let $T \in (0, \infty] \setminus \mathcal{A}(\alpha_0)$. Let $B = M/S^1$ and assume further that $\mathrm{H}_*(B; \mathbb{Q})$ vanishes in odd degrees. Then the shift operator D can be (partially) computed in the following way. Let $\Delta : \mathrm{H}_*(B; \mathbb{Q}) \rightarrow \mathrm{H}_{*-2}(B; \mathbb{Q})$ be the shift operator given by the Gysin exact sequence associated to the S^1 -bundle $S^1 \rightarrow M \rightarrow B$

$$\dots \rightarrow \mathrm{H}_*(M; \mathbb{Q}) \rightarrow \mathrm{H}_*(B; \mathbb{Q}) \xrightarrow{\Delta} \mathrm{H}_{*-2}(B; \mathbb{Q}) \rightarrow \mathrm{H}_{*-1}(M; \mathbb{Q}) \rightarrow \dots$$

Suppose that there exist non-zero elements $v_i \in \mathrm{H}_{2i}(B; \mathbb{Q})$, $i = 0, \dots, n$, such that $\Delta(v_{i+1}) = v_i$ for every $i \in \{0, \dots, n-1\}$.

Remark 2.1. In this work, we will use the techniques presented in this section in the particular case where $M = L_p^{2n+1}(1, \dots, 1)$ is endowed with the contact form α_0 that generates the obvious free circle action (induced by the Hopf fibration) whose orbit space B is $\mathbb{C}P^n$. In this case, clearly $\mathrm{H}_*(B; \mathbb{Q})$ vanishes in odd degrees, $\mathrm{H}_{2i}(B; \mathbb{Q}) \cong \mathbb{Q}$ for every $i \in \{0, \dots, n\}$ and $\Delta : \mathrm{H}_{2i+2}(B; \mathbb{Q}) \rightarrow \mathrm{H}_{2i}(B; \mathbb{Q})$ is an isomorphism for every $i \in \{0, \dots, n-1\}$.

Let γ be a simple periodic orbit of α_0 and denote by $A(\gamma)$ the action of γ . A standard Morse-Bott computation shows that there exists $\epsilon > 0$ such that

$$\mathrm{HC}_*^{a, (A(\gamma^{(k-1)p+1}) - \epsilon, A(\gamma^{(k-1)p+1}) + \epsilon)}(\alpha_0) \cong \mathrm{H}_{*-\mu(\gamma^{(k-1)p+1})}(B; \mathbb{Q}).$$

for every $k \in \mathbb{N}$. It turns out that this isomorphism is equivariant with respect to D and $((k-1)p+1)\Delta$; see [18, Proposition 2.22]. (Note that $((k-1)p+1)\Delta$ is the map in the Gysin exact sequence of the S^1 -bundle induced by the $((k-1)p+1)$ -th iterate of the Reeb flow of α_0 .) From this (using our assumption that $H_*(B; \mathbb{Q})$ vanishes in odd degrees) we can conclude that

$$\mathrm{HC}_*^{a,T}(\alpha_0) \cong \bigoplus_{k \in \{j \in \mathbb{N}; A(\gamma^{(j-1)p+1}) < T\}} H_{*-\mu(\gamma^{(k-1)p+1})}(B; \mathbb{Q}) \quad (2.3)$$

and that (using our hypothesis on Δ) if $T > A(\gamma)$ then there exist non-zero elements $w_i \in \mathrm{HC}_{\mu(\gamma)+2i}^{a,T}(\alpha_0)$, $i = 0, \dots, n$, such that $Dw_{i+1} = w_i$ for every $i \in \{0, \dots, n-1\}$. (Note here that D respects the action filtration and therefore given non-zero elements w_i in the first summand $\mathrm{HC}_*^{a,(A(\gamma)-\epsilon, A(\gamma)+\epsilon)}(\alpha_0) \cong \mathrm{HC}_*^{a,(0, A(\gamma)+\epsilon)}(\alpha_0)$ such that $D^{A(\gamma)+\epsilon}w_{i+1} = w_i$, where $D^{A(\gamma)+\epsilon} : \mathrm{HC}_*^{a,(0, A(\gamma)+\epsilon)}(\alpha_0) \rightarrow \mathrm{HC}_*^{a,(0, A(\gamma)+\epsilon)}(\alpha_0)$ is the shift operator in the corresponding action window, then the corresponding elements w_i in $\mathrm{HC}_*^{a,T}(\alpha_0)$ also satisfy $Dw_{i+1} = w_i$.)

Taking $T = \infty$ (so that $\mathrm{HC}_*^{a,T}(\alpha_0) = \mathrm{HC}_*^a(M)$ does not depend on the contact form) we conclude in particular that

$$\mathrm{HC}_*^a(M) \cong \bigoplus_{k \in \mathbb{N}} H_{*-\mu(\gamma^{(k-1)p+1})}(B; \mathbb{Q}) \quad (2.4)$$

and that there exist non-zero elements $w_i \in \mathrm{HC}_{\mu(\gamma)+2i}^a(M)$, $i = 0, \dots, n$, such that $Dw_{i+1} = w_i$. From this and (2.2) we infer that there exists an injective map

$$\psi : \{0, \dots, n\} \rightarrow \mathcal{P}^a(\alpha), \quad (2.5)$$

called carrier map, where $\mathcal{P}^a(\alpha)$ is the set of closed orbit of α with homotopy class a (note the difference between α and α_0), such that if $\gamma_i = \psi(i)$ then $A(\gamma_i) = c_{w_i}(\alpha)$ and $\mathrm{HC}_{\mu(\gamma)+2i}(\gamma_i) \neq 0$; see [18].

Using the functoriality of D , we can refine this carrier map in the following way. Consider contact forms $\alpha < \alpha'$ and let $T \in (0, \infty] \setminus (\mathcal{A}(\alpha) \cup \mathcal{A}(\alpha'))$. Suppose that there exist non-zero elements $w_i \in \mathrm{HC}_{k_i}^{a,T}(\alpha')$, $i = 0, \dots, n$, such that $Dw_{i+1} = w_i$ for every $i \in \{0, \dots, n-1\}$ (for some k_i such that $k_i = k_{i+1} - 2$) and that the continuation map $\phi_{\alpha', \alpha} : \mathrm{HC}_*^{a,T}(\alpha') \rightarrow \mathrm{HC}_*^{a,T}(\alpha)$ is injective. Then there exists an action filtered injective carrier map

$$\psi^T : \{0, \dots, n\} \rightarrow \mathcal{P}^{a,T}(\alpha), \quad (2.6)$$

where $\mathcal{P}^{a,T}(\alpha)$ is the set of closed orbit of α with homotopy class a and period less than T , such that if $\gamma_i = \psi(i)$ then $A(\gamma_i) = c_{\phi_{\alpha', \alpha}(w_i)}(\alpha)$ and $\mathrm{HC}_{k_i}(\gamma_i) \neq 0$.

2.3. Grading. The grading of the positive equivariant symplectic homology is defined in [9] as follows. In what follows, let $M = L_p^{2n+1}(\ell_0, \dots, \ell_n)$ be endowed with the contact structure ξ induced from $(S^{2n+1}, \xi_{\mathrm{std}})$. Given a homotopy class $a \in \pi_1(M)$, choose a reference loop ψ^a in M and a symplectic trivialization of $(\psi^a)^*\xi$. When $a = 0$ we ask that both ψ^a and the trivialization are constant. Given a closed orbit $\gamma : S^1 \rightarrow M$ with homotopy class a and a homotopy between γ and ψ^a we have an induced trivialization of $\gamma^*\xi$. The assumption that $c_1(\xi)|_{H_2(M, \mathbb{R})} = 0$ implies that the homotopy class of this trivialization does not depend on the choice of the homotopy. Then the index of γ is the Conley-Zehnder index of the symplectic path given by the linearized Reeb flow along γ (restricted to the contact structure) with respect to the trivialization of $\gamma^*\xi$. Note that, for non-trivial homotopy classes, this grading depends on the choice of the reference loops ψ^a and of the trivializations of $(\psi^a)^*\xi$. In general, this grading has the following issue: this trivialization might be not *closed under iterations*,

that is, the trivialization induced on γ^j might be not homotopic with the j -th iterate of the trivialization over γ . This is a problem when we try to use index theory.

In order to fix this issue, we will define a *fractional* grading using sections of the determinant line bundle in the following way [30, 31]. Note that $c_1(\xi)$ is torsion and let N be the smallest positive integer such that $Nc_1(\xi) = 0$ (N can be easily computed from the weights ℓ_0, \dots, ℓ_n ; see Proposition 2.2). Then $(\Lambda_{\mathbb{C}}^n \xi)^{\otimes N}$ is a trivial line bundle. Choose a trivialization $\tau : (\Lambda_{\mathbb{C}}^n \xi)^{\otimes N} \rightarrow M \times \mathbb{C}$ which corresponds to a choice of a non-vanishing section \mathfrak{s} of $(\Lambda_{\mathbb{C}}^n \xi)^{\otimes N}$. The choice of this trivialization furnishes a unique way to symplectically trivialize $\bigoplus_1^N \xi$ along periodic orbits of α up to homotopy. As a matter of fact, given a periodic orbit γ , let $\Phi : \gamma^* \bigoplus_1^N \xi \rightarrow S^1 \times \mathbb{C}^{nN}$ be a trivialization of $\bigoplus_1^N \xi$ over γ as a Hermitian vector bundle such that its highest complex exterior power coincides with τ . This condition fixes the homotopy class of Φ : given any other such trivialization Ψ we have, for every $t \in S^1$, that $\Phi_t \circ \Psi_t^{-1} : \mathbb{C}^{nN} \rightarrow \mathbb{C}^{nN}$ has complex determinant equal to one and therefore the Maslov index of the symplectic path $t \mapsto \Phi_t \circ \Psi_t^{-1}$ vanishes, where $\Phi_t := \pi_2 \circ \Phi|_{\gamma^* \bigoplus_1^N \xi(t)}$ and $\Psi_t := \pi_2 \circ \Psi|_{\gamma^* \bigoplus_1^N \xi(t)}$ with $\pi_2 : S^1 \times \mathbb{C}^{nN} \rightarrow \mathbb{C}^{nN}$ being the projection onto the second factor; cf. [30]. Notice that this trivialization is closed under iterations, that is, the trivialization induced on γ^j coincides, up to homotopy, with the j -th iterate of the trivialization over γ .

Now, one can define the Conley-Zehnder index $\mu(\gamma; \mathfrak{s})$ of a closed orbit γ in the following way. By the previous discussion, \mathfrak{s} induces a unique up to homotopy symplectic trivialization $\Phi : \gamma^* \bigoplus_1^N \xi \rightarrow S^1 \times \mathbb{R}^{2nN}$. Using this trivialization, the linearized Reeb flow gives the symplectic path

$$\Gamma(t) = \Phi_t \circ \bigoplus_1^N d\phi_\alpha^t(\gamma(0))|_{\xi} \circ \Phi_0^{-1},$$

where ϕ_α^t is the Reeb flow of α . Then the Conley-Zehnder index is defined as

$$\mu(\gamma; \mathfrak{s}) = \frac{\mu(\Gamma)}{N}$$

where the Conley-Zehnder index of Γ is defined as the lowersemicontinuous extension of the usual Conley-Zehnder index for non-degenerate paths. It turns out that, since $H^1(M; \mathbb{Q}) = 0$, this index does not depend on the choice of \mathfrak{s} since every two such sections are homotopic; see [30, Lemma 4.3].

Note that this grading is fractional in general. Even though the idea of a fractional grading may seem unnatural at first, it can be thought of as a way of keeping track of some information about the homotopy classes of the orbits. Indeed, given two homotopic orbits we have that their index difference is an integer. As a matter of fact, fixed a homotopy class a , the gradings obtained using sections of $(\Lambda_{\mathbb{C}}^n \xi)^{\otimes N}$ and reference loops coincide up to a constant. To see this, choose a reference loop ψ^a and a trivialization of $(\psi^a)^* \xi$. It induces an obvious (product) trivialization Ψ^a of $(\psi^a)^* \bigoplus_1^N \xi$. Given a closed orbit γ with homotopy class a , we have the induced trivialization of $\gamma^* \bigoplus_1^N \xi$. Let $\mu(\gamma; \Psi^a)$ be the index of γ using this trivialization. Let Φ^a be a trivialization $(\psi^a)^* \bigoplus_1^N \xi$ induced by a non-vanishing section \mathfrak{s} of $(\Lambda_{\mathbb{C}}^n \xi)^{\otimes N}$. Then we have that

$$\mu(\gamma; \mathfrak{s}) - \mu(\gamma; \Psi^a) = c^a := \frac{2\mu_{\text{Maslov}}(\Phi_t^a \circ (\Psi_t^a)^{-1})}{N}. \quad (2.7)$$

Let $\bar{\gamma}$ be the underlying simple orbit of γ , that is, $\bar{\gamma}$ is a simple orbit and $\gamma = \bar{\gamma}^j$ for some $j \in \mathbb{N}$. Denote by \bar{a} the homotopy class of $\bar{\gamma}$ (so that $a = \bar{a}^j$). We say that γ is good if

$$\mu(\gamma; \Psi^a) - \mu(\bar{\gamma}; \Psi^{\bar{a}}) \in 2\mathbb{Z}.$$

Otherwise, it is called bad. From the previous discussion we have that γ is good if and only if

$$\mu(\gamma; \mathfrak{s}) - \mu(\bar{\gamma}; \mathfrak{s}) \in c^a - c^{\bar{a}} + 2\mathbb{Z}.$$

2.4. Index computations and non-vanishing of the first Chern class in lens spaces.

This section contains index computations relevant for this work and shows how to deal with the non-vanishing of the first Chern class in lens spaces. First, we have the following result regarding the computation of N in terms of the weights that define the lens space.

Proposition 2.2. *There is an isomorphism $H^2(L_p^{2n+1}(\ell_0, \ell_1, \dots, \ell_n); \mathbb{Z}) \cong \mathbb{Z}_p$ such that the first Chern class $c_1(\xi)$ is given by*

$$c_1(\xi) = \sum_{i=0}^n \ell_i \pmod{p}.$$

Hence,

$$m \cdot c_1(\xi) = 0 \Leftrightarrow m \cdot \sum_{i=0}^n \ell_i = 0 \pmod{p}.$$

In particular, we have that $N = \min\{m \in \mathbb{N}; m \sum_i \ell_i = 0 \pmod{p}\}$.

Proof. This is a particular case of Proposition 2.16 in [1] and its proof. The main points are the following:

- (i) The quotient map from S^{2n+1} to $L_p^{2n+1}(\ell_0, \ell_1, \dots, \ell_n)$ is a principal \mathbb{Z}_p -bundle and its classifying map $f : L_p^{2n+1}(\ell_0, \ell_1, \dots, \ell_n) \rightarrow B\mathbb{Z}_p$ induces an isomorphism $f^* : H^2(B\mathbb{Z}_p; \mathbb{Z}) \rightarrow H^2(L_p^{2n+1}(\ell_0, \ell_1, \dots, \ell_n); \mathbb{Z})$.
- (ii) The \mathbb{Z}_p -action on \mathbb{C}^{n+1} gives rise to an associated vector bundle $S^{2n+1} \times_{\mathbb{Z}_p} \mathbb{C}^{n+1}$ over $L_p^{2n+1}(\ell_0, \ell_1, \dots, \ell_n)$, and

$$c_1(\xi) = c_1(S^{2n+1} \times_{\mathbb{Z}_p} \mathbb{C}^{n+1}) = f^* c_1(E\mathbb{Z}_p \times_{\mathbb{Z}_p} \mathbb{C}^{n+1}).$$

- (iii) $H^2(B\mathbb{Z}_p; \mathbb{Z}) \cong \mathbb{Z}_p \cong$ character group of \mathbb{Z}_p and $c_1(E\mathbb{Z}_p \times_{\mathbb{Z}_p} \mathbb{C}^{n+1})$ is the sum of the characters that determine the \mathbb{Z}_p -action on \mathbb{C}^{n+1} .

□

Given $a \in \pi_1(L_p^{2n+1}(\ell_0, \dots, \ell_n))$, consider on $\mathbb{R}^{(2n+2)N} \simeq \mathbb{C}^{(n+1)N}$ the Hamiltonian

$$G_a(z_0, \dots, z_{n+(N-1)(n+1)}) = \frac{\pi}{p} \left(\sum_{j=0}^{N-1} \sum_{i=0}^n \ell_i^a \|z_{i+j(n+1)}\|^2 - \left(N \sum_{i=0}^n \ell_i^a \right) \|z_{n+(N-1)(n+1)}\|^2 \right) \quad (2.8)$$

so that its Hamiltonian flow given by

$$\varphi_t^{G_a}(z_0, \dots, z_{n+(N-1)(n+1)}) = \left(e^{\frac{2\pi\sqrt{-1}\ell_0^a t}{p}} z_0, \dots, e^{\frac{2\pi\sqrt{-1}\ell_n^a t}{p}} z_n, \dots, e^{\frac{2\pi\sqrt{-1}\ell_0^a t}{p}} z_{(n+1)(N-1)}, \dots, \right. \\ \left. e^{\frac{2\pi\sqrt{-1}\ell_{n-1}^a t}{p}} z_{n+(n+1)(N-1)-1}, e^{\frac{2\pi\sqrt{-1}(\ell_n^a - N \sum_i \ell_i^a) t}{p}} z_{n+(N-1)(n+1)} \right) \quad (2.9)$$

generates a loop with Maslov index zero. We have that

$$\sum_{i=0}^n \ell_i^a = j_a \sum_{i=0}^n \ell_i \pmod{p}. \quad (2.10)$$

and so, by Proposition 2.2, $N \sum_{i=0}^n \ell_i^a = 0 \pmod{p}$. Therefore,

$$\begin{aligned} \varphi_1^{G^a}(z_0, \dots, z_{n+(N-1)(n+1)}) &= \left(e^{\frac{2\pi\sqrt{-1}\ell_0^a}{p}} z_0, \dots, e^{\frac{2\pi\sqrt{-1}\ell_n^a}{p}} z_n, \dots, e^{\frac{2\pi\sqrt{-1}\ell_0^a}{p}} z_{(n+1)(N-1)}, \dots, \right. \\ &\quad \left. e^{\frac{2\pi\sqrt{-1}\ell_n^a}{p}} z_{n+(N-1)(n+1)} \right) \\ &= (\psi^{j^a}(z_0, \dots, z_n), \dots, \psi^{j^a}(z_{(n+1)(N-1)}, \dots, z_{n+(N-1)(n+1)})). \end{aligned}$$

Remark 2.3. Note that the key reason why we take N copies of \mathbb{C}^{n+1} is to generate a loop with *zero* Maslov index whose time one map is $(\psi^{j^a}, \dots, \psi^{j^a})$.

Let α be a contact form on $L_p^{2n+1}(\ell_0, \dots, \ell_n)$ and γ a closed Reeb orbit with non-trivial homotopy class a . Multiplying α by a constant if necessary, we can assume that the period of γ is 1. Let β be the lift of α to S^{2n+1} and $\hat{\gamma}$ a segment of Reeb orbit of β that lifts γ . Let $H_\beta : \mathbb{R}^{2n+2} \rightarrow \mathbb{R}$ be a homogeneous of degree two Hamiltonian such that $H_\beta^{-1}(1) = \Sigma_\beta$ and consider $\hat{\gamma}$ as a segment of Hamiltonian orbit of H_β on Σ_β .

Denote by $\Gamma_\beta : [0, 1] \rightarrow \text{Sp}((2n+2)N)$ the symplectic path given by N copies of the linearized Hamiltonian flow of H_β along $\hat{\gamma}$ with respect to the constant symplectic trivialization of $T\mathbb{R}^{(2n+2)N}$, that is,

$$\Gamma_\beta(t) = \oplus_1^N D\varphi_t^{H_\beta}(\hat{\gamma}(0)) \quad (2.11)$$

with $t \in [0, 1]$. The next result shows how to compute the index of γ in terms of Γ_β . Its proof follows [3, Proposition 3.1].

Proposition 2.4. *We have that*

$$\mu(\gamma) = \frac{\mu(\varphi_{-t}^{G^a} \circ \Gamma_\beta)}{N} + 1.$$

Proof. Consider the symplectization $W = L_p^{2n+1}(\ell_0, \dots, \ell_n) \times \mathbb{R}$ of $L_p^{2n+1}(\ell_0, \dots, \ell_n)$ endowed with the symplectic form $\omega = d(e^r \alpha)$, where r is the coordinate in the \mathbb{R} -component. Let $H : W \rightarrow \mathbb{R}$ be the Hamiltonian given by $H(x, r) = e^r$ so that its Hamiltonian vector field is given by $X_H(x, r) = (R_\alpha(x), 0)$, where R_α is the Reeb vector field. Let γ_H be the Hamiltonian closed orbit of H at $H^{-1}(1)$ corresponding to γ .

Choose a section \mathfrak{s} of $\Lambda_{\mathbb{C}}^n \xi^{\otimes N}$ and consider a trivialization Φ^ξ of $\gamma_H^* \oplus_1^N \xi$ as a Hermitian vector bundle such that its highest complex exterior power coincides with \mathfrak{s} ; see Section 2.3. We have that $\oplus_1^N TW = \oplus_1^N (\xi \oplus \xi^\omega)$, where ξ^ω is the symplectic orthogonal to ξ which admits a global (symplectic) frame given by R_α and the vertical vector field ∂_r . Let Φ^{ξ^ω} be the symplectic trivialization of $\gamma_H^* \oplus_1^N \xi^\omega$ that sends the frame $\{R_\alpha, \partial_r\}^{\oplus N}$ to a fixed symplectic basis. Consider the symplectic trivialization of $\gamma_H^* \oplus_1^N TW$ given by $\Phi := \Phi^\xi \oplus \Phi^{\xi^\omega}$ and let $\Gamma_H = \Gamma_H^\xi \oplus \Gamma_H^{\xi^\omega}$ be the symplectic path given by $\Phi_t \circ \oplus_1^N D\varphi_t^H \circ \Phi_0^{-1}$, where $\Gamma_H^\xi = \Phi_t^\xi \circ \oplus_1^N D\varphi_t^H|_\xi \circ (\Phi_0^\xi)^{-1}$ and $\Gamma_H^{\xi^\omega} = \Phi_t^{\xi^\omega} \circ \oplus_1^N D\varphi_t^H|_{\xi^\omega} \circ (\Phi_0^{\xi^\omega})^{-1}$.

We have that

$$\mu(\gamma_H) = \mu(\Gamma_H)/N = (\mu(\Gamma_H^\xi) + \mu(\Gamma_H^{\xi^\omega}))/N.$$

By our choice of H , we clearly have that $\mu(\gamma) = \mu(\Gamma_H^\xi)/N$ and, since R_α and ∂_r are both invariant under the linearized flow of H , $\Gamma_H^{\xi^\omega}$ is the path constant equal the identity and therefore

$$\mu(\Gamma_H^{\xi^\omega}) = -N.$$

Consequently, we arrive at

$$\mu(\gamma_H) = \mu(\gamma) - 1.$$

Hence, in order to prove the proposition, we need to show that

$$\mu(\gamma_H) = \frac{\mu(\varphi_{-t}^{G_a} \circ \Gamma_\beta)}{N}. \quad (2.12)$$

Let $(u_1, \dots, u_{(2n+2)N})$ be a symplectic frame of $\gamma_H^* \oplus_1^N TW$ obtained via the trivialization Φ . Let $\pi : \mathbb{R}^{2n+2} \setminus \{0\} \rightarrow TW$ be the quotient projection with respect to the \mathbb{Z}_p -action in $\mathbb{R}^{2n+2} \setminus \{0\}$. Let $v_i = (d\pi)^{-1}(u_i)$ be the frame of $\hat{\gamma}^* T\mathbb{R}^{(2n+2)N}$ given by the lift of (u_i) . Denote by Ψ the trivialization of $\hat{\gamma}^* T\mathbb{R}^{(2n+2)N}$ induced by (v_i) . Clearly,

$$\mu(\gamma_H) = \mu(\hat{\gamma}; \Psi)$$

where the index on the left hand side is computed using the frame (u_i) .

Now, consider the symplectic frame (w_i) of $\hat{\gamma}^* T\mathbb{R}^{(2n+2)N}$ given by $w_i(t) = D\varphi_t^{G_a}(\hat{\gamma}(0))(v_i(0))$ and let Θ be the trivialization of $\hat{\gamma}^* T\mathbb{R}^{(2n+2)N}$ induced by (w_i) . (Although $\varphi_t^{G_a}$ is a linear flow, we take its derivative because the proof does not use this linearity; see Remark 2.5.) It is easy to see that

$$\mu(\hat{\gamma}; \Theta) = \frac{\mu(D\varphi_t^{G_a}(\hat{\gamma}(0))^{-1} \circ \Gamma_\beta)}{N},$$

where the index on the right hand side is computed using the constant trivialization of $T\mathbb{R}^{(2n+2)N}$. Therefore, in order to prove (2.12), it is enough to show that

$$\mu(\hat{\gamma}; \Psi) = \mu(\hat{\gamma}; \Theta). \quad (2.13)$$

To accomplish this equality, let m be the smallest positive integer such that γ^m is contractible and consider the lift $\tilde{\gamma}$ of γ^m to \mathbb{R}^{2n+2} which is a closed orbit of H_β . Let $g := \psi^{ja}$ and consider the extensions of the frames (v_i) and (w_i) to $\tilde{\gamma}^* T\mathbb{R}^{(2n+2)N}$ given by

$$v'_i(t+j) = \oplus_1^N Dg^j(\tilde{\gamma}(t))(v_i(t)) \text{ and } w'_i(t+j) = \oplus_1^N Dg^j(\tilde{\gamma}(t))(w_i(t))$$

for every $t \in [0, 1]$ and $j \in \{0, \dots, m-1\}$. Since $g^j \circ \pi = \pi$ we have that (v'_i) is the lift of the obvious extension (u'_i) of the frame (u_i) to $(\gamma_H^m)^* \oplus_1^N TW$. It follows from this that

$$\mu(\tilde{\gamma}; \Psi') = \mu(\gamma_H^m), \quad (2.14)$$

where Ψ' is the trivialization of $\tilde{\gamma}^* T\mathbb{R}^{(2n+2)N}$ induced by (v'_i) and the index on the right hand side is computed using the frame (u'_i) . We also conclude from $g^j \circ \pi = \pi$ that

$$w'_i(t) = D\varphi_t^{G_a}(v_i(0)).$$

Since $\varphi_t^{G_a}$ generates a loop with vanishing Maslov index,

$$\mu(\tilde{\gamma}; \Theta') = \mu(\tilde{\gamma}) \quad (2.15)$$

where Θ' is the trivialization of $\tilde{\gamma}^* T\mathbb{R}^{(2n+2)N}$ induced by (w'_i) and the index on the right hand side is computed using the constant trivialization of $T\mathbb{R}^{(2n+2)N}$. It is easy to see that the right hand sides of (2.14) and (2.15) are equal. (Indeed, since γ^m is contractible, the index of its lift $\tilde{\gamma}$ (computed using the constant trivialization of $T\mathbb{R}^{(2n+2)N}$) is equal to the index of γ_H^m using a trivialization defined over a capping disk (c.f. [1, Lemma 3.4]) which coincides with the index of γ_H^m using the frame (u'_i) .) Hence, we arrive at

$$\mu(\tilde{\gamma}; \Psi') = \mu(\tilde{\gamma}; \Theta'). \quad (2.16)$$

Consider the map $A' : [0, m] \rightarrow \text{Sp}((2n+2)N)$ uniquely defined by the property that $A'_t v'_i(t) = w'_i(t)$ for every i and t . Using the fact that $g^j \circ \pi = \pi$ we conclude that

$$v'_i(j) = w'_i(j) \text{ for every } i \implies A'_j = id$$

for every $j \in \{0, \dots, m\}$. We have that

$$\mu(\bar{\gamma}; \Psi') = \mu(\bar{\gamma}; \Theta') + \frac{2\mu_{\text{Maslov}}(A')}{N} \quad (2.17)$$

which implies, by (2.16), that $\mu_{\text{Maslov}}(A') = 0$. Similarly,

$$\mu(\hat{\gamma}; \Psi) = \mu(\hat{\gamma}; \Theta) + \frac{2\mu_{\text{Maslov}}(A)}{N} \quad (2.18)$$

where $A = A'|_{[0,1]}$. We claim that

$$\mu_{\text{Maslov}}(A') = m\mu_{\text{Maslov}}(A). \quad (2.19)$$

which would imply (2.13), finishing the proof. To prove the claim, note that

$$\begin{aligned} \bigoplus_1^N Dg^j(\hat{\gamma}(t))A_t v'_i(t) &= \bigoplus_1^N Dg^j(\hat{\gamma}(t))w'_i(t) \\ &= w'_i(t+j) \\ &= A'_{t+j} v'_i(t+j) \\ &= A'_{t+j} \bigoplus_1^N Dg^j(\hat{\gamma}(t))v'_i(t) \end{aligned}$$

for every i and consequently

$$A'_{t+j} = \bigoplus_1^N Dg^j(\hat{\gamma}(t))A_t \bigoplus_1^N Dg^j(\hat{\gamma}(t))^{-1} \quad (2.20)$$

for every $t \in [0, 1]$ and $j \in \{0, \dots, m-1\}$. Now, let g_s be a homotopy between $g_0 = g$ and $g_1 = Id$ (the existence of this homotopy follows from the fact that ψ^{j_a} is the time one map of a flow). Note that the curve $\bigoplus_1^N Dg^j(\hat{\gamma}(t))A_t \bigoplus_1^N Dg^j(\hat{\gamma}(t))^{-1}$ is homotopic to the curve A_t via the homotopy $\bigoplus_1^N Dg_s^j(\hat{\gamma}(t))A_t \bigoplus_1^N Dg_s^j(\hat{\gamma}(t))^{-1}$. Thus, all the loops $t \in [0, 1] \mapsto A'_{t+j}$, $j \in \{0, \dots, m-1\}$, are homotopic to the loop $t \in [0, 1] \mapsto A_t$. Since A' is a concatenation of these loops, we conclude the equality (2.19). \square

Remark 2.5. The previous proof does not use the linearity of the flow $\varphi_t^{G_a}$: it uses only that it generates a loop with zero Maslov index. Therefore, if $(\psi^{j_a}, \dots, \psi^{j_a})$ is the time one map of a Hamiltonian flow $\varphi_t^{G_a}$ that generates a loop with vanishing Maslov index then

$$\mu(\gamma) = \frac{\mu(D\varphi_t^{G_a}(\hat{\gamma}(0))^{-1} \circ \Gamma_\beta)}{N} + 1.$$

The next proposition shows how to compute k_a easily from the weights that determine the lens space.

Proposition 2.6. *We have that*

$$k_a = w_-^a - w_+^a + \frac{2 \sum_i \ell_i^a}{p} + 1,$$

Proof. Consider the Hamiltonian $H : \mathbb{R}^{2n+2} \rightarrow \mathbb{R}$ given by

$$H(z_0, \dots, z_n) = \frac{j_a \pi}{p} \|z_0\|^2 + \sum_{i=1}^n \pi \epsilon_i \|z_i\|^2$$

where the coefficients $j_a/p, \epsilon_1, \dots, \epsilon_n$ are rationally independent and the ϵ_i 's are positive and very small. Consider the ellipsoid $E = H^{-1}(1)$ and let β be the contact form on S^{2n+1} such that $\Sigma_\beta = E$. Let α be the induced contact form on $L_p^{2n+1}(\ell_0, \dots, \ell_n)$ (note that E is invariant under the \mathbb{Z}_p -action).

Let γ be the closed orbit of H given by $\gamma(t) = (e^{2\pi\sqrt{-1}tj_a/p}, 0, \dots, 0)$. Since $\ell_0^a = j_a \pmod{p}$ (by our normalization of the homotopy weights), given $a \in \pi_1(L_p^{2n+1}(\ell_0, \dots, \ell_n))$, we have that the closed orbit γ_a of α given by the projection of $\gamma|_{[0,1]}$ (we are tacitly identifying E with the unit sphere) has homotopy class a .

Choosing $\epsilon_1, \dots, \epsilon_n$ sufficiently small, it is easy to see that γ_a is the closed orbit with smallest index among the closed orbits of α with homotopy class a . Thus, it is enough to show that

$$\mu(\gamma_a) = w_-^a - w_+^a + \frac{2 \sum_i \ell_i^a}{p} + 1.$$

Let $\Gamma_\beta : [0, 1] \rightarrow \text{Sp}((2n+2)N)$ be the path given by $\Gamma_\beta(t) = \bigoplus_1^N D\varphi_t^H(\gamma(0))$. By Proposition 2.4, we have to prove that

$$\frac{\mu(\varphi_{-t}^{G_a} \circ \Gamma_\beta)}{N} = w_-^a - w_+^a + \frac{2 \sum_i \ell_i^a}{p}. \quad (2.21)$$

But

$$\Gamma_\beta(t) = \bigoplus_1^N (e^{2\pi\sqrt{-1}j_a t/p}, e^{2\pi\sqrt{-1}\epsilon_1 t}, \dots, e^{2\pi\sqrt{-1}\epsilon_n t}).$$

From this and (2.9), we have that

$$\begin{aligned} \varphi_{-t}^{G_a} \circ \Gamma_\beta = & \left(e^{\frac{-2\pi\sqrt{-1}(\ell_0^a - j_a)t}{p}}, e^{\frac{-2\pi\sqrt{-1}\ell_1^a t}{p} + 2\pi\sqrt{-1}\epsilon_1 t}, \dots, e^{\frac{-2\pi\sqrt{-1}\ell_n^a t}{p} + 2\pi\sqrt{-1}\epsilon_n t}, \dots, \right. \\ & e^{\frac{-2\pi\sqrt{-1}(\ell_0^a - j_a)t}{p}}, e^{\frac{-2\pi\sqrt{-1}\ell_1^a t}{p} + 2\pi\sqrt{-1}\epsilon_1 t}, \dots, e^{\frac{-2\pi\sqrt{-1}\ell_{n-1}^a t}{p} + 2\pi\sqrt{-1}\epsilon_{n-1} t}, \\ & \left. e^{\frac{-2\pi\sqrt{-1}(\ell_n^a - N \sum_i \ell_i^a)t}{p} + 2\pi\sqrt{-1}\epsilon_n t} \right). \end{aligned}$$

Since the index of the path $e^{\frac{-2\pi\sqrt{-1}(\ell_0^a - j_a)t}{p}}$, $t \in [0, 1]$, is -1 if $\ell_0^a > 0$ (i.e. $\ell_0^a = j_a$) and 1 if $\ell_0^a < 0$ (i.e. $\ell_0^a = j_a - p$), $N \sum_i \ell_i^a = 0 \pmod{p}$ (by Proposition 2.2 and (2.10)) and the ϵ_i 's are very small, we can easily see that

$$\mu(\varphi_{-t}^{G_a} \circ \Gamma_\beta) = N(w_-^a - w_+^a) + \frac{2N \sum_i \ell_i^a}{p}$$

concluding (2.21). \square

2.5. Bott's function. Let $\Gamma : [0, T] \rightarrow \text{Sp}(2n)$ be a symplectic path starting at the identity and $P := \Gamma(T)$ its endpoint. Following [27, 28], one can associate to Γ its Bott's function $\mathcal{B} : S^1 \rightarrow \mathbb{Z}$ which will be a crucial tool throughout this work. It has the following properties:

- (a) (Bott's formula) We have that $\mu(\Gamma^k) = \sum_{z^k=1} \mathcal{B}(z)$ for every $k \in \mathbb{N}$. In particular, the mean index $\hat{\mu}(\Gamma) := \lim_{k \rightarrow \infty} \frac{\mu(\Gamma^k)}{k}$ satisfies

$$\hat{\mu}(\Gamma) = \int_{S^1} \mathcal{B}(z) dz,$$

where the total measure of the circle is normalized to be equal to one.

- (b) If $\Gamma = \Gamma_1 \oplus \Gamma_2$ then $\mathcal{B} = \mathcal{B}_1 + \mathcal{B}_2$ where \mathcal{B}_i is the Bott's function associated to Γ_i for $i = 1, 2$.
- (c) If Γ_1 and Γ_2 are homotopic with fixed endpoints then $\mathcal{B}_1 = \mathcal{B}_2$.
- (d) The discontinuity points of \mathcal{B} are contained in $\sigma(P) \cap S^1$, where $\sigma(P)$ is the spectrum of P .
- (e) $\mathcal{B}(z) = \mathcal{B}(\bar{z})$ for every $z \in S^1$.
- (f) The *splitting numbers* $S_z^\pm(P) := \lim_{\epsilon \rightarrow 0^+} \mathcal{B}(e^{\pm\sqrt{-1}\epsilon}z) - \mathcal{B}(z)$ depend only on P and satisfy, for every $z \in S^1$,

$$S_z^\pm(P) = S_{\bar{z}}^\mp(P), \quad (2.22)$$

$$S_z^\pm(P_1 \oplus P_2) = S_z^\pm(P_1) + S_z^\pm(P_2), \quad (2.23)$$

$$0 \leq S_z^\pm(P) \leq \nu_z(P) \quad (2.24)$$

and

$$S_z^+(P) + S_z^-(P) \leq \eta_z(P) \quad (2.25)$$

where $\nu_z(P)$ and $\eta_z(P)$ are the geometric and algebraic multiplicities of z (viewing P as a complex matrix) respectively if $z \in \sigma(P) \cap S^1$ and zero otherwise. Moreover,

$$\mathcal{B}(e^{\sqrt{-1}\theta}) = \mathcal{B}(1) + S_1^+(P) + \sum_{\phi \in (0, \theta)} (S_{e^{\sqrt{-1}\phi}}^+(P) - S_{e^{\sqrt{-1}\phi}}^-(P)) - S_{e^{\sqrt{-1}\theta}}^-(P) \quad (2.26)$$

for every $\theta \in [0, 2\pi)$. (Note that the sum above makes sense since $S_z^\pm(P) \neq 0$ only for finitely many points $z \in S^1$.)

- (g) $\mathcal{B}_\Gamma(z)$ is lower semicontinuous with respect to Γ in the C^0 -topology. More precisely, let $\mathcal{P}([0, T], \text{Sp}(2n))$ be the set of continuous paths in $\text{Sp}(2n)$ starting at the identity endowed with the C^0 -topology. Then, for a fixed $z \in S^1$, the map

$$\mathcal{P}([0, T], \text{Sp}(2n)) \rightarrow \mathbb{Z}$$

that sends Γ to $\mathcal{B}_\Gamma(z)$, where \mathcal{B}_Γ denotes the Bott's function associated to Γ , is lower semicontinuous, that is,

$$\mathcal{B}_\Gamma(z) = \sup_U \inf_{\Gamma' \in U} \mathcal{B}_{\Gamma'}(z),$$

where the supremum runs over all C^0 -neighborhoods U of Γ in $\mathcal{P}([0, T], \text{Sp}(2n))$.

We refer to [28] for a proof of these properties. In the proof of Theorem 1.4 the following comparison result, proved in [19] using the spectral flow, will play a crucial role.

Theorem 2.7. ([19, Theorem 2.2]) *Let $\Gamma_i : [0, T] \rightarrow \text{Sp}(2n)$ ($i = 1, 2$) be two symplectic paths starting at the identity and satisfying the differential equation*

$$\frac{d}{dt}\Gamma_i(t) = JA_i(t)\Gamma_i(t),$$

where $A_i(t)$ is a path of symmetric matrices. Suppose that $A_1(t) \geq A_2(t)$ for every t and let \mathcal{B}_i be the Bott's function associated to Γ_i . Then

$$\mathcal{B}_1(z) \geq \mathcal{B}_2(z)$$

for every $z \in S^1$.

3. PROOF OF THEOREM 1.4

3.1. Idea of the proof. Let β be the lift of α to S^{2n+1} and H_β be the convex (resp. strictly convex) homogeneous of degree two Hamiltonian such that $H_\beta^{-1}(1) = \Sigma_\beta$. Let $\hat{\gamma}$ be a segment of Hamiltonian orbit of H_β that projects onto γ . Let $\Gamma_\beta : [0, 1] \rightarrow \text{Sp}((2n+2)N)$ be the linearized Hamiltonian flow along $\hat{\gamma}$ (or, more precisely, N copies of it as defined in (2.11)). It is not true, in general, that the index of γ equals the index of Γ_β : as showed in Proposition 2.4, we need to “correct” the path Γ_β taking $\Gamma := \varphi_{-t}^{G_a} \circ \Gamma_\beta$ so that the “correction term” $\varphi_{-t}^{G_a}$ plays a key role.

The convexity (resp. strict convexity) of α implies, by our comparison result given by Theorem 2.7, that

$$\mathcal{B}_\Gamma(z) \geq \mathcal{B}_{G_a}(z) \quad (\text{resp. } \mathcal{B}_\Gamma(z) \geq \mathcal{B}_{G_a^\epsilon}(z))$$

where G_a^ϵ is a small “negative” perturbation of G_a given by (3.1) and \mathcal{B}_Γ , \mathcal{B}_{G_a} and $\mathcal{B}_{G_a^\epsilon}$ are the Bott’s functions associated to Γ , $\varphi_{-t}^{G_a}$ and $\varphi_{-t}^{G_a^\epsilon}$ respectively.

Then, a careful analysis of \mathcal{B}_{G_a} and $\mathcal{B}_{G_a^\epsilon}$ allows us to show that the Bott’s function \mathcal{B}_Γ associated to the linearized Reeb flow along γ satisfies

- (A) $\mathcal{B}_\Gamma(1) \geq k_a$.
- (B) There exists $z \in S^1 \setminus \{1\}$ such that $\mathcal{B}_\Gamma(z) \geq h_a$ (resp. $\mathcal{B}_\Gamma(z) \geq \tilde{h}_a$).
- (C) Under the assumptions of Assertion 3, there exists $z \in S^1 \setminus \{1\}$ such that $\mathcal{B}_\Gamma(z) \geq k_a + n$ (indeed, under these assumptions $h_a = k_a + n$ (resp. $\tilde{h}_a = k_a + n$)).

Assertion A implies, by Bott’s formula, that $\mu(\gamma) \geq k_a$, proving Assertion 1. If γ is hyperbolic then \mathcal{B}_Γ must be constant (because all the splitting numbers vanish). Hence, by Assertion B, if $\mathcal{B}_\Gamma(1) = \mu(\gamma) < h_a$ (resp. $\mu(\gamma) < \tilde{h}_a$) then γ is non-hyperbolic, proving Assertion 2. Finally, we can show that if $\mathcal{B}_\Gamma(1) = \mu(\gamma) = k_a$ and $\mathcal{B}_\Gamma(z) \geq k_a + n$ for some $z \in S^1$ (so that there is a big enough jump of Bott’s function) then γ must be elliptic (Proposition 3.2). Thus, Assertion 3 follows from Assertion C.

3.2. Proof of the theorem. Assume, without loss of generality, that γ has period 1. As in the previous section, let β be the lift of α to S^{2n+1} and H_β be the convex (resp. strictly convex) homogeneous of degree two Hamiltonian such that $H_\beta^{-1}(1) = \Sigma_\beta$. Let $\hat{\gamma}$ be a segment of Hamiltonian orbit of H_β that projects onto γ (here, as before, we are tacitly identifying S^{2n+1} and Σ_β). Let $\Gamma_\beta : [0, 1] \rightarrow \text{Sp}((2n+2)N)$ be the path given by (2.11) and $\Gamma = \varphi_{-t}^{G_a} \circ \Gamma_\beta$. The symplectic path Γ satisfies the differential equation

$$\frac{d}{dt}\Gamma(t) = JA(t)\Gamma(t)$$

where J is the canonical complex structure in $\mathbb{R}^{(2n+2)N}$ and $A(t)$ is a path of symmetric matrices. It follows from [19, Lemma 7.5] that

$$A(t) = -\text{Hess } G_a + (\varphi_t^{G_a})^* \left(\oplus_1^N \text{Hess } H_\beta(\hat{\gamma}(t)) \right) \varphi_t^{G_a}$$

where $(\varphi_t^{G_a})^*$ denotes the transpose of $\varphi_t^{G_a}$ and $\oplus_1^N \text{Hess } H_\beta(\hat{\gamma}(t))$ is the quadratic form on $\mathbb{R}^{(2n+2)N}$ given by N copies of $\text{Hess } H_\beta(\hat{\gamma}(t))$.

Remark 3.1. If $\varphi_t^{G_a}$ is not linear and $\Gamma := D\varphi_t^{G_a}(\hat{\gamma}(0))^{-1} \circ \Gamma_\beta$ (see Remark 2.5) then $A(t) = -\text{Hess } G_a + D\varphi_t^{G_a}(\hat{\gamma}(0))^* \left(\oplus_1^N \text{Hess } H_\beta(\hat{\gamma}(t)) \right) D\varphi_t^{G_a}(\hat{\gamma}(0))$.

Given $\epsilon > 0$, define the Hamiltonian

$$G_a^\epsilon(z_0, \dots, z_{n+(N-1)(n+1)}) = \frac{\pi}{p} \left(\sum_{j=0}^{N-1} \sum_{i=0}^n (\ell_i^a - p\epsilon) \|z_{i+j(n+1)}\|^2 - \left(N \sum_{i=0}^n \ell_i^a \right) \|z_{n+(N-1)(n+1)}\|^2 \right) \quad (3.1)$$

whose flow is clearly linear. Since H_β is convex (resp. strictly convex),

$$\langle (\varphi_t^{G_a})^* (\oplus_1^N \text{Hess } H_\beta(\hat{\gamma}(t))) \varphi_t^{G_a} v, v \rangle = \langle (\oplus_1^N \text{Hess } H_\beta(\hat{\gamma}(t))) \varphi_t^{G_a} v, \varphi_t^{G_a} v \rangle \geq 0 \quad (\text{resp. } > 0)$$

for every t and $v \in \mathbb{R}^{(2n+2)N}$. Hence, $A(t) \geq -\text{Hess } G_a$ (resp. $A(t) \geq -\text{Hess } G_a^\epsilon = 2\epsilon \text{Id} - \text{Hess } G_a$ for some $\epsilon > 0$ sufficiently small). In what follows, we will take ϵ sufficiently small such that $\varphi_{-1}^{G_a^\epsilon}$ has no eigenvalue -1 .

Thus, Theorem 2.7 yields the inequality

$$\mathcal{B}_\Gamma(z) \geq \mathcal{B}_{G_a}(z) \quad (\text{resp. } \mathcal{B}_\Gamma(z) \geq \mathcal{B}_{G_a^\epsilon}(z)) \quad (3.2)$$

for every $z \in S^1$, where \mathcal{B}_Γ and \mathcal{B}_{G_a} (resp. $\mathcal{B}_{G_a^\epsilon}$) are the Bott's functions associated to Γ and $\varphi_{-t}^{G_a}$ (resp. $\varphi_{-t}^{G_a^\epsilon}$) respectively. Using the arguments in the proofs of Proposition 2.4 and [19, Proposition 4.1] we easily conclude that

$$\mathcal{B}_\Gamma(z) = \begin{cases} \mathcal{B}_\Gamma(1)/N + 1 & \text{if } z = 1 \\ \mathcal{B}_\Gamma(z)/N & \text{otherwise,} \end{cases} \quad (3.3)$$

for every $z \in S^1$.

Let us study \mathcal{B}_{G_a} . First, we easily derive from (2.9) that

$$\mathcal{B}_{G_a}(1) = N(w_-^a - w_+^a) + \frac{2N \sum_i \ell_i^a}{p} = N(k_a - 1), \quad (3.4)$$

where the last equality follows from Proposition 2.6. Thus,

$$\mu(\gamma) = \mathcal{B}_\Gamma(1)/N + 1 \geq \mathcal{B}_{G_a}(1)/N + 1 = k_a,$$

proving the first assertion of the theorem.

The proof of the remaining assertions relies on an analysis of $\mathcal{B}_{G_a}(z)$ and $\mathcal{B}_{G_a^\epsilon}(z)$ for $z \neq 1$. Once we know the Bott's function at 1, it is completely determined by its splitting numbers; see (2.26). The eigenvalues of $\varphi_{-1}^{G_a}$ are the following:

- Those with negative imaginary part, given by $e^{-2\pi\sqrt{-1}\ell_i^a/p}$ with $\ell_i^a > 0$ and $\ell_i^a \neq p/2$, having multiplicity μ_i^a , and their complex conjugates.
- Those with non-negative imaginary part, given by $e^{-2\pi\sqrt{-1}\ell_i^a/p}$ with $\ell_i^a < 0$ or $\ell_i^a = p/2$, having multiplicity ν_i^a , and their complex conjugates.

Since the splitting numbers satisfy (2.22), it is enough to determine them in the eigenvalues with non-negative imaginary part given by $e^{2\pi\sqrt{-1}\bar{\ell}_i^a/p}$ for $i \in \{1, \dots, k\}$. (Recall that $\bar{\ell}_1^a < \dots < \bar{\ell}_k^a$ are the absolute values of the homotopy weights.) From the multiplicity of the eigenvalues discussed above, the additivity of the splitting numbers (2.23) and [28, List 9.1.12, page 198], we conclude the following:

- $S^+(e^{2\pi\sqrt{-1}\bar{\ell}_i^a/p}) = \mu_i^a$ and $S^-(e^{2\pi\sqrt{-1}\bar{\ell}_i^a/p}) = \nu_i^a$ if $\ell_i^a \neq p/2$;
- $S^+(e^{2\pi\sqrt{-1}\bar{\ell}_i^a/p}) = S^-(e^{2\pi\sqrt{-1}\bar{\ell}_i^a/p}) = \nu_i^a$ if $\ell_i^a = p/2$ (in this case $i = k$).

From this and (3.4) we can conclude that \mathcal{B}_{G_a} is given by the following. Since $\mathcal{B}_{G_a}(z) = \mathcal{B}_{G_a}(\bar{z})$ for every $z \in S^1$, it is enough to establish the values of \mathcal{B}_{G_a} in the upper (closed) half-circle. We have that if $\bar{\ell}_k^a \neq p/2$ (resp. $\bar{\ell}_k^a = p/2$) then

$$\frac{\mathcal{B}_{G_a}(e^{\sqrt{-1}\theta})}{N} = \begin{cases} k_a - 1 & \text{if } \theta \in [0, 2\pi\bar{\ell}_1^a/p) \\ k_a - 1 + \sum_{i=1}^{j-1} \mu_i^a - \sum_{i=1}^j \nu_i^a & \text{if } \theta = 2\pi\bar{\ell}_j^a/p, j = 1, \dots, k \text{ (resp. } j = 1, \dots, k-1) \\ k_a - 1 + \sum_{i=1}^j \mu_i^a - \sum_{i=1}^j \nu_i^a & \text{if } \theta \in (2\pi\bar{\ell}_j^a/p, 2\pi\bar{\ell}_{j+1}^a/p), j = 1, \dots, k-1 \\ k_a - 1 + \sum_{i=1}^k \mu_i^a - \sum_{i=1}^k \nu_i^a & \text{if } \theta \in (2\pi\bar{\ell}_k^a/p, \pi] \text{ (resp. } \theta = 2\pi\bar{\ell}_k^a/p = \pi), \end{cases} \quad (3.5)$$

for every $\theta \in [0, \pi]$; see Figures 1 and 2. Note that the function is lower semicontinuous since the splitting numbers are non-negative.




FIGURE 1. The function \mathcal{B}_{G_a}/N with the jumps μ_i^a and ν_i^a when $\bar{\ell}_k^a \neq p/2$. The first jump goes down and the second goes up (when we move counterclockwise in the circle or, equivalently, move to the right in the figure).




FIGURE 2. The function \mathcal{B}_{G_a}/N with the jumps μ_i^a and ν_i^a when $\bar{\ell}_k^a = p/2$. The jump at $\pi = 2\pi\bar{\ell}_k^a/p$ goes down.

From this, we conclude that

$$\max_{z \in S^1} \mathcal{B}_{G_a} = \max_{z \in S^1 \setminus \{1\}} \mathcal{B}_{G_a} = Nh_a.$$

Using this, (3.2) and (3.3) we infer that there exists $z \in S^1 \setminus \{-1\}$ such that $\mathcal{B}_\gamma(z) \geq h_a$. Hence, if $\mu(\gamma) < h_a$ then $\mathcal{B}_\gamma(1) < \mathcal{B}_\gamma(z)$ which implies that \mathcal{B}_γ is not constant and therefore γ is not hyperbolic, proving the second assertion of the theorem when α is not necessarily strictly convex.

When α is strictly convex, we have to study \mathcal{B}_{G_ϵ} . Taking $\epsilon > 0$ sufficiently small, we conclude that the eigenvalues of $\varphi_{-1}^{G_\epsilon}$ are the following:

- Those with negative imaginary part, given by $e^{-2\pi\sqrt{-1}(\ell_i^a/p-\epsilon)}$ with $\ell_i^a > 0$, having multiplicity $\tilde{\mu}_i^a$, and their complex conjugates.
- Those with positive imaginary part, given by $e^{-2\pi\sqrt{-1}(\ell_i^a/p-\epsilon)}$ with $\ell_i^a < 0$, having multiplicity $\tilde{\nu}_i^a$, and their complex conjugates.

The possibly non-zero splitting numbers of \mathcal{B}_{G_ϵ} in the upper-half circle are the following (see [28, List 9.1.12, page 198]):

- $S^+(e^{2\pi\sqrt{-1}(\bar{\ell}_i^a/p-\epsilon)}) = \tilde{\mu}_i^a$ and $S^-(e^{2\pi\sqrt{-1}(\bar{\ell}_i^a/p-\epsilon)}) = 0$;
- $S^+(e^{2\pi\sqrt{-1}(\bar{\ell}_i^a/p+\epsilon)}) = 0$ and $S^-(e^{2\pi\sqrt{-1}(\bar{\ell}_i^a/p+\epsilon)}) = \tilde{\nu}_i^a$ if $\bar{\ell}_i^a \neq p/2$.

(Recall here that, by our choice of ϵ , $\varphi_{-1}^{G_\epsilon}$ has no eigenvalue -1 .) Now, note that an argument analogous to the one to derive (3.4) shows that $\mathcal{B}_{G_\epsilon} = N(k_a - 1)$. Using this and our knowledge about the splitting numbers we can deduce that if $\bar{\ell}_k^a \neq p/2$ (resp. $\bar{\ell}_k^a = p/2$) then

$$\frac{\mathcal{B}_{G_\epsilon}(e^{\sqrt{-1}\theta})}{N} = \begin{cases} k_a - 1 & \text{if } \theta \in [0, 2\pi(\bar{\ell}_1^a/p - \epsilon)] \\ k_a - 1 + \sum_{i=1}^j \tilde{\mu}_i^a - \sum_{i=0}^{j-1} \tilde{\nu}_i^a & \text{if } \theta \in (2\pi(\bar{\ell}_j^a/p - \epsilon), 2\pi(\bar{\ell}_j^a/p + \epsilon)] \\ k_a - 1 + \sum_{i=1}^j \tilde{\mu}_i^a - \sum_{i=1}^j \tilde{\nu}_i^a & \text{if } \theta \in [2\pi(\bar{\ell}_j^a/p + \epsilon), 2\pi(\bar{\ell}_{j+1}^a/p - \epsilon)] \\ k_a - 1 + \sum_{i=1}^k \tilde{\mu}_i^a - \sum_{i=0}^{k-1} \tilde{\nu}_i^a & \text{if } \theta \in (2\pi(\bar{\ell}_k^a/p - \epsilon), 2\pi(\bar{\ell}_k^a/p + \epsilon)] \text{ (resp. } \theta \in (\pi(1 - 2\epsilon), \pi]) \\ k_a - 1 + \sum_{i=1}^k \tilde{\mu}_i^a - \sum_{i=1}^k \tilde{\nu}_i^a & \text{if } \theta \in [2\pi(\bar{\ell}_k^a/p + \epsilon), \pi] \text{ and } \bar{\ell}_k^a \neq p/2 \end{cases} \quad (3.6)$$

for every $\theta \in [0, \pi]$ and $j \in \{1, \dots, k-1\}$; see Figures 3 and 4. Note that when $\bar{\ell}_k^a = p/2$, \mathcal{B}_{G_ϵ} has a jump that goes up at $\theta = \pi(1 - 2\epsilon)$ while \mathcal{B}_{G_a} has a jump that goes down at $\theta = \pi$.




FIGURE 3. The function $\mathcal{B}_{G_\epsilon}/N$ with the jumps $\tilde{\mu}_i^a$ and $\tilde{\nu}_i^a$ when $\bar{\ell}_k^a \neq p/2$. The first jump goes up and happens when $\theta = 2\pi(\frac{\bar{\ell}_1^a}{p} - \epsilon)$. The second one goes down and happens when $\theta = 2\pi(\frac{\bar{\ell}_1^a}{p} + \epsilon)$.




FIGURE 4. The function $\mathcal{B}_{G_a^\epsilon}/N$ with the jumps $\tilde{\mu}_i^a$ and $\tilde{\nu}_i^a$ when $\bar{\ell}_k^a = p/2$. The jump at $\pi(1 - 2\epsilon) = 2\pi(\frac{\ell_k^a}{p} - \epsilon)$ goes up.

From this, we conclude that

$$\max_{z \in S^1} \mathcal{B}_{G_a^\epsilon} = \max_{z \in S^1 \setminus \{-1\}} \mathcal{B}_{G_a^\epsilon} = N\tilde{h}_a.$$

Using this, (3.2) and (3.3) we conclude that there exists $z \in S^1 \setminus \{-1\}$ such that $\mathcal{B}_\gamma(z) \geq \tilde{h}_a$. Thus, if $\mu(\gamma) < \tilde{h}_a$ then \mathcal{B}_γ is not constant and therefore γ is not hyperbolic, proving the second assertion of the theorem when α is strictly convex.

To prove the third assertion, note that from our assumptions that $\ell_i^a > 0$ and $\ell_i^a \neq p/2$ (resp. $\ell_i^a > 0$) for every i we have that $h_a = k_a + n$ (resp. $\tilde{h}_a = k_a + n$) and therefore $\max_{z \in S^1 \setminus \{-1\}} \mathcal{B}_{G_a} = N(k_a + n)$ (resp. $\max_{z \in S^1 \setminus \{-1\}} \mathcal{B}_{G_a^\epsilon} = N(k_a + n)$). Using this, (3.2) and (3.3) we conclude that there exists $z \in S^1 \setminus \{-1\}$ such that $\mathcal{B}_\gamma(z) \geq k_a + n$. Now, the result readily follows from the following proposition. Recall that a symplectic path $\Gamma : [0, T] \rightarrow \text{Sp}(2n)$ is elliptic if every eigenvalue of $\Gamma(T)$ has modulus one.

Proposition 3.2. *Let $\Gamma : [0, T] \rightarrow \text{Sp}(2n)$ be a symplectic path starting at the identity such that $\mathcal{B}_\Gamma(z) - \mathcal{B}_\Gamma(1) \geq n$ for some $z \in S^1$. Then Γ is elliptic.*

Proof. Let θ be the smallest positive number such that $z = e^{\sqrt{-1}\theta}$. Let $P = \Gamma(T)$ and $\{e^{\sqrt{-1}\theta_1}, \dots, e^{\sqrt{-1}\theta_l}\}$ the eigenvalues of P with modulus one and argument $0 < \theta_k < \theta$, such that $\theta_k < \theta_{k+1}$ for every $1 \leq k \leq l - 1$. By (2.26),

$$\mathcal{B}_\Gamma(z) - \mathcal{B}_\Gamma(1) = (S_1^+(P) - S_z^-(P)) + \sum_{k=1}^l (S_{e^{\sqrt{-1}\theta_k}}^+(P) - S_{e^{\sqrt{-1}\theta_k}}^-(P)).$$

Since the splitting numbers are non-negative (see (2.24)), it follows from our assumption that

$$S_1^+(P) + \sum_{k=1}^l S_{e^{\sqrt{-1}\theta_k}}^+(P) \geq n.$$

By (2.22), this implies that

$$S_1^-(P) + \sum_{k=1}^l S_{e^{-\sqrt{-1}\theta_k}}^-(P) \geq n.$$

Putting these two inequalities together we arrive at

$$\begin{aligned} 2n &\geq \sum_{z \in S^1} \eta_z(P) \geq \sum_{z \in S^1} S_z^+(P) + S_z^-(P) \\ &\geq S_1^+(P) + S_1^-(P) + \sum_{k=1}^l (S_{e^{\sqrt{-1}\theta_k}}^+(P) + S_{e^{-\sqrt{-1}\theta_k}}^-(P)) \geq 2n, \end{aligned}$$

where the first inequality is trivial by a dimensional reason, the second inequality holds by (2.25) and the third inequality follows again from the fact that the splitting numbers are non-negative. This ensures that all of the above inequalities are in fact equalities. Hence, Γ is elliptic. \square

4. PROOF OF COROLLARY 1.10

We have only to prove the existence/non-existence of positive homotopy classes. In all the assertions, we have that the weights assume only two values 1 and q . Let a be the homotopy class such that $j_a = 1$. Firstly, note that when $1 \leq q \leq p/2$ we have that a is obviously positive.

Suppose now that $-p/2 < q \leq -1$. The integers mq , with $m \in \mathbb{N}$, form a negative arithmetic sequence with first term bigger than $-p/2$ and with distance less than $p/2$ between consecutive terms. Hence, there exists a smallest positive $k \in \mathbb{N}$ such that

$$-p < kq \leq -p/2.$$

Moreover, since $-p/2 < q \leq -1$, we also have that

$$2 \leq k \leq [p/2]$$

and

$$k > p/2 \text{ iff } p \text{ is odd and } q = -1.$$

Hence, when $q = -1$ and p is odd, the homotopy weights of a^j are j and $-j$ for all $1 \leq j \leq p/2$, and there are no positive homotopy classes in the fundamental group of $L_p^{2n+1}(\ell_0, \dots, \ell_n)$. On the other hand, when $-p/2 < q < -1$ or when $q = -1$ and p is even, the homotopy weights of a^k are $k > 0$, $p + kq > 0$, and we have that the homotopy class a^k is positive. Note that when $q = -1$ and p is even we have that $k = p/2$ and hence $a^{p/2}$ is the unique positive homotopy class.

5. PROOF OF THEOREM 1.14

Let $M = L_p^{2n+1}(1, \dots, 1)$ be the lens space endowed with the induced contact structure ξ and N be the minimal positive integer such that $Nc_1(\xi) = 0$. In the proof we will take advantage of the following useful fact. A contact form α on a lens space $L_p^{2n+1}(\ell_0, \dots, \ell_n)$ is called toric if its lift β to S^{2n+1} satisfies the property that Σ_β is a linear ellipsoid. (Although this is not the standard definition, the standard definition is equivalent to this one [23].)

Proposition 5.1. *Let $a \in \pi_1(L_p^{2n+1}(\ell_0, \dots, \ell_n))$ and m be a positive integer such that j_a and m are coprime. Given $i \in \{0, \dots, n\}$ let $\hat{\ell}_i = \ell_i \pmod{p}$ such that $1 \leq \hat{\ell}_i \leq p-1$. Then there exists a toric contact form on L whose Reeb flow generates a circle action whose generic (simple) orbit has homotopy class a and Conley-Zehnder index $\frac{2(\sum_i \hat{\ell}_i)}{p} j_a + 2m - n$.*

Proof. We will use the toric geometry description of $L_p^{2n+1}(\ell_0 = 1, \ell_1, \dots, \ell_n)$ and its toric Reeb vectors presented in [3, Section 2] (see in particular Subsection 2.3.3).

The moment cone $C \subset \mathbb{R}^{n+1}$ of $L_p^{2n+1}(\ell_0 = 1, \ell_1, \dots, \ell_n)$ can be chosen to have primitive integral exterior normals $\nu_0, \dots, \nu_n \in \mathbb{Z}^{n+1}$ given by

$$\nu_0 = \begin{bmatrix} -\hat{\ell}_1 \\ \vdots \\ -\hat{\ell}_{n-1} \\ p \\ -\sum_{i=1}^n \hat{\ell}_i \end{bmatrix}, \quad \nu_j = \begin{bmatrix} e_j \\ 0 \\ 1 \end{bmatrix} \text{ for } j = 1, \dots, n-1, \quad \nu_n = \begin{bmatrix} 0 \\ \vdots \\ 0 \\ 0 \\ 1 \end{bmatrix},$$

where $e_1, \dots, e_{n-1} \in \mathbb{Z}^{n-1}$ are the canonical coordinate basis vectors of \mathbb{Z}^{n-1} . The fundamental group $\pi_1(L_p^{2n+1}(\ell_0 = 1, \ell_1, \dots, \ell_n))$ is naturally isomorphic to $\mathbb{Z}^{n+1}/\mathcal{N} \cong \mathbb{Z}_p$, where \mathcal{N} is the \mathbb{Z} -span of $\{\nu_0, \dots, \nu_n\}$. More precisely,

$$[(a_0, \dots, a_n)] = a_{n-1} \pmod{p}.$$

Any toric Reeb vector on $L_p^{2n+1}(\ell_0 = 1, \ell_1, \dots, \ell_n)$ can be written as a positive linear combination of the normals $\nu_0, \dots, \nu_n \in \mathbb{Z}^{n+1}$. Consider the following toric Reeb vector:

$$R_a = \left(\sum_{i=0}^n \frac{\hat{\ell}_i j_a}{p} \nu_i \right) + m \nu_n = \begin{bmatrix} 0 \\ \vdots \\ 0 \\ j_a \\ m \end{bmatrix}.$$

Its generic (simple) orbit γ has homotopy class a , corresponding to j_a , and period 1. We will now use the set-up of Section 2.4 to compute its Conley-Zehnder index.

The lift of γ to $\mathbb{R}^{2(n+1)} = \mathbb{C}^{n+1}$ corresponds to the linear symplectic path

$$\Gamma_t(z_0, z_1, \dots, z_n) = \left(e^{2\pi\sqrt{-1}\frac{\hat{\ell}_0 j_a}{p}t} z_0, \dots, e^{2\pi\sqrt{-1}\frac{\hat{\ell}_{n-1} j_a}{p}t} z_{n-1}, e^{2\pi\sqrt{-1}\left(\frac{\hat{\ell}_n j_a}{p} + m\right)t} z_n \right).$$

Hence, we have that

$$\begin{aligned} \mu\left(\varphi_{-t}^{G_a} \circ (\oplus_1^N \Gamma_t)\right) &= 2N \left(\sum_{i=0}^n \frac{\hat{\ell}_i j_a - \ell_i^a}{p} - \frac{n+1}{2} + m + \sum_{i=0}^n \frac{\ell_i^a}{p} \right) \\ &= 2N \left(\left(\sum_{i=0}^n \hat{\ell}_i \right) \frac{j_a}{p} + m - \frac{n+1}{2} \right), \end{aligned}$$

which implies that

$$\mu(\gamma) = \frac{\mu\left(\varphi_{-t}^{G_a} \circ (\oplus_1^N \Gamma_t)\right)}{N} + 1 = 2 \left(\left(\sum_{i=0}^n \hat{\ell}_i \right) \frac{j_a}{p} + m - \frac{n}{2} \right),$$

as desired. \square

5.1. Proof of Assertion 1. Let $a \in \pi_1(M)$ be such that $j_a = p - 1$ so that $k_a = \frac{2n+2}{p}(p - 1) - n$. Let $m = 1$ and $\bar{\alpha}$ be the contact form given by Proposition 5.1 corresponding to a and m . The Conley-Zehnder index of the generic orbit of $\bar{\alpha}$ is given by

$$\frac{2n+2}{p}(p-1) + 2 - n = 2n + 4 - \frac{2n+2}{p} - n.$$

We will need the following elementary lemma.

Lemma 5.2. *Suppose that $p \geq 3$. If $n = 1$ (resp. $n = 2$), assume further that $p \geq 5$ (resp. $p \geq 4$). Then there exists an even integer Δ such that*

$$\frac{2n+4}{p} \leq \Delta - \frac{2n+2}{p} < 2n+2 - \frac{2n+2}{p}. \quad (5.1)$$

Proof. Note that (5.1) is equivalent to

$$\frac{4n+6}{p} \leq \Delta < 2n+2.$$

Hence, we have to show that the interval $[(4n+6)/p, 2n+2)$ contains at least one even integer. It is clear if $n = 1$ and $p \geq 5$ or $n = 2$ and $p \geq 4$. To prove it for $n \geq 3$, note that, since $p \geq 3$, it is enough to show that the interval $[(4n+6)/3, 2n+2)$ contains at least two integer numbers. But it follows from the fact that the Lebesgue measure of $[(4n+6)/3, 2n+2)$ is bigger than or equal to two whenever $n \geq 3$. \square

Now, let $F : \mathbb{R}^2 \rightarrow \mathbb{R}$ be defined as $F(q, p) = \pi(q^2 + p^2)$. Given $\epsilon > 0$ consider the Hamiltonian $G : \mathbb{R}^{2n} \rightarrow \mathbb{R}$ given by

$$G(q_1, p_1, \dots, q_n, p_n) = -\left(\frac{2n+4-\Delta+\epsilon}{2}F(q_1, p_1) + \sum_{i=2}^n \epsilon F(q_i, p_i)\right).$$

Let $f : \mathbb{R} \rightarrow \mathbb{R}$ be a smooth function such that

- $f(0) = 1, f'(0) = 1$;
- $f'(r) \in (0, 1)$ and $f''(r) > 0$ for every $r \in (-r_0, 0)$ and some $r_0 > 0$;
- $f(r) = C$ for every $r \leq -r_0$, where C is a constant bigger than $1/2$.

Define the Hamiltonian

$$H = f \circ G,$$

where the constants ϵ and r_0 will be properly chosen.

Remark 5.3. Note that, choosing $\epsilon < 1$, r_0 very small and C close enough to 1, we can make H arbitrarily uniformly C^1 -close to the constant function equal to one.

Consider the lens space M as the prequantization circle bundle of an orbifold B given by the quotient of M by the circle action generated by the Reeb flow of $\bar{\alpha}$. Let $\pi : M \rightarrow B$ be the quotient projection. Take a point x_0 in the smooth part of B and a neighborhood U of x_0 with Darboux coordinates $(q_1, p_1, \dots, q_n, p_n)$ identifying x_0 with the origin. Taking r_0 sufficiently small and viewing H as an Hamiltonian on U , extend H to B setting $H|_{B \setminus U} \equiv C$. Define the contact form

$$\alpha = \bar{\alpha}/\hat{H},$$

where $\hat{H} = H \circ \pi$. The Reeb vector field of α is given by

$$R_\alpha = \hat{H}R_{\bar{\alpha}} + \hat{X}_H,$$

where $R_{\bar{\alpha}}$ is the Reeb vector field of $\bar{\alpha}$ (assume that the generic orbit of $\bar{\alpha}$ has minimal period one) and \hat{X}_H is the horizontal lift of the Hamiltonian vector field of H . By the construction of H , clearly $W := \pi^{-1}(U)$ is invariant under the Reeb flow of α and outside W the Reeb vector field of α is a constant multiple of $R_{\bar{\alpha}}$.

Lemma 5.4. *The contact form α is dynamically convex.*

Let γ_0 be the simple closed orbit of R_{α} over x_0 . Recall that, by construction, the homotopy class of γ_0 is a .

Lemma 5.5. *We have that $\mu(\gamma_0) < k_a$.*

The proof of first assertion of Theorem 1.14 follows immediately from Lemmas 5.4 and 5.5. These two lemmas are proved in the next section.

5.2. Proof of Lemmas 5.4 and 5.5.

5.2.1. *Proof of Lemma 5.4.* Consider on $W \simeq U \times S^1$ the coordinates (x, θ) , where $x = (q_1, p_1, \dots, q_n, p_n)$ and θ is the coordinate along the fiber such that, with respect to these coordinates, x_0 is the origin and

$$\bar{\alpha}|_W = \lambda + d\theta,$$

where $\lambda = \frac{1}{2} \sum_{i=1}^n (q_i dp_i - p_i dq_i)$ is the Liouville form. In what follows, for a shorter notation, let $TU^N = \bigoplus_1^N TU$ and $\xi^N = \bigoplus_1^N \xi$. Let γ be a contractible periodic orbit of α . Clearly, if γ lies outside W then $\mu(\gamma) \geq n + 2$ because $\bar{\alpha}$ is toric (and therefore dynamically convex). So suppose that the image of γ is contained in W . The Darboux coordinates induce an obvious (constant) trivialization $D : TU^N \rightarrow U \times \mathbb{R}^{2nN}$. From this we get a trivialization of $\xi^N|_W$ given by

$$\Phi(v_1, \dots, v_N) = \pi_2(D(\pi_* v_1, \dots, \pi_* v_N)),$$

where $\pi_2 : U \times \mathbb{R}^{2nN} \rightarrow \mathbb{R}^{2nN}$ is the projection onto the second factor. It is clear that

$$\mu(\gamma, \Phi) = \mu(\gamma_H), \quad (5.2)$$

where $\mu(\gamma, \Phi)$ stands for the index of γ with respect to the trivialization Φ and $\gamma_H = \pi \circ \gamma$ is the corresponding orbit of H with the index computed using the trivialization D . Let \mathfrak{f} be the generator of $\pi_1(W) \simeq \mathbb{Z}$ given by the homotopy class of a simple orbit of $R_{\bar{\alpha}}$ contained in W . Let $q \in \mathbb{Z}$ be such that $[\gamma] = q\mathfrak{f}$, where $[\gamma]$ is the homotopy class of γ in W . It turns out that q is given by the Hamiltonian action of γ_H . Indeed,

$$q = \int_{\gamma} d\theta = \int_0^T \bar{\alpha}(R_{\alpha}(\gamma(t))) - \lambda(X_H(\gamma_H(t))) dt = \int_0^T H(\gamma_H(t)) - \lambda(X_H(\gamma_H(t))) dt, \quad (5.3)$$

where T is the period of γ_H . Therefore, the action $A_H(\gamma_H)$ is an integer number. Note that, since γ is contractible and a is a generator of $\pi_1(M)$, q is a positive multiple of p .

Consider a trivialization Ψ of $\gamma^* \xi^N$ induced by our choice of a section of $(\Lambda^n \xi)^{\otimes N}$. The relation between the trivializations Φ and Ψ are given by the following lemma.

Lemma 5.6. *We have that*

$$\mu(\gamma, \Psi) = \mu(\gamma, \Phi) + q(2n + 4 - \frac{2n + 2}{p}).$$

Proof. Choose a simple orbit $\phi(t)$ of $R_{\bar{\alpha}}$ contained in W and let $Q : [0, 1] \times S^1 \rightarrow W$ be a homotopy between γ and ϕ^q . We can extend the trivialization Ψ to $Q^*\xi^N$ inducing a trivialization of $(\phi^q)^*\xi^N$. We have that

$$\mu(\gamma, \Psi) - \mu(\gamma, \Phi) = \mu(\phi^q, \Psi) - \mu(\phi^q, \Phi).$$

But an easy computation shows that

$$\mu(\phi^q, \Psi) - \mu(\phi^q, \Phi) = \mu(\phi^q, \Psi) + n = q(2n + 4 - \frac{2n + 2}{p}).$$

□

From now on, if the trivialization is not explicitly stated we use a trivialization given by a section of $(\Lambda^n \xi)^{\otimes N}$ and the trivialization D for closed orbits of Hamiltonians on U . Note that the index with respect to D coincides with the index using the constant trivialization of $T\mathbb{R}^{2n}$. Let $U_0 = \{x \in U; -G(x) < r_0\}$. Clearly, U_0 is invariant under the Hamiltonian flow of H and if the image of γ_H is not contained in U_0 then $\mu(\gamma_H) = -n$ which implies, by Lemma 5.6, that $\mu(\gamma) = -n + q(2n + 4) - q(2n + 2)/p \geq n + 2$ since $q \geq p \geq 3$.

Thus suppose that the image of γ_H lies in U_0 . If $\gamma_H(t) = 0$ for some t then $\gamma_H = (\gamma_H^0)^{pk}$ for some $k \in \mathbb{N}$, where $\gamma_H^0 : [0, 1] \rightarrow U_0$ is the constant solution $\gamma_H^0(t) \equiv 0$. A direct computation shows that in this case, due to our choice of f , the linearized Hamiltonian flows of H and G along γ_H coincide and therefore have the same index. But the Hamiltonian flow of G is given by

$$\varphi_t^G(z_1, \dots, z_n) = (e^{-(2n+4-\Delta+\epsilon)\pi\sqrt{-1}t}z_1, e^{-2\epsilon\pi\sqrt{-1}t}z_2, \dots, e^{-2\epsilon\pi\sqrt{-1}t}z_n), \quad (5.4)$$

where we are identifying (q_i, p_i) with $z_i = q_i + \sqrt{-1}p_i$. Thus, it is clear that if $k = 1$ then, choosing $\epsilon < 1/p$, we have

$$\mu(\gamma_H) = -p(2n + 4) + p\Delta - 1 - (n - 1) = -p(2n + 4) + p\Delta - n.$$

This implies, by Lemma 5.6, that $\mu(\gamma) = p\Delta - (2n + 2) - n \geq n + 2$ (note that $q = pk = p$), where we used in the last inequality that $p\Delta - (2n + 2) \geq 2n + 2$ by (5.1). The case where $k > 1$ follows from the fact that if a symplectic path Γ in $\text{Sp}(2n)$ starting at the identity satisfies $\mu(\Gamma) \geq n + 2$ then $\mu(\Gamma^k) \geq n + 2$ for every k .

Now, let us consider the remaining case, where γ_H lies inside U_0 and $\gamma_H(t) \neq 0$ for every t . Let $\gamma_G(t) = \gamma_H(t/f'(G(\gamma_H(0))))$ be the corresponding orbit of G . The next three lemmas are taken from [19]. For the sake of completeness, we will provide their proofs.

Lemma 5.7. *We have that $|\mu(\gamma_H) - \mu(\gamma_G)| \leq 1$.*

Proof. Firstly, notice that it is enough to find some symplectic trivialization $\Phi^t : T_{\gamma_G(t)}\mathbb{R}^{2n} \rightarrow \mathbb{R}^{2n}$ such that

$$|\mu(\gamma_G, \Phi^t) - \mu(\gamma_H, \Phi^{f'(e)t})| \leq 1,$$

where $e = G(\gamma_H(0))$ and the indexes above are the indexes of the symplectic paths defined using the corresponding trivializations. Let T and $T_G = f'(G(\gamma_H(0)))T$ be the periods of γ_H and γ_G respectively. We claim that there exists a symplectic plane P such that $X_G(x) \in P$ and $d\varphi_{T_G}^G(x)P = P$, where $x = \gamma_H(0)$. Indeed, write $x = (z_1, \dots, z_n)$ and let $v_i = (0, \dots, 0, z_i, 0, \dots, 0)$, $w_i = (0, \dots, 0, \sqrt{-1}z_i, 0, \dots, 0)$ and $P_i = \text{span}\{v_i, X_G(x)\}$. Note that if $z_i \neq 0$ then P_i is a symplectic plane because $X_G(x) = -\pi\sqrt{-1}((2n + 4) - \Delta + \epsilon)z_1, 2\epsilon z_2, \dots, 2\epsilon z_n$ and $\omega(v_i, w_i) \neq 0$. Moreover, $d\varphi_{T_G}^G(x)P_i = P_i$ for every i because clearly

$d\varphi_{T_G}^G(x)X_G(x) = X_G(x)$ and $d\varphi_{T_G}^G(x)v_i = v_i$ whenever $v_i \neq 0$ by (5.4). Thus, take $P = P_i$ for some i such that $z_i \neq 0$.

Now, let $S = G^{-1}(e)$. Define

$$D_1(t) = d\varphi_t^G(x)P \quad \text{and} \quad D_2(t) = d\varphi_t^G(x)P^\omega = D_1(t)^\omega,$$

where P^ω is the symplectic orthogonal to P . Since $X_G(\gamma_G(t)) \in D_1(t)$ we have that $D_2(t) \subset T_{\gamma_G(t)}S$ for every t . By construction, D_1 and D_2 are both invariant under the linearized Hamiltonian flow of G . But $\varphi_t^H|_S = \varphi_{f'(e)t}^G|_S$ which implies that D_2 is also invariant under the linearized Hamiltonian flow of H and therefore the same holds for D_1 (since $D_1 = D_2^\omega$). Moreover, $X_H(\gamma_G(t)) = f'(e)X_G(\gamma_G(t)) \in D_1(t)$ for every t .

Take symplectic trivializations $\Phi_1^t : D_1(t) \rightarrow \mathbb{R}^2$ and $\Phi_2^t : D_2(t) \rightarrow \mathbb{R}^{2n-2}$. We can choose Φ_1 such that $\Phi_1^t(X_G(\gamma_G(t))) = v$ for every t , where v is a fixed vector in \mathbb{R}^2 . Let $\Gamma_1^G(t) = \Phi_1^t \circ d\varphi_t^G(x) \circ (\Phi_1^0)^{-1}$, $\Gamma_2^G(t) = \Phi_2^t \circ d\varphi_t^G(x) \circ (\Phi_2^0)^{-1}$ and $\Gamma^G(t) = \Phi^t \circ d\varphi_t^G(x) \circ (\Phi^0)^{-1}$ be the corresponding symplectic paths, where $\Phi = \Phi_1 \oplus \Phi_2$. Similarly, define $\Gamma_1^H(t) = \Phi_1^{f'(e)t} \circ d\varphi_t^H(x) \circ (\Phi_1^0)^{-1}$, $\Gamma_2^H(t) = \Phi_2^{f'(e)t} \circ d\varphi_t^H(x) \circ (\Phi_2^0)^{-1}$ and $\Gamma^H(t) = \Phi^{f'(e)t} \circ d\varphi_t^H(x) \circ (\Phi^0)^{-1}$. Since the Hamiltonian flows of G and H restricted to S differ by a constant reparametrization,

$$\mu(\Gamma_2^G) = \mu(\Gamma_2^H).$$

By our choice of Φ_1 , we have that the spectra of $\Gamma_1^G(t)$ and $\Gamma_1^H(t)$ are equal to $\{1\}$ for every t . This implies that

$$\mu(\Gamma_1^G) \in \{0, -1\} \quad \text{and} \quad \mu(\Gamma_1^H) \in \{0, -1\}$$

and consequently $|\mu(\Gamma_1^G) - \mu(\Gamma_1^H)| \leq 1$. Thus,

$$|\mu(\Gamma^G) - \mu(\Gamma^H)| \leq 1$$

as desired. \square

Now, let $k : \mathbb{R} \rightarrow \mathbb{R}$ be a smooth function such that $k'(r) > 0$ for every $r > 0$. Define $\delta : \mathbb{R} \rightarrow \mathbb{Z}$ as

$$\delta(x) = \begin{cases} 2x + 1 & \text{if } x \in \mathbb{Z}, \\ 2[x] - 1 & \text{otherwise,} \end{cases}$$

where $[x] = \min\{k \in \mathbb{Z}; k \geq x\}$. In what follows, recall that $F : \mathbb{R}^2 \rightarrow \mathbb{R}$ is the Hamiltonian given by $F(q, p) = \pi(q^2 + p^2)$.

Lemma 5.8. *Let $K = -k \circ F$ with k as above. Given a periodic orbit γ_K of K with period T_K then*

$$\mu(\gamma_K) \geq -\delta(k'(F(\gamma_K(0)))T_K).$$

Proof. The flow of K is given by $\varphi_t^K(z) = e^{-2\pi k'(F(z))\sqrt{-1}t}z$, where, as before, we are identifying (q, p) with $z = q + \sqrt{-1}p$. An easy computation shows that if $\gamma_K(t) \equiv 0$ then the linearized Hamiltonian flow on γ_K is given by $e^{-2\pi k'(0)\sqrt{-1}t}z$ and consequently $\mu(\gamma_K) = -\delta(k'(0)T_K)$.

If γ_K is away from the origin, we proceed similarly as in the proof of the previous lemma. Let $\gamma_{-F}(t) = \gamma_K(t/k'(F(\gamma_K(0))))$ be the corresponding periodic orbit of $-F$ with period $T_{-F} = k'(F(\gamma_K(0)))T_K$. Take a symplectic trivialization $\Phi^t : T_{\gamma_{-F}(t)}\mathbb{R}^2 \rightarrow \mathbb{R}^2$ such that $\Phi^t(X_{-F}(\gamma_{-F}(t))) = v$ and $\Phi^t(-\nabla F(\gamma_{-F}(t))/\|\nabla F(\gamma_{-F}(t))\|) = w$ (here the gradient and the norm are taken with respect to the Euclidean metric) where $\{v, w\}$ is a fixed symplectic basis

in \mathbb{R}^2 . Then clearly the linearized Hamiltonian flow of $-F$ with respect to this trivialization is constant equal to the identity and therefore

$$\mu(\gamma_{-F}, \Phi) = -1.$$

Since $X_K(\gamma_K(t)) = k'(F(\gamma_K(0)))X_{-F}(\gamma_K(t))$ is preserved under the linearized Hamiltonian flow of K , we see that the spectrum of the symplectic path

$$\Gamma^K(t) = \Phi^{k'(F(\gamma_K(0)))t} \circ d\varphi_t^K(x) \circ (\Phi^0)^{-1}$$

is constant and equal to $\{1\}$. Thus,

$$\mu(\gamma_K, \Phi) = \mu(\Gamma^K) \in \{-1, 0\}.$$

In particular, we conclude that $\mu(\gamma_K, \Phi) \geq \mu(\gamma_{-F}, \Phi)$. But this implies that

$$\mu(\gamma_K) \geq \mu(\gamma_{-F}),$$

where the indexes above are computed using the canonical trivialization of \mathbb{R}^2 . Finally, an easy computation shows that

$$\mu(\gamma_{-F}) = -\delta(k'(F(\gamma_K(0)))T_K).$$

□

Therefore, it follows from (5.4) and Lemma 5.8 that

$$\mu(\gamma_G) \geq -\delta\left(\frac{2n+4-\Delta+\epsilon}{2}T_G\right) - (n-1)\delta(\epsilon T_G).$$

Let q be the integer number such that $[\gamma] = qf$ given by (5.3). It is clear from the previous inequality and Lemmas 5.6 and 5.7 that

$$\mu(\gamma) \geq q\left(2n+4 - \frac{2n+2}{p}\right) - \delta\left(\frac{2n+4-\Delta+\epsilon}{2}T_G\right) - (n-1)\delta(\epsilon T_G) - 1. \quad (5.5)$$

(Recall that $\mu(\gamma)$ is the index of γ with respect to a trivialization given by a section of $(\Lambda^n \xi)^{\otimes N}$.) Thus, to prove that $\mu(\gamma) \geq n+2$ it is enough to show that

$$q^{-1}(n+2 + \delta\left(\frac{2n+4-\Delta+\epsilon}{2}T_G\right) + (n-1)\delta(\epsilon T_G) + 1) \leq 2n+4 - \frac{2n+2}{p}. \quad (5.6)$$

In order to prove this inequality, we need the following result.

Lemma 5.9. *If γ_H is a non-constant periodic orbit of H then $A_H(\gamma_H) > T_G$, where T_G is the period of the corresponding orbit of G .*

Proof. Let T be the period of γ_H . We have that

$$\begin{aligned} A_H(\gamma_H) &= \int_0^T f(G(\gamma_H(t))) - f'(G(\gamma_H(t)))\lambda(X_G(\gamma_H(t))) dt \\ &= Tf(G(\gamma_H(0))) - f'(G(\gamma_H(0))) \int_0^T \lambda(X_G(\gamma_H(t))) dt. \end{aligned}$$

An easy computation shows that

$$\lambda(X_G(q_1, p_1, \dots, q_n, p_n)) = G(q_1, p_1, \dots, q_n, p_n).$$

Thus,

$$A_H(\gamma_H) = T(f(G) - f'(G)G),$$

where, to simplify the notation, we have omitted the dependence of G on $\gamma_H(0)$.

Let $T_G = Tf'(G)$ be the period of the corresponding orbit of G . Since $f'(G) > 0$, we arrive at

$$\begin{aligned} A_H(\gamma_H) &= T(f(G) - f'(G)G) \\ &= T_G \frac{f(G) - f'(G)G}{f'(G)} \\ &= s(G)T_G, \end{aligned} \tag{5.7}$$

where $s : (-r_0, 0] \rightarrow \mathbb{R}$ is given by

$$s(r) = f(r)/f'(r) - r.$$

We claim that $s(G) > 1$. As a matter of fact, since γ_H is non-constant, $G(\gamma_H(0)) \in (-r_0, 0)$. But, by our choice of f , $s(0) = 1$ and

$$s'(r) = \frac{-f''(r)f(r)}{f'(r)^2} < 0$$

for every $r \in (-r_0, 0)$. Consequently,

$$A_H(\gamma_H) > T_G, \tag{5.8}$$

as desired. \square

Now notice that

$$\delta(cT_G) \leq 2c[T_G] + 1$$

for every positive real number c . Hence,

$$\begin{aligned} & q^{-1}(n+2 + \delta(\frac{2n+4-\Delta+\epsilon}{2}T_G) + (n-1)\delta(\epsilon T_G) + 1) \\ & \leq \frac{2n+3}{q} + \frac{2[T_G]}{q}(\frac{2n+4-\Delta+\epsilon}{2} + (n-1)\epsilon). \end{aligned}$$

In order to estimate the last expression, note that, by Lemma 5.9 and the fact that $q = A_H(\gamma_H)$ is an integer,

$$\frac{[T_G]}{q} \leq 1.$$

Hence,

$$\begin{aligned} \frac{2n+3}{q} + \frac{2[T_G]}{q}(\frac{2n+4-\Delta+\epsilon}{2} + (n-1)\epsilon) & \leq \frac{2n+3}{q} + 2n+4-\Delta+\epsilon + (2n-2)\epsilon \\ & = 2n+4 - (\Delta - \frac{2n+3}{q}) + (2n-1)\epsilon \\ & \leq 2n+4 - \frac{2n+2}{p}, \end{aligned}$$

choosing ϵ such that $(2n-1)\epsilon < 2/p$, where the last inequality follows from the facts that $\Delta - \frac{2n+3}{q} > \Delta - \frac{2n+3}{p}$ because $q \geq p$ and $\Delta - \frac{2n+2}{p} \geq \frac{2n+4}{p}$ by (5.1). This proves (5.6) finishing the proof the lemma.

5.2.2. *Proof of Lemma 5.5.* As in the proof of Lemma 5.4, let $\gamma_H^0 : [0, 1] \rightarrow U_0$ be the constant solution $\gamma_H^0(t) \equiv 0$ of period one so that γ_0 is the corresponding closed Reeb orbit of α . As explained in the previous section, it is clear from (5.4) that

$$\mu(\gamma_H^0) = -(2n + 4) + \Delta - n.$$

Hence, we conclude from Lemma 5.6 that

$$\begin{aligned} \mu(\gamma) &= -(2n + 4) + \Delta - n + 2n + 4 - \frac{2n + 2}{p} \\ &= \Delta - \frac{2n + 2}{p} - n \\ &< n + 2 - \frac{2n + 2}{p}, \end{aligned}$$

where the last inequality follows from (5.1). But note that

$$k_a = \frac{2n + 2}{p}(p - 1) - n = n + 2 - \frac{2n + 2}{p}.$$

5.3. **Proof of Assertion 2.** As in Section 5.1, let $M = L_{11}^5(1, 1, 1)$, with p odd, be the lens space endowed with the induced contact structure ξ and $N = 11$ be the minimal number such that $Nc_1(\xi) = 0$. Let $a \in \pi_1(M)$ be such that $j_a = 5$ so that $k_a = \frac{2n+2}{11}j_a - n = 8/11$. Let $m = n - 1$ and $\bar{\alpha}$ be the contact form given by Proposition 5.1 corresponding to a and m . Although j_a , n , N and p are fixed in what follows, in view of Remark 1.15, we will write them as variables unless explicitly stated.

Remark 5.10. For Remark 1.15, let $M = L_p^{2n+1}(1, \dots, 1)$, with p odd, be the lens space endowed with the induced contact structure ξ and $a \in \pi_1(M)$ be such that $j_a = \lfloor p/2 \rfloor$ so that $h_a = \frac{2n+2}{p}\lfloor p/2 \rfloor$ (see Example 1.2). Suppose that $p \geq 5$, j_a and $n - 1$ are coprime, a is a generator of $\pi_1(M)$ and $n \geq p + 2$. Let $m = n - 1$ and $\bar{\alpha}$ be the contact form given by Proposition 5.1 corresponding to a and m . Under these hypotheses, we have several examples with vanishing first Chern class. For instance, $L_5^{29}(1, \dots, 1)$.

Let $F_h : \mathbb{R}^2 \rightarrow \mathbb{R}$ be the Hamiltonian whose graph and phase portrait are depicted in Figure 5. The Hamiltonian vector field has precisely one hyperbolic singularity p_0 at the origin and two elliptic ones p_{\pm} . The regular orbits are two homoclinic connections and periodic orbits. We ask that

- F_h attains its global minimum at p_{\pm} and $F_h(p_{\pm}) = 0$;
- F_h is strictly convex near the elliptic singularities.

Note that F_h can be chosen arbitrarily C^∞ -small.

As in the previous section, let $F : \mathbb{R}^2 \rightarrow \mathbb{R}$ be defined as $F(q, p) = \pi(q^2 + p^2)$. Consider the Hamiltonians $G : \mathbb{R}^{2n} \rightarrow \mathbb{R}$ and $G_h : \mathbb{R}^{2n} \rightarrow \mathbb{R}$ given by

$$G(q_1, p_1, \dots, q_n, p_n) = f\left(\sum_{i=1}^n F(q_i, p_i)\right) \quad \text{and} \quad G_h(q_1, p_1, \dots, q_n, p_n) = f_h\left(\sum_{i=1}^n F_h(q_i, p_i)\right)$$

where $f : \mathbb{R} \rightarrow \mathbb{R}$ and $f_h : \mathbb{R} \rightarrow \mathbb{R}$ are smooth functions such that

- $f_h(x) = x$ for every $x \leq \delta/2$ for some $\delta > 0$ small;
- $f_h'(x) > 0$ for every $x < \delta$;
- $f_h(x) = C_h$ for every $x \geq \delta$ and some constant C_h ;




FIGURE 5. The graph and phase portrait of the Hamiltonian F_h .

- $f(r) = -r$ for every $r \leq r_0$, where r_0 is such that $G_h^{-1}([0, C_h])$ is contained in the interior of the ball $B_0 := \{(q_1, p_1, \dots, q_n, p_n) \in \mathbb{R}^{2n}; \pi \sum_i q_i^2 + p_i^2 \leq r_0\}$;
- $f(r) = C$ for every $r \geq r_1$ and some constant C , where $r_1 > r_0$;
- $-1 < f'(r) < 0$ for every $r \in (r_0, r_1)$.

We also ask that $G_h^{-1}((0, C_h))$ contains the origin.

Define the time-dependent Hamiltonian

$$H_t(x) = (G \# G_h)_t(x) = G(x) + G_h(\varphi_G^{-t}(x)),$$

where φ_G^t is the Hamiltonian flow of G . The flow of H is given by $\varphi_H^t = \varphi_G^t \circ \varphi_{G_h}^t$. Note that on B_0 , the Hamiltonian flow of G generates a loop of period one. This, together with the fact that G_h is constant outside the ball B_0 , implies that H is 1-periodic in time.

As in Section 5.1, consider the lens space M as the prequantization circle bundle of an orbifold B given by the quotient of the sphere by the circle action generated by the Reeb flow of $\bar{\alpha}$ and let $\pi : M \rightarrow B$ be the quotient projection. Take a point x_0 in the smooth part of B and consider the corresponding periodic orbit $\bar{\gamma}_0$ of $\bar{\alpha}$ over x_0 . Consider a neighborhood $W \simeq U \times S^1$ of $\bar{\gamma}_0$ and coordinates $(q_1, p_1, \dots, q_n, p_n, t)$ on V , where t is the coordinate along the fiber such that, with respect to these coordinates, x_0 is the origin and

$$\bar{\alpha}|_W = \lambda + dt,$$

where $\lambda = \frac{1}{2} \sum_{i=1}^n (q_i dp_i - p_i dq_i)$ is the Liouville form. Take r_1 above small enough such that the ball $B_1 := \{(q_1, p_1, \dots, q_n, p_n) \in \mathbb{R}^{2n}; \pi \sum_i q_i^2 + p_i^2 \leq r_1\}$ is contained in U . Using these coordinates, define the contact form α on M such that

$$\alpha|_W = \lambda + \frac{H_t}{C + C_h} dt,$$

and extending α to the complement of W as $\bar{\alpha}$. By the construction of H_t , one can easily check that α is smooth.

Remark 5.11. Note that G_h can be chosen C^∞ -small. On the other hand, G can be chosen C^1 -small but not C^2 -small. Hence, α can be chosen C^1 -close to the convex contact form $\bar{\alpha}$ but it is not C^2 -close.

The next two lemmas will be proved in Section 5.4.

Lemma 5.12. *The contact form α is dynamically convex.*

Let γ_0 be the simple closed orbit of R_α over x_0 . Recall that, by construction, the homotopy class of γ_0 is a .

Lemma 5.13. *The periodic orbit γ_0 is hyperbolic and satisfies $\mu(\gamma_0) = k_a < h_a$.*

Remark 5.14. Under the context of Remark 5.10, we have that the corresponding periodic orbit γ_0 is hyperbolic and satisfies $\mu(\gamma_0) < h_a$.

5.4. Proof of Lemmas 5.12 and 5.13.

5.4.1. *Proof of Lemma 5.12.* We need first the following lemmata.

Lemma 5.15. *All the periodic orbits of F_h have non-negative index. Moreover, the hyperbolic singularity has index zero.*

Proof. Let γ be a periodic orbit of F_h of period T and let $\Phi : [0, T] \rightarrow \text{Sp}(2)$ be the corresponding symplectic path obtained via the canonical (constant) trivialization of $T\mathbb{R}^2$. Let $z \in \mathbb{C} \setminus \{0\}$ and $\rho(t)$ be a continuous argument of $z(t) = \Phi(t)z$, that is, $\rho : [0, T] \rightarrow \mathbb{R}$ is a continuous function such that $e^{2\pi\sqrt{-1}\rho(t)} = z(t)/|z(t)|$. Let $\Delta(z) := \rho(T) - \rho(0)$ and $I(\Phi) = \{\Delta(z); z \in \mathbb{C} \setminus \{0\}\}$. The set $I(\Phi)$ is an interval of length less than $1/2$ and therefore it is between two integers or contains an integer. Define

$$\tilde{\mu}(\Phi) = \begin{cases} 2k + 1 & \text{if } I(\Phi) \subset (k - 1, k), \\ 2k & \text{if } k \in I(\Phi). \end{cases}$$

It is well known that $\mu(\gamma)$ equals $\tilde{\mu}(\Phi)$ if Φ is non-degenerate and equals $\tilde{\mu}(\Phi)$ or $\tilde{\mu}(\Phi) - 1$ in general. It is easy to see that if γ is hyperbolic then $\mu(\gamma) = \tilde{\mu}(\gamma) = 0$. If γ is one of the elliptic singularities then we have that $\mu(\gamma) \geq 2$ by the convexity of H near these singularities.

Now, suppose that γ is regular. The Hamiltonian vector field X_{F_h} satisfies $\Phi(t)X_{F_h}(\gamma(0)) = X_{F_h}(\gamma(t))$ for every t and therefore one can easily see that $\tilde{\mu}(\Phi) \geq 1$. Hence, $\mu(\gamma) \geq 0$. \square

Lemma 5.16. *All the periodic orbits of G_h have index bounded from below by $-n$.*

Proof. Consider the Hamiltonian $\bar{G}_h(q_1, p_1, \dots, q_n, p_n) = \sum_{i=1}^n F_h(q_i, p_i)$. It follows from Lemma 5.15 that every periodic orbit of \bar{G}_h is non-negative. In particular, every such periodic orbit has non-negative mean index $\hat{\mu}$. From this we can conclude that every periodic orbit of G_h in $G_h^{-1}([0, C_h])$ has index bounded from below by $-n$. Indeed, let γ be a closed orbit of G_h in $G_h^{-1}([0, C_h])$ and denote by $\bar{\gamma}$ the corresponding periodic orbit of \bar{G}_h . It is well known that $\hat{\mu}(\bar{\gamma}) = \hat{\mu}(\gamma)$; see [17, Lemma 2.6]. Since $\hat{\mu}(\bar{\gamma}) \geq 0$ and $|\hat{\mu}(\bar{\gamma}) - \mu(\gamma)| \leq n$ we conclude that $\mu(\gamma) \geq -n$.

Finally, if γ is a closed orbit lying outside $G_h^{-1}([0, C_h])$ then, by construction of f_h , we have that the linearized Hamiltonian flow along γ is constant equal to the identity and therefore $\mu(\gamma) = -n$. \square

Lemma 5.17. *Let γ be a closed orbit of G with integral period T and whose image is contained in B_1 . Then $\mu(\gamma) \geq -n(2T + 1)$.*

Proof. Consider the Hamiltonian $\bar{G}(q_1, p_1, \dots, q_n, p_n) = -\sum_{i=1}^n F(q_i, p_i)$ and let $\bar{\gamma}$ be the corresponding closed orbit of \bar{G} . The period of $\bar{\gamma}$ is $\bar{T} = Tf'(\bar{G}(\gamma))$. It is easy to see

that the index of $\bar{\gamma}$ is bounded from below by $-n(2\bar{T} + 1)$. By our choice of f , $\bar{T} \leq T \implies -n(2\bar{T} + 1) \geq -n(2T + 1)$ with the equality if and only if γ lies in B_0 . On B_0 , clearly $\mu(\bar{\gamma}) = \mu(\gamma)$ and therefore the result holds if γ lies in B_0 . If γ lies in $B_1 \setminus B_0$, we have the strict inequality $-n(2\bar{T} + 1) > -n(2T + 1)$ and one can easily check that $|\mu(\bar{\gamma}) - \mu(\gamma)| \leq 1$. Therefore, $\mu(\gamma) \geq -n(2T + 1)$, where we are using the fact that T is an integer. \square

Proof of Lemma 5.12. Let $W_i = \pi^{-1}(B_i)$ ($i = 0, 1$), where $\pi : W \simeq U \times S^1 \rightarrow U$ is the projection. Since H_t is constant outside B_1 and $\bar{\alpha}$ is toric, we have only to check that every contractible periodic orbit γ of α contained in W_1 satisfies $\mu(\gamma) \geq n + 2$. Note that the projection γ_H of γ to B_1 corresponds to a periodic orbit of H_t with period T given by a positive multiple of p (since a is a generator of $\pi_1(M)$).

As in Section 5.2.1, the Darboux coordinates induce an obvious (constant) trivialization $D : TU^N \rightarrow U \times \mathbb{R}^{2nN}$. From this we get a trivialization of $\xi^N|_W$ given by

$$\Phi(v_1, \dots, v_N) = \pi_2(D(\pi_*v_1, \dots, \pi_*v_N)),$$

where $\pi_2 : U \times \mathbb{R}^{2nN} \rightarrow \mathbb{R}^{2nN}$ is the projection onto the second factor. It is clear that

$$\mu(\gamma, \Phi) = \mu(\gamma_H), \quad (5.9)$$

where $\mu(\gamma, \Phi)$ stands for the index of γ with respect to the trivialization Φ and the index of γ_H is computed using the trivialization D .

Consider a trivialization Ψ of $\gamma^*\xi^N$ induced by our choice of a section of $(\Lambda^n\xi)^{\otimes N}$. The relation between the trivializations Φ and Ψ is given by the following lemma whose proof is analogous to the one of Lemma 5.6.

Lemma 5.18. *We have that*

$$\mu(\gamma, \Psi) = \mu(\gamma, \Phi) + T\left(\frac{2n+2}{p}j_a + 2(n-1)\right).$$

From now on, if the trivialization is not explicitly stated we use a trivialization given by a section of $(\Lambda^n\xi)^{\otimes N}$ and the trivialization D for closed orbits of Hamiltonians on B_1 . Note that the index with respect to D coincides with the index using the constant trivialization of $T\mathbb{R}^{2n}$. Clearly, B_0 and B_1 are both invariant under the Hamiltonian flow of H and if the image of γ_H is not contained in B_1 then $\mu(\gamma_H) = -n$. Write $T = kp$ with k being a positive integer (recall that T is a positive multiple of p). By Lemma 5.18, we have that $\mu(\gamma) = (2n+2)kj_a + 2kp(n-1) - n \geq n + 2$.

Now suppose that the image of γ lies in W_0 . Since on B_0 the Hamiltonian flow of G generates a loop of period one and Conley-Zehnder index $-3n$ we have, by Lemmas 5.16 and 5.18,

$$\begin{aligned} \mu(\gamma) &\geq T((2n+2)j_a/p + 2(n-1) - 2n) - n \\ &= k(2n+2)j_a - 2kp - n \\ &= 2k((n+1)j_a - p) - n \\ &\geq 2(3 \cdot 5 - 11) - 2 = 6 > n + 2 = 4 \end{aligned} \quad (5.10)$$

where in the second inequality and the last equality we used that $j_a = 5$, $k \geq 1$, $n = 2$ and $p = 11$. (Here we are using the fact that the index of a symplectic path Γ in $\text{Sp}(2n)$ composed with a loop with Conley-Zehnder index $-3n$ is equal to the index of Γ minus $2n$.)

Remark 5.19. Regarding Remark 5.10, we have the relation

$$\begin{aligned}
\mu(\gamma) &\geq T\left(\frac{2n+2}{p}j_a + 2(n-1) - 2n\right) - n \\
&= 2k((n+1)j_a - p) - n \\
&\geq 2((n+1)\lfloor p/2 \rfloor - p) - n \\
&= 2((n+1)(p/2 - 1/2) - p) - n \\
&= n(p-2) - p - 1 \geq 2n - p \geq n + 2
\end{aligned} \tag{5.11}$$

where in the second equality we used the fact that p is odd, the third inequality follows from the assumption that $p \geq 5$ (and the fact that $n \geq 1$) and the last one holds because $n \geq p + 2 \implies 2n - p \geq n + p + 2 - p = n + 2$.

Finally, suppose that the image of γ lies in $W_1 \setminus W_0$. Since G_h restricted to $B_1 \setminus B_0$ is constant, γ_H must be a closed orbit of G with integral period. By Lemmas 5.17 and 5.18 we have, as in (5.10),

$$\mu(\gamma) \geq T((2n+2)j_a/p + 2(n-1) - 2n) - n > n + 2$$

using the fact that $j_a = 5$, $k \geq 1$, $n = 2$ and $p = 11$.

Remark 5.20. Concerning Remark 5.10, inequality (5.11) also holds under the assumptions that $p \geq 5$ and $n \geq p + 2$ as explained in Remark 5.19. □

5.4.2. *Proof of Lemma 5.13.* Since the hyperbolic periodic orbit of G_h has index zero and on B_0 the flow of G generates a loop of period one and Conley-Zehnder index $-3n$, we have from Lemma 5.18 that

$$\mu(\gamma) = \frac{2n+2}{p}j_a + 2(n-1) - 2n = \frac{2n+2}{p}j_a - 2 = k_a \tag{5.12}$$

where we are using the fact that $n = 2$. (Here, as before, we are using the fact that the index of a symplectic path Γ in $\text{Sp}(2n)$ composed with a loop with Conley-Zehnder index $-3n$ is equal to the index of Γ minus $2n$.)

Remark 5.21. Under the assumptions of Remark 5.10, note that by equality (5.12) we have that

$$\mu(\gamma) = \frac{2n+2}{p}j_a + 2(n-1) - 2n = \frac{2n+2}{p}j_a - 2 = h_a - 2$$

where we are using the fact that $j_a < p/2$ (because p is odd).

6. PROOF OF THEOREM 1.6

6.1. Proof of Assertion 1. First, note that if α is non-degenerate, then, as mentioned before, $\text{HC}_*^a(L_p^{2n+1}(\ell_0, \dots, \ell_n))$ can be obtained as the homology of a chain complex generated by the good periodic orbits of α with homotopy class a . Therefore, if every closed orbit γ of α with homotopy class a satisfies $\mu(\gamma) > k_a$ we would conclude that $\text{HC}_{k_a}^a(L_p^{2n+1}(\ell_0, \dots, \ell_n)) = 0$, a contradiction. Hence, we must have a periodic orbit γ with homotopy class a such that $\mu(\gamma) \leq k_a$ which implies, by the first assertion of Theorem 1.4, that $\mu(\gamma) = k_a$.

If α is degenerate, we proceed as follows. Suppose that every closed orbit of α with homotopy class a satisfies $\mu(\gamma) > k_a$. Since $\text{HC}_{k_a}^a(L_p^{2n+1}(\ell_0, \dots, \ell_n)) \neq 0$ we have that $\text{HC}_{k_a}^{a,T}(\alpha) \neq 0$

for some $T > 0$ sufficiently large. Let $\bar{\alpha}$ be a non-degenerate perturbation of α such that every periodic orbit of $\bar{\alpha}$ with action less than T is close to some periodic orbit of α . Choosing T away from the action spectrum of α , we have that $\mathrm{HC}_{k_a}^{a,T}(\bar{\alpha}) \cong \mathrm{HC}_{k_a}^{a,T}(\alpha) \neq 0$. Since the index is lower semicontinuous, we have that every periodic orbit $\bar{\gamma}$ of $\bar{\alpha}$ with action less than T satisfies $\mu(\bar{\gamma}) > k_a$ which implies that $\mathrm{HC}_{k_a}^{a,T}(\bar{\alpha}) = 0$, a contradiction. This implies that α has a periodic orbit γ with homotopy class a such that $\mu(\gamma) \leq k_a$ which implies, by the convexity of α as before, that $\mu(\gamma) = k_a$.

6.2. Proof of Assertion 2. The proof of this assertion follows the argument of the proof the second assertion of Theorem 1.14 but it is actually much simpler. We consider the contact form $\bar{\alpha}$ on $M = L_p^{2n+1}(1, \dots, 1)$ whose Reeb flow generates a free circle action ($\bar{\alpha}$ is induced from a constant multiple of the Liouville form restricted to the unit sphere in \mathbb{R}^{2n+2}). In this way, we consider the lens space M as the prequantization circle bundle of the complex projective space $\mathbb{C}P^n$ given by the quotient of the sphere by the circle action generated by the Reeb flow of $\bar{\alpha}$. Let $\pi : M \rightarrow \mathbb{C}P^n$ be the quotient projection.

Take a point x_0 in $\mathbb{C}P^n$ and a neighborhood U of x_0 with Darboux coordinates $(q_1, p_1, \dots, q_n, p_n)$ identifying x_0 with the origin. Consider the Hamiltonian G_h defined in Section 5.3 as an Hamiltonian on U (taking δ sufficiently small) and extend it to $\mathbb{C}P^n$ setting $G_h|_{\mathbb{C}P^n \setminus U} \equiv C_h$. Define the contact form

$$\alpha = \bar{\alpha}/(1 + \hat{G}_h),$$

where $\hat{G}_h = G_h \circ \pi$. The Reeb vector field of α is given by

$$R_\alpha = (1 + \hat{G}_h)R_{\bar{\alpha}} + \hat{X}_{G_h},$$

where $R_{\bar{\alpha}}$ is the Reeb vector field of $\bar{\alpha}$ (assume that the simple orbits of $\bar{\alpha}$ have minimal period one) and \hat{X}_{G_h} is the horizontal lift of the Hamiltonian vector field of G_h . By Remark 5.11, α can be chosen C^∞ -close to $\bar{\alpha}$. Therefore, it is strictly convex.

Let γ_0 be the periodic orbit of α over x_0 . By construction, it is hyperbolic. Let us compute its index. By Lemma 5.15 and the construction of G_h , we have from (5.2) and an inspection of the proof of Lemma 5.6 that

$$\mu(\gamma_0) = \frac{2n + 2}{p}$$

since the right hand side is the Conley-Zehnder index of the (simple) orbits of $\bar{\alpha}$ plus n and the hyperbolic orbit of G_h has index zero. Now, note that the homotopy class a of γ_0 satisfies $j_a = 1$ and therefore $h_a = \tilde{h}_a = (2n + 2)/p$. Consequently, γ_0 is hyperbolic and satisfies $\mu(\gamma_0) = h_a = \tilde{h}_a$, as desired.

6.3. Proof of Assertion 3. Consider the contact form α constructed in the previous section with $n = 1$ and $p = 4$. It is a strictly convex contact form carrying a hyperbolic closed orbit γ_0 with homotopy class a satisfying $\mu(\gamma_0) = h_a = \tilde{h}_a$ (since $j_a = 1 < p/2 = 2$). But we have that $k_a = \tilde{h}_a - n = \tilde{h}_a - 1 \implies \mu(\gamma_0) = k_a + 1$.

7. PROOF OF THEOREM 1.18

Let α be one of the contact forms given by Theorem 1.14 and β its lift to S^{2n+1} . Given a contactomorphism $\bar{\varphi} : S^{2n+1} \leftarrow$ that commutes with ψ , we have to show that there exists a C^1 -neighborhood U of $\bar{\varphi}$ such that $\varphi^*\beta$ cannot be convex for any $\varphi \in U$. Let $\bar{\phi} : L_p^{2n+1}(1, \dots, 1) \leftarrow$ be the contactomorphism induced by $\bar{\varphi}$. As discussed in the introduction, the action $\bar{\phi}_*$ on

$\pi_1(L_p^{2n+1}(1, \dots, 1))$ induced by $\bar{\phi}$ is trivial (because $k_a \neq k_b$ for distinct homotopy classes a and b). We have that $\varphi^*\beta$ is invariant under the conjugated \mathbb{Z}_p -action generated by $\varphi^{-1}\psi\varphi$.

Let $L_p^{2n+1}(1, \dots, 1; \varphi)$ be the quotient of S^{2n+1} by the action generated by $\varphi^{-1}\psi\varphi$. Denote by $\phi : L_p^{2n+1}(1, \dots, 1; \varphi) \rightarrow L_p^{2n+1}(1, \dots, 1)$ the contactomorphism induced by φ . Since $\phi = \bar{\phi}(\bar{\phi}^{-1}\phi)$ and $\bar{\phi}_*$ is the identity, we have that $\phi_* = (\bar{\phi}^{-1}\phi)_*$. Therefore, it is enough to show that if $\phi^*\alpha$ is convex (the definition of convex contact forms on $L_p^{2n+1}(1, \dots, 1; \varphi)$ is analogous to the one for contact forms on $L_p^{2n+1}(1, \dots, 1)$) and γ is a periodic orbit of $\phi^*\alpha$ with homotopy class $(\bar{\phi}^{-1}\phi)_*^{-1}a$ then

- (1) $\mu(\gamma) \geq k_a$;
- (2) if $\mu(\gamma) < h_a$ (resp. $\mu(\gamma) < \tilde{h}_a$) then γ is non-hyperbolic;
- (3) if $\ell_i^a > 0$ and $\ell_i^a \neq p/2$ (resp. $\ell_i^a > 0$) for every i and $\mu(\gamma) = k_a$ then γ is elliptic.

The proof of Theorem 1.4 carries out word by word to prove this except that, instead of G_a , we have to take into account the Hamiltonian $G_a^\varphi := G_a \circ (\bar{\varphi}^{-1}\varphi, \dots, \bar{\varphi}^{-1}\varphi)$ whose flow might not be linear; see Remarks 2.5 and 3.1. Taking U sufficiently small, we have that $\bar{\varphi}^{-1} \circ \varphi$ is C^1 -close to identity and therefore the symplectic path $D\varphi_t^{G_a^\varphi}(\hat{\gamma}(0))^{-1}$ is C^0 -close to the path $\varphi_{-t}^{G_a} = D\varphi_{-t}^{G_a}$.

By the lower semicontinuity of the Bott's function with respect to the symplectic path (in the C^0 -topology) we infer that $\mathcal{B}_{G_a^\varphi}(z) \geq \mathcal{B}_{G_a}(z)$ for every $z \in S^1$, where $\mathcal{B}_{G_a^\varphi}$ is the Bott's function of the symplectic path $D\varphi_t^{G_a^\varphi}(\hat{\gamma}(0))^{-1}$. Since the proof of Theorem 1.4 follows from a lower bound of \mathcal{B}_{G_a} ($\mathcal{B}_{G_a}(1) \geq N(k_a - 1)$ for the first assertion, $\mathcal{B}_{G_a}(z) \geq Nh_a$ and $\mathcal{B}_{G_a}(z) \geq N\tilde{h}_a$ for some $z \in S^1 \setminus \{-1\}$ for the second assertion and $\mathcal{B}_{G_a}(z) \geq N(k_a + n)$ for some $z \in S^1 \setminus \{-1\}$ for the third assertion) we have the same bound for $\mathcal{B}_{G_a^\varphi}$, concluding the desired result.

8. PROOF OF THEOREM 1.20

Let $M = L_p^{2n+1}(1, \dots, 1)$ and let $a \in \pi_1(M)$ be the homotopy class of a simple orbit γ of the obvious contact form α_0 on M whose Reeb flow generates a free circle action (induced from the contact form on S^{2n+1} given by the Liouville form restricted to the unit sphere). Clearly M and α_0 satisfy the hypotheses of Section 2.2 used to conclude (2.4). An easy computation shows that

$$\mu(\gamma^k) = (2n + 2)k/p - n$$

for every $k \in \mathbb{N}$. Therefore, we have from (2.4) that

$$\mathrm{HC}_*^a(M) \cong \bigoplus_{k \in \mathbb{N}} \mathrm{H}_{* - (2n+2)((k-1)p+1)/p+n}(\mathbb{C}P^n; \mathbb{Q})$$

From Example 1.2, we have that $h_a = (2n + 2)/p$. Thus, we have precisely $\lfloor (n + 1)/2 \rfloor$ non-trivial elements in $\mathrm{HC}_*^a(M)$ with degrees less than h_a .

Now, let α be any contact form on M satisfying the hypothesis of the theorem (in particular, α is convex). From the discussion in Section 2.2, we have an injective map

$$\psi : \{0, \dots, \lfloor (n - 1)/2 \rfloor\} \rightarrow \mathcal{P}^a(\alpha)$$

such that if $\gamma_i = \psi(i)$ then $\mathrm{HC}_{(2n+2)/p-n+2i}(\gamma_i) \neq 0$. Since $\mathrm{HC}_*(\gamma_i)$ is supported in $[\mu(\gamma_i), \mu(\gamma_i) + \nu(\gamma_i)]$, we conclude that $\mu(\gamma_i) \leq (2n + 2)/p - n + 2\lfloor (n - 1)/2 \rfloor \leq (2n + 2)/p - 1 < h_a$. Therefore, by Theorem 1.4, we conclude the existence of $\lfloor (n - 1)/2 \rfloor + 1 = \lfloor (n + 1)/2 \rfloor$ non-hyperbolic periodic orbits. Now, we will use our pinching condition to ensure that these closed orbits are simple:

Lemma 8.1. *Under our pinching conditions the periodic orbits $\gamma_i = \psi(i)$ are simple in the sense that they are not an iterate of another periodic orbit with homotopy class a . In particular, the orbits γ_i are geometrically distinct.*

Proof. Let γ be a periodic orbit of α with homotopy class a and period T . The (second or more) iterates of γ with homotopy class a are given by γ^{kp+1} with $k \in \mathbb{N}$. Since $\mu(\gamma_i) < h_a$, it is enough to show that $\mu(\gamma^{kp+1}) \geq n + 2$ because $h_a = (2n + 2)/p < n + 2$ since $p \geq 2$. Let β be the lift of α to S^{2n+1} and $H_\beta : \mathbb{R}^{2n+2} \rightarrow \mathbb{R}$ the homogeneous of degree two Hamiltonian such that $H_\beta^{-1}(1) = \Sigma_\beta$.

Let $\Gamma_\beta : [0, (kp+1)T] \rightarrow \text{Sp}((2n+2)N)$ be the N copies of the linearized Hamiltonian flow of H_β along a lift $\hat{\gamma}$ of γ^{kp+1} as discussed in Section 2.4 and defined in (2.11). By Proposition 2.4 we have that

$$\mu(\gamma^{kp+1}) = \frac{\mu(\varphi_{-t/(kp+1)T}^{G_a} \circ \Gamma_\beta)}{N} + 1. \quad (8.1)$$

We claim that the symplectic path $\Gamma(t) = (\varphi_{-t/(kp+1)T}^{G_a} \circ \Gamma_\beta)(t)$ is positive. As explained in Section 3, Γ satisfies the differential equation

$$\frac{d}{dt}\Gamma(t) = JA(t)\Gamma(t)$$

with

$$A(t) = -\frac{\text{Hess } G_a}{(kp+1)T} + (\varphi_{t/(kp+1)T}^{G_a})^* \left(\bigoplus_1^N \text{Hess } H_\beta(\hat{\gamma}(t)) \right) \varphi_{t/(kp+1)T}^{G_a}.$$

From our pinching condition, $\text{Hess } H_\beta \geq \frac{Id_{2n}}{R^2}$ and therefore

$$A(t) \geq -\frac{\text{Hess } G_a}{(kp+1)T} + \frac{Id_{2nN}}{R^2},$$

where in the last equation we used that $\varphi_t^{G_a}$ is unitary. The right hand side of the last inequality is positive if and only if

$$\frac{\|w\|^2}{R^2} > \frac{\langle \text{Hess } G_a w, w \rangle}{(kp+1)T} \quad (8.2)$$

for every $w \in \mathbb{R}^{(2n+2)N}$. From (2.8) and the fact that $\ell_i^a = 1$ for every i we conclude that the eigenvalues of $\text{Hess } G_a$ are $2\pi/p$ (with multiplicity $2N(n+1) - 2$) and $-2\pi N(n+1)$ (with multiplicity 2). Hence,

$$\frac{\langle \text{Hess } G_a w, w \rangle}{(kp+1)T} \leq \frac{2\pi\|w\|^2}{(kp+1)Tp}. \quad (8.3)$$

Now, we claim that

$$T \geq 2\pi r^2/p. \quad (8.4)$$

Indeed, let $H_r : \mathbb{R}^{2n+2} \rightarrow \mathbb{R}$ be the Hamiltonian given by

$$H_r(x) = \frac{1}{2r^2}\|x\|^2.$$

The inequality $\text{Hess } H_\beta(x)(v, v) \leq r^{-2}\|v\|^2$ applied to $v = x$ implies that

$$H_\beta(x) \leq H_r(x).$$

(Indeed, note that, by homogeneity, $H_\beta(x) = \frac{1}{2}\langle \text{Hess } H(x)x, x \rangle$.) A theorem due to Croke and Weinstein [12, Theorem A] establishes that if $H : \mathbb{R}^{2n+2} \rightarrow \mathbb{R}$ is a convex Hamiltonian homogeneous of degree two such that $H(x) \leq H_r(x)$ then every non-constant periodic solution

of H has period at least $2\pi r^2$. Thus, since the lift of γ^p (with period pT) is a periodic orbit of H_β (note that γ^p is contractible), we conclude that $pT \geq 2\pi r^2$, as desired.

Consequently, from (8.3) and (8.4) we arrive at

$$\frac{\langle \text{Hess } G_a w, w \rangle}{(kp+1)T} \leq \frac{\|w\|^2}{r^2(kp+1)}. \quad (8.5)$$

On the other hand,

$$\frac{\|w\|^2}{R^2} > \frac{\|w\|^2}{r^2(kp+1)} \quad (8.6)$$

because $\frac{r^2}{R^2} > \frac{1}{kp+1} \iff \frac{R^2}{r^2} < kp+1$ which is a consequence of the inequality $\frac{R}{r} < \sqrt{p+1}$ since $k \geq 1$. From (8.5) and (8.6) we conclude (8.2), showing that Γ is positive.

From the positivity of Γ , we have that $\mu(\Gamma) \geq (n+1)N$. Therefore, from (8.1),

$$\mu(\gamma^{kp+1}) = \mu(\Gamma)/N + 1 \geq n+2$$

as desired. \square

9. PROOF OF THEOREM 1.21

Let α_1 be the contact form on $L_p^{2n+1}(1, \dots, 1)$ induced by the Liouville form λ restricted to the unit sphere $S^{2n+1} \subset \mathbb{R}^{2n+2}$. Let $\alpha_t = t^2 \alpha_1$ for $t \in [r, R]$. Note that our pinching condition means that $\alpha_r < \alpha < \alpha_R$. Indeed, write $\alpha = f\lambda|_{S^{2n+1}}$ for a positive function $f : S^{2n+1} \rightarrow \mathbb{R}$, and note that our pinching condition means that $r \leq \|x\| \leq R$ for every $x \in \Sigma_\beta$. But $x \in \Sigma_\beta$ if and only if $x = \sqrt{f(x/\|x\|)}x/\|x\|$ and therefore our pinching condition is equivalent to the inequality $r^2 \leq f(v) \leq R^2$ for every $v \in S^{2n+1}$.

It is easy to see that the Reeb flow of α_t generates a free circle action with period $\pi t^2/p$. Let a be the homotopy class of the (simple) orbits of α_t and note that

$$\mathcal{A}^a(\alpha_t) = \{\pi((k-1)p+1)t^2/p; k \in \mathbb{N}\}.$$

Choose $\epsilon > 0$ sufficiently small such that $\pi R^2/p + \epsilon \notin \mathcal{A}(\alpha_R)$ and, by (2.3),

$$\text{HC}_*^{a, \pi R^2/p + \epsilon}(\alpha_R) \cong \text{H}_{* - \mu(\gamma)}(\mathbb{C}P^n; \mathbb{Q}), \quad (9.1)$$

where γ is a simple orbit of α_R (an easy computation shows that $\mu(\gamma) = (2n+2)/p - n$). From the discussion in Section 2.2, we conclude that this isomorphism is equivariant with respect to the shift operators D and Δ and therefore there exist non-zero elements $w_i \in \text{HC}_{(2n+2)/p - n + 2i}^{a, \pi R^2/p + \epsilon}(\alpha_0)$, $i = 0, \dots, n$, such that $Dw_{i+1} = w_i$ ($\Delta : H_*(\mathbb{C}P^n; \mathbb{Q}) \rightarrow H_{*-2}(\mathbb{C}P^n; \mathbb{Q})$ is an isomorphism).

We claim that ϵ can be chosen such that $\pi R^2/p + \epsilon \notin \mathcal{A}^a(\alpha_t)$ for every $t \in [r, R]$. As a matter of fact, note that if $k \geq 2$ then

$$\frac{\pi((k-1)p+1)t^2}{p} \geq \frac{\pi(p+1)r^2}{p} > \frac{\pi R^2}{p}$$

for every $t \in [r, R]$, where the last inequality holds because $(p+1)r^2 > R^2 \iff \frac{R^2}{r^2} < p+1$. Thus, it is enough to take $\epsilon > 0$ such that

$$\frac{\pi((k-1)p+1)t^2}{p} > \frac{\pi R^2}{p} + \epsilon$$

for every $t \in [r, R]$ and $k \geq 2$.

Thus, the continuation map

$$\phi_{\alpha_R, \alpha_r} : \mathrm{HC}_*^{a, \pi R^2/p+\epsilon}(\alpha_R) \rightarrow \mathrm{HC}_*^{a, \pi R^2/p+\epsilon}(\alpha_r)$$

is an isomorphism. Take ϵ such that $\pi R^2/p + \epsilon \notin \mathcal{A}^a(\alpha)$. From the commutative diagram (see (2.1))

$$\begin{array}{ccc} \mathrm{HC}_*^{a, \pi R^2/p+\epsilon}(\alpha_R) & \xrightarrow{\phi_{\alpha_R, \alpha_r}} & \mathrm{HC}_*^{a, \pi R^2/p+\epsilon}(\alpha_r) \\ & \searrow \phi_{\alpha_R, \alpha} & \nearrow \phi_{\alpha, \alpha_r} \\ & \mathrm{HC}_*^{a, \pi R^2/p+\epsilon}(\alpha) & \end{array}$$

we infer that $\phi_{\alpha_R, \alpha}$ is injective. Hence from the discussion in Section 2.2 and the equivariant isomorphism (9.1) we conclude that there exists an injective map (see (2.6))

$$\psi^T : \{0, \dots, n\} \rightarrow \mathcal{P}^{a, \pi R^2/p+\epsilon}(\alpha),$$

such that if $\gamma_i = \psi(i)$ then $A(\gamma_i) = c_{\phi_{\alpha_R, \alpha}(w_i)}(\alpha)$ and $\mathrm{HC}_{(2n+2)/p-n+2i}(\gamma_i) \neq 0$.

We claim that every γ_i is simple in its homotopy class in the sense that it cannot be an iterate of a periodic orbit with homotopy class a . Indeed, let β be the lift of α to S^{2n+1} . From the inequality $H_\beta(x) \leq r^{-2}\|x\|^2$ and the convexity of H_β we conclude from [12, Theorem A] that every periodic orbit α has period bigger than or equal to $\pi r^2/p$ (c.f. the proof of Lemma 8.1). Arguing by contradiction, assume that $\gamma_i = \gamma^k$ for some closed orbit γ with homotopy class a and $k \geq 2$. Then $k = jp + 1$ for some $j \geq 1$. Since $A(\gamma) \geq \pi r^2/p$, we arrive at

$$A(\gamma_i) = (jp + 1)A(\gamma) \geq (p + 1)\frac{\pi r^2}{p} > \pi R^2/p + \epsilon,$$

contradicting the fact that $\gamma_i \in \mathcal{P}^{a, \pi R^2/p+\epsilon}(\alpha)$. Consequently, the periodic orbits γ_i are geometrically distinct.

REFERENCES

- [1] M. Abreu and L. Macarini, *Contact homology of good toric contact manifolds*. *Compositio Mathematica* **148** (2012), 304–334.
- [2] M. Abreu, L. Macarini, *Dynamical convexity and elliptic periodic orbits for Reeb flows*. *Math. Ann.* **369** (2017), 331–386.
- [3] M. Abreu, L. Macarini, M. Moreira, *On contact invariants of non-simply connected Gorenstein toric contact manifolds*. Preprint arXiv:1812.10361, 2018.
- [4] P. Albers, D. Hein, J. Gutt, *Periodic Reeb orbits on prequantization bundles*. *J. Mod. Dyn.* **12** (2018), 123–150.
- [5] M. Arnaud, *Existence d’orbites périodiques complètement elliptiques des Hamiltoniens convexes présentant certaines symétries*. *C. R. Acad. Sci. Paris.* **328** (1999), 1035–1038.
- [6] F. Bourgeois, A. Oancea, *Fredholm theory and transversality for the parametrized and for the S^1 -invariant symplectic action*. *J. Eur. Math. Soc. (JEMS)* **12** (2010), no. 5, 1181–1229.
- [7] F. Bourgeois, A. Oancea, *The index of Floer moduli problems for parametrized action functionals*. *Geom. Dedicata* **165** (2013), 5–24.
- [8] F. Bourgeois, A. Oancea, *The Gysin exact sequence for S^1 -equivariant symplectic homology*. *J. Topol. Anal.* **5** (2013), no. 4, 361–407.
- [9] F. Bourgeois, A. Oancea, *S^1 -equivariant symplectic homology and linearized contact homology*. *Int. Math. Res. Not. IMRN* 2017, no. 13, 3849–3937.
- [10] J. Chaidez, O. Edtmair, *3d convex contact forms and the Ruelle invariant*. Preprint arXiv:2012.12869, 2020.

- [11] D. Cristofaro-Gardiner, M. Hutchings, *From one Reeb orbit to two*. J. Differential Geom. **102** (2016), no. 1, 25–36.
- [12] C. Croke, A. Weinstein, *Closed curves on convex hypersurfaces and periods of nonlinear oscillations*. Invent. Math. **64** (1981), no. 2, 199–202.
- [13] G. Dell’Antonio, B. D’Onofrio, I. Ekeland, *Periodic solutions of elliptic type for strongly nonlinear Hamiltonian systems*. The Floer memorial volume, 327–333, Progr. Math., 133, Birkhäuser, Basel, 1995.
- [14] H. Duan, H. Liu, *Multiplicity and ellipticity of closed characteristics on compact star-shaped hypersurfaces in \mathbb{R}^{2n}* . Calc. Var. Partial Differential Equations **56** (2017), no. 3, Art. 65, 30 pp.
- [15] I. Ekeland, J. Lasry, *On the number of periodic trajectories for a Hamiltonian flow on a convex energy surface*. Ann. of Math. **112** (1980), no. 2, 283–319.
- [16] V. Ginzburg, D. Hein, U. Hryniewicz, L. Macarini, *Closed Reeb orbits on the sphere and symplectically degenerate maxima*. Acta Math. Vietnam. **38** (2013), no. 1, 55–78.
- [17] V. Ginzburg, B. Gürel, *Periodic orbits of twisted geodesic flows and the Weinstein–Moser theorem*. Comment. Math. Helv. **84** (2009), 865–907.
- [18] V. Ginzburg, B. Gürel, *Lusternik–Schnirelmann theory and closed Reeb orbits*. Math. Z. **295** (2020), 515–582.
- [19] V. Ginzburg, L. Macarini, *Dynamical convexity and closed orbits on symmetric spheres*. Duke Math. J. **170** (2021), 1201–1250.
- [20] H. Hofer, K. Wysocki and E. Zehnder. *The dynamics of strictly convex energy surfaces in \mathbb{R}^4* . Ann. of Math. **148** (1998), 197–289.
- [21] U. Hryniewicz, L. Macarini, *Local contact homology and applications*. J. Topol. Anal. **7** (2015), 167–238.
- [22] E. Kerman, *Rigid constellations of closed Reeb orbits*. Compos. Math. **153** (2017), no. 11, 2394–2444.
- [23] E. Lerman, *Contact toric manifolds*, J. Symplectic Geom. **1** (2003), 785–828.
- [24] H. Liu, Y. Long, *The existence of two closed characteristics on every compact star-shaped hypersurface in \mathbb{R}^4* . Acta Math. Sin. (Engl. Ser.) **32** (2016), no. 1, 40–53.
- [25] C. Liu, Y. Long, C. Zhu, *Multiplicity of closed characteristics on symmetric convex hypersurfaces in \mathbb{R}^{2n}* . Math. Ann. **323** (2002), 201–215.
- [26] H. Liu, C. Wang, D. Zhang, *Elliptic and non-hyperbolic closed characteristics on compact convex P -cyclic symmetric hypersurfaces in \mathbb{R}^{2n}* . Calc. Var. Partial Differential Equations **59** (2020), no. 1, Paper No. 24, 20 pp.
- [27] Y. Long. *Bott formula of the Maslov-type index theory*. Pacific J. Math. **187** (1999), 113–149.
- [28] Y. Long, *Index Theory for Symplectic Paths with Applications*. Birkhäuser, Basel, 2002.
- [29] Y. Long, C. Zhu, *Closed characteristics on compact convex hypersurfaces in \mathbb{R}^{2n}* . Ann. of Math. **155** (2002), no. 2, 317–368.
- [30] M. McLean, *Reeb orbits and the minimal discrepancy of an isolated singularity*. Invent. Math. **204** (2016), no. 2, 505–594.
- [31] P. Seidel, *A biased view of symplectic cohomology*. In Current developments in mathematics, 2006, 211–253. Int. Press, Somerville, MA, 2008.
- [32] C. Viterbo, *Functors and computations in Floer homology with applications. I*. Geom. Funct. Anal. **9** (1999), no. 5, 985–1033.
- [33] W. Wang, *Existence of closed characteristics on compact convex hypersurfaces in \mathbb{R}^{2n}* . Calc. Var. Partial Differential Equations, **55** (2016), 1–25.

CENTER FOR MATHEMATICAL ANALYSIS, GEOMETRY AND DYNAMICAL SYSTEMS, INSTITUTO SUPERIOR TÉCNICO, UNIVERSIDADE DE LISBOA, AV. ROVISCO PAIS, 1049-001 LISBOA, PORTUGAL
E-mail address: mabreu@math.tecnico.ulisboa.pt

CENTER FOR MATHEMATICAL ANALYSIS, GEOMETRY AND DYNAMICAL SYSTEMS, INSTITUTO SUPERIOR TÉCNICO, UNIVERSIDADE DE LISBOA, AV. ROVISCO PAIS, 1049-001 LISBOA, PORTUGAL
E-mail address: macarini@math.tecnico.ulisboa.pt