

# Analysis 1, Solutions to Übungsblatt Nr. 7

---

## Problem 1 (Sequences: sup, inf, lim sup, lim inf)

In what follows, let  $n$  be a natural number greater than or equal to 1.

A preliminary observation, common to all the items below, is that  $(-1)^n = 1$  if  $n$  is even and  $(-1)^n = -1$  if  $n$  is odd.

(a) In light of the preliminary observation, note that

$$x_n = \begin{cases} (2 + \frac{3}{n}) & \text{if } n \text{ is odd,} \\ -(2 + \frac{3}{n}) & \text{if } n \text{ is even.} \end{cases}$$

The sequence  $(2 + \frac{3}{n})_{n \geq 1}$  is monotonically decreasing, and so the sequence  $(-(2 + \frac{3}{n}))_{n \geq 1}$  is monotonically increasing. Moreover we have that

$$\lim_{n \rightarrow \infty} (2 + \frac{3}{n}) = \inf_{n \geq 1} (2 + \frac{3}{n}) = 2 \text{ and } \lim_{n \rightarrow \infty} \left\{ -\left(2 + \frac{3}{n}\right) \right\} = \sup_{n \geq 1} \left\{ -\left(2 + \frac{3}{n}\right) \right\} = -2.$$

The proof of these facts entirely parallels the proof that  $\lim_{n \rightarrow \infty} \frac{1}{n} = 0$ , and as such will be omitted.

It follows that

$$\sup_{n \geq 1} x_n = x_1 = 2 + \frac{3}{1} = 5$$

and that

$$\inf_{n \geq 1} x_n = x_2 = -\left(2 + \frac{3}{2}\right) = -\frac{7}{2}.$$

Moreover,

$$\sup_{n \geq m} x_n = \begin{cases} 2 + \frac{3}{m} & \text{if } m \text{ is odd,} \\ 2 + \frac{3}{m+1} & \text{if } m \text{ is even,} \end{cases}$$

and

$$\inf_{n \geq m} x_n = \begin{cases} -(2 + \frac{3}{m+1}) & \text{if } m \text{ is odd,} \\ -(2 + \frac{3}{m}) & \text{if } m \text{ is even.} \end{cases}$$

It follows that

$$\limsup_{n \rightarrow \infty} x_n = \inf_{m \geq 1} \left( \sup_{n \geq m} x_n \right) = \begin{cases} \inf_{m \geq 1} \left( 2 + \frac{3}{m} \right) & \text{if } m \text{ is odd} \\ \inf_{m \geq 1} \left( 2 + \frac{3}{m+1} \right) & \text{if } m \text{ is even} \end{cases} = 2.$$

and that

$$\liminf_{n \rightarrow \infty} x_n = \sup_{m \geq 1} \left( \inf_{n \geq m} x_n \right) = \begin{cases} \sup_{m \geq 1} \left\{ -\left(2 + \frac{3}{m+1}\right) \right\} & \text{if } m \text{ is odd} \\ \sup_{m \geq 1} \left\{ -\left(2 + \frac{3}{m}\right) \right\} & \text{if } m \text{ is even} \end{cases} = -2.$$

(b) The crucial observation is that

$$x_n = \begin{cases} 6 & \text{if } n = 4k + 1 \text{ for some } k \in \mathbb{N}, \\ -4 & \text{if } n = 4k + 2 \text{ for some } k \in \mathbb{N}, \\ 0 & \text{if } n = 4k + 3 \text{ for some } k \in \mathbb{N}, \\ 2 & \text{if } n = 4k + 4 \text{ for some } k \in \mathbb{N}. \end{cases}$$

Assuming this, it follows at once that

$$\sup_{n \geq 1} x_n = \limsup_{n \rightarrow \infty} x_n = 6$$

and that

$$\inf_{n \geq 1} x_n = \liminf_{n \rightarrow \infty} x_n = -4.$$

The proof of the claim amounts to a straightforward analysis of the four different cases that arise. For instance, if  $n = 4k + 1$  for some natural number  $k$ , then

$$\begin{aligned} x_n &= 1 + 2(-1)^{n+1} + 3(-1)^{\frac{n(n-1)}{2}} = 1 + 2(-1)^{(4k+1)+1} + 3(-1)^{\frac{(4k+1)(4k)}{2}} \\ &= 1 + 2(-1)^{2(2k+1)} + 3(-1)^{2k(4k+1)} = 1 + 2 \cdot 1 + 3 \cdot 1 = 6 \end{aligned}$$

since both  $2(2k + 1)$  and  $2k(4k + 1)$  are even numbers. The analysis of the other three cases is similar and is left to the reader.

(c) Note that

$$x_n = \begin{cases} n & \text{if } n \text{ is even,} \\ \frac{1}{n} & \text{if } n \text{ is odd.} \end{cases}$$

In particular, the subsequence  $(x_{2n} = 2n)$  is monotonically increasing and unbounded, and

$$\sup_{n \geq 1} x_n = \limsup_{n \rightarrow \infty} x_n = \infty.$$

On the other hand, the subsequence  $(x_{2n+1} = \frac{1}{2n+1})$  is monotonically decreasing and satisfies

$$\lim_{n \rightarrow \infty} x_{2n+1} = 0.$$

It follows that

$$\inf_{n \geq 1} x_n = \liminf_{n \rightarrow \infty} x_n = 0.$$

## Problem 2 (Series)

(a) Start by noticing that, for every natural number  $n \geq 1$ ,

$$\frac{(n+1)^n}{n^{n+1}} = \frac{1}{n} \frac{(n+1)^n}{n^n} = \frac{1}{n} \left(1 + \frac{1}{n}\right)^n.$$

In Problem 4(a) from Übungsblatt 6 we showed that the sequence  $\alpha_n := (1 + \frac{1}{n})^n$  is monotonically increasing. In particular,  $\alpha_n \geq \alpha_1$  for every  $n \geq 1$ . Since  $\alpha_1 = 2$ , it follows that

$$\frac{(n+1)^n}{n^{n+1}} = \frac{1}{n} \left(1 + \frac{1}{n}\right)^n \geq \frac{2}{n},$$

and so

$$\sum_{n=1}^{\infty} \frac{(n+1)^n}{n^{n+1}} \geq \sum_{n=1}^{\infty} \frac{2}{n} = 2 \sum_{n=1}^{\infty} \frac{1}{n}. \quad (1)$$

We know from Problem 3(c) of Übungsblatt 6 that the harmonic series  $\sum_{n=1}^{\infty} \frac{1}{n}$  diverges. From (1), it follows that the series

$$\sum_{n=1}^{\infty} \frac{(n+1)^n}{n^{n+1}} = \infty$$

diverges as well.

(b) For  $n \in \mathbb{N}$ , define  $a_n := \frac{n^3}{n!}$ . Let us compute the quotient

$$\frac{a_{n+1}}{a_n} = \frac{(n+1)^3}{(n+1)!} \frac{n!}{n^3} = \frac{\cancel{(n+1)}(n+1)^2}{\cancel{(n+1)}n!} \frac{n!}{n^3} = \frac{(n+1)^2}{n!} \frac{n!}{n^3} = \frac{(n+1)^2}{n^3} = \frac{1}{n} \left(1 + \frac{1}{n}\right)^2.$$

Observe that the sequence

$$\frac{a_{n+1}}{a_n} = \frac{1}{n} \left(1 + \frac{1}{n}\right)^2 = \frac{1}{n} \left(1 + \frac{2}{n} + \frac{1}{n^2}\right) = \frac{1}{n} + \frac{2}{n^2} + \frac{1}{n^3}$$

is monotonically decreasing and that

$$\frac{a_4}{a_3} = \frac{1}{3} \left(1 + \frac{1}{3}\right)^2 = \frac{16}{27} < 1.$$

It follows that

$$\frac{a_{n+1}}{a_n} \leq \frac{16}{27}$$

for every  $n \geq 3$ , and so the conditions of Problem 2(b), Übungsblatt 6 are fulfilled (for  $N = 3$  and  $y = 16/27$ ). We can use that result to establish the convergence of the sequence

$$\sum_{n=0}^{\infty} \frac{n^3}{n!} < \infty.$$

(c) We proceed similarly to part (b). For  $n \in \mathbb{N}$ , define  $b_n := \frac{(n!)^2}{(2n)!}$ . Then

$$\frac{b_{n+1}}{b_n} = \frac{((n+1)!)^2 (2n)!}{(2(n+1))! (n!)^2} = \frac{(n+1) \cancel{n!} (n+1) \cancel{n!} (2n)!}{(2n+2)! \cancel{n!} \cancel{n!}} = (n+1)^2 \frac{(2n)!}{(2n+2)(2n+1)(2n)!} = \frac{(n+1)^2}{(2n+2)(2n+1)}.$$

Again, the sequence

$$\frac{b_{n+1}}{b_n} = \frac{(n+1)^2}{(2n+2)(2n+1)} = \frac{1}{2} \frac{n+1}{2n+1} = \frac{1}{2} \left(1 - \frac{n}{2n+1}\right) = \frac{1}{2} \left(1 - \frac{1}{2 + \frac{1}{n}}\right)$$

is monotonically decreasing, and

$$\frac{b_1}{b_0} = \frac{(0+1)^2}{(2 \cdot 0 + 2)(2 \cdot 0 + 1)} = \frac{1}{2} < 1.$$

Problem 2(b) from Übungsblatt 6 (with  $N = 0$  and  $y = 1/2$ ) implies that the series

$$\sum_{n=0}^{\infty} \frac{(n!)^2}{(2n)!} < \infty$$

converges.

(d) Start by observing that, for every natural number  $n \geq 1$ ,

$$\frac{\sqrt{n+1} - \sqrt{n}}{n} = \frac{(\sqrt{n+1} - \sqrt{n})(\sqrt{n+1} + \sqrt{n})}{n(\sqrt{n+1} + \sqrt{n})} = \frac{(n+1) - n}{n(\sqrt{n+1} + \sqrt{n})} = \frac{1}{n(\sqrt{n+1} + \sqrt{n})}.$$

Moreover, we have that

$$\frac{1}{n(\sqrt{n+1} + \sqrt{n})} \leq \frac{1}{n\sqrt{n}}$$

since  $\sqrt{n} \leq \sqrt{n+1} + \sqrt{n}$ . Since this holds for every natural  $n \geq 1$ , it follows that

$$\sum_{n=1}^{\infty} \frac{1}{n(\sqrt{n+1} + \sqrt{n})} \leq \sum_{n=1}^{\infty} \frac{1}{n\sqrt{n}}.$$

From Problem 3(c), Übungsblatt 6, we know that the series

$$\sum_{n=1}^{\infty} \frac{1}{n\sqrt{n}} = \sum_{n=1}^{\infty} \frac{1}{n^{3/2}} < \infty$$

converges since  $3/2 > 1$ . It follows that the series

$$\sum_{n=1}^{\infty} \frac{1}{n(\sqrt{n+1} + \sqrt{n})} < \infty$$

converges as well.

**Problem 3 (Fubini without Tonelli)**

Let  $f = (f_n(m))_{m,n}$ , where  $m \in \mathbb{N}$  designates the number of the (horizontal) row and  $n \in \mathbb{N}$  designates the number of the (vertical) column, be the following infinite matrix:

$$\begin{pmatrix} 0 & \frac{1}{2} & \frac{1}{4} & \frac{1}{8} & \frac{1}{16} & \frac{1}{32} & \cdots \\ -\frac{1}{2} & 0 & \frac{1}{2} & \frac{1}{4} & \frac{1}{8} & \frac{1}{16} & \cdots \\ -\frac{1}{4} & -\frac{1}{2} & 0 & \frac{1}{2} & \frac{1}{4} & \frac{1}{8} & \cdots \\ -\frac{1}{8} & -\frac{1}{4} & -\frac{1}{2} & 0 & \frac{1}{2} & \frac{1}{4} & \cdots \\ \vdots & \vdots & \vdots & \vdots & \vdots & \vdots & \ddots \end{pmatrix}.$$

In other words,

$$f_n(m) = \begin{cases} 2^{m-n} & \text{if } n > m, \\ 0 & \text{if } n = m, \\ -2^{n-m} & \text{if } n < m. \end{cases}$$

It follows that, for  $m = 0, 1, 2, \dots$ ,

$$\begin{aligned} \sum_{n=0}^{\infty} f_n(m) &= \sum_{n=0}^{m-1} (-2^{n-m}) + 0 + \sum_{n=m+1}^{2m} 2^{m-n} + \sum_{n=2m+1}^{\infty} 2^{m-n} = \sum_{n=2m+1}^{\infty} 2^{m-n} \\ &= \sum_{k=1}^{\infty} 2^{-m-k} = 2^{-m} \sum_{k=1}^{\infty} 2^{-k} = 2^{-m} \frac{1}{2} \sum_{k=0}^{\infty} 2^{-k} = 2^{-m} \frac{1}{2} \frac{1}{1-1/2} = 2^{-m}, \end{aligned}$$

where in the passage from the first to the second line we changed summation index  $k = n - 2m$ . It follows that

$$\sum_{m=0}^{\infty} \sum_{n=0}^{\infty} f_n(m) = \sum_{m=0}^{\infty} 2^{-m} = \frac{1}{1-1/2} = 2.$$

On the other hand, for  $n = 0, 1, 2, \dots$ ,

$$\sum_{m=0}^{\infty} f_n(m) = -2^{-n} \frac{1}{2} \sum_{k=0}^{\infty} 2^{-k} = -2^{-n},$$

and so

$$\sum_{n=0}^{\infty} \sum_{m=0}^{\infty} f_n(m) = \sum_{n=0}^{\infty} (-2^{-n}) = -2.$$

Since  $2 \neq -2$ , the sequence (of sequences)  $f$  does the required job. Note that, in light of Theorem 3.40 (Fubini-Tonelli), such a result could never hold if instead we considered sequences of (extended) positive real numbers.

**Problem 4 (A – perhaps surprising? – convergent series)**

For a natural number  $n \geq 1$ , define the set

$$S_n := \{k \in \mathbb{N} : k \geq 1 \text{ and } k \text{ has exactly } n \text{ digits and none of them is } 9\}.$$

Our first claim is that *the set  $S_n$  has  $8 \cdot 9^{n-1}$  elements*. This can be verified via induction on  $n$ . For  $n = 1$ , we have that

$$S_1 = \{1, 2, 3, 4, 5, 6, 7, 8\},$$

and so  $S_1$  has  $8 = 8 \cdot 9^{1-1}$  elements, as desired. Let us assume that the set  $S_n$  has  $8 \cdot 9^{n-1}$  elements, and prove that the set  $S_{n+1}$  has  $8 \cdot 9^n$  elements. Any elements of  $S_{n+1}$  can be obtained in a unique way from an element of  $S_n$  by concatenating a single digit to its end. Conversely, any element of  $S_n$  gives rise to an element of  $S_{n+1}$  by concatenation of a single digit to its end. For each element of  $S_n$ , there are 9 possibilities for doing this (one chooses one of 0,1,2,3,4,5,6,7,8). It follows that the number of elements of  $S_{n+1}$  is exactly 9 times the number of elements of  $S_n$ , which by induction hypothesis is  $8 \cdot 9^{n-1}$ . Therefore the number of elements of the set  $S_{n+1}$  is  $9 \cdot (8 \cdot 9^{n-1}) = 8 \cdot 9^n$ , as desired.

Our next observation is that *any element of  $S_n$  is at least  $10^{n-1}$* . This follows from noting that

$$10^{n-1} = \underbrace{100\dots0}_{n \text{ digits}} \in S_n,$$

and that

$$10^{n-1} - 1 = \underbrace{99\dots9}_{n-1 \text{ digits}} \in S_{n-1}.$$

As a consequence, the elements of the set  $S_n$  contribute to the original sum by less than  $\frac{8 \cdot 9^{n-1}}{10^{n-1}}$ . Indeed, denoting the number of elements of the set  $S_n$  by  $\#S_n$ , we have that

$$\sum_{k \in S_n} \frac{\epsilon_n}{n} \leq \sum_{k \in S_n} \frac{1}{10^{n-1}} = \frac{1}{10^{n-1}} \sum_{k \in S_n} 1 = \frac{1}{10^{n-1}} (\#S_n) = \frac{8 \cdot 9^{n-1}}{10^{n-1}}.$$

We can finally conclude that

$$\sum_{n=1}^{\infty} \frac{\epsilon_n}{n} = \sum_{n=1}^{\infty} \left( \sum_{k \in S_n} \frac{\epsilon_n}{n} \right) \leq \sum_{n=1}^{\infty} \frac{8 \cdot 9^{n-1}}{10^{n-1}} = 8 \sum_{n=1}^{\infty} \left( \frac{9}{10} \right)^{n-1} = 8 \sum_{m=1}^{\infty} \left( \frac{9}{10} \right)^m = 8 \frac{1}{1 - 9/10} = 80 < \infty.$$

Note that we proved a stronger result. Not only does the series converge, but we also have the quantitative estimate

$$\sum_{n=1}^{\infty} \frac{\epsilon_n}{n} \leq 80.$$

As a final remark, let us mention that this series is an example of what in the literature goes under the name of “Kempner series”. It can be shown that, up to 10 decimals, the actual sum is 22.9206766193, but the methods for proving that result are beyond the scope of this class.