

# Modular class and integrability in Poisson and related geometries

Raquel Caseiro

CMUC, Department of Mathematics  
University of Coimbra

XVI IFWGP 2007

# Outline

- 1 Modular class on Poisson manifolds
  - Curl operator
  - Properties of Modular class
- 2 Modular classes on Lie algebroids
  - Definition
  - Examples
  - Cohomology pairing
  - Generalizing the curl operator
- 3 Modular classes of Nijenhuis operators
  - Relative modular class
  - Poisson-Nijenhuis algebroids
  - Jacobi-Nijenhuis algebroids
- 4 References

## It was used by...

- J. L. Koszul (1985)
- J. P. Dufour and A. Haraki (1991)
- Z. -L. Liu and P. Xu (1992)
- J. Grabowski, G. Marmo and A. M. Perelomov (1993)

Classify quadratic Poisson structures

# What is the Curl Operator?

On a Poisson manifold  $(M, \pi)$  choose a volume form  $\mu$ :

$$\phi : \mathfrak{X}^k(M) \xrightarrow{\cong} \Omega^{n-k}(M), \quad P \longmapsto i_P \mu$$

# What is the Curl Operator?

On a Poisson manifold  $(M, \pi)$  choose a volume form  $\mu$ :

$$\phi : \mathfrak{X}^k(M) \xrightarrow{\cong} \Omega^{n-k}(M), \quad P \longmapsto i_P \mu$$

- The **curl operator** is defined as

$$\begin{array}{ccccccc}
 D_\mu : \mathfrak{X}^k(M) & \rightarrow & \Omega^{n-k}(M) & \rightarrow & \Omega^{n-k+1}(M) & \rightarrow & \mathfrak{X}^{k-1}(M) \\
 P & \xrightarrow{\phi} & i_P \mu & \xrightarrow{d} & di_P \mu & \xrightarrow{\phi^{-1}} & \underbrace{D_\mu P}_{\text{curl of } P}
 \end{array}$$

# What is the Curl Operator?

On a Poisson manifold  $(M, \pi)$  choose a volume form  $\mu$ :

$$\phi : \mathfrak{X}^k(M) \xrightarrow{\cong} \Omega^{n-k}(M), \quad P \longmapsto i_P \mu$$

- The **curl operator** is defined as

$$\begin{array}{ccccccc}
 D_\mu : \mathfrak{X}^k(M) & \rightarrow & \Omega^{n-k}(M) & \rightarrow & \Omega^{n-k+1}(M) & \rightarrow & \mathfrak{X}^{k-1}(M) \\
 P & \xrightarrow{\phi} & i_P \mu & \xrightarrow{d} & di_P \mu & \xrightarrow{\phi^{-1}} & \underbrace{D_\mu P}_{\text{curl of } P}
 \end{array}$$

- The curl of a vector field is its divergence with respect to  $\mu$ :

$$D_\mu X = \operatorname{div}_\mu(X) = \frac{\mathcal{L}_X \mu}{\mu}, \quad X \in \mathfrak{X}^1(M).$$

## Definition

The **modular vector field** of  $(M, \pi)$  is the curl of a Poisson bivector  $\pi$ :

$$X_\mu = D_\mu \pi = \phi^{-1}(\mathbf{d}i_\pi \mu).$$

## Definition

The **modular vector field** of  $(M, \pi)$  is the curl of a Poisson bivector  $\pi$ :

$$X_\mu = D_\mu \pi = \phi^{-1}(\mathbf{d}i_\pi \mu).$$

## Remark

This modular vector field depends on the choice of  $\mu$ .



# Weinstein geometric approach

## Question

Given  $(M, \pi)$  is there a volume form  $\mu$  invariant under the flow of all hamiltonian vector fields, i.e, such that:

$$\mathcal{L}_{X_f}\mu = 0, \forall f \in C^\infty(M)?$$

# Weinstein geometric approach

## Question

Given  $(M, \pi)$  is there a volume form  $\mu$  invariant under the flow of all hamiltonian vector fields, i.e, such that:

$$\mathcal{L}_{X_f}\mu = 0, \forall f \in C^\infty(M)?$$

The modular vector field relative to  $\mu$  gives:

$$X_\mu f = \mathcal{L}_{X_f}\mu/\mu = \operatorname{div}_\mu \pi^\sharp(df).$$

# Weinstein geometric approach

## Question

Given  $(M, \pi)$  is there a volume form  $\mu$  invariant under the flow of all hamiltonian vector fields, i.e, such that:

$$\mathcal{L}_{X_f}\mu = 0, \forall f \in C^\infty(M)?$$

The modular vector field relative to  $\mu$  gives:

$$X_\mu f = \mathcal{L}_{X_f}\mu/\mu = \operatorname{div}_\mu \pi^\sharp(df).$$

- $X_\mu$  is a 1-cocycle in Poisson cohomology

$$d_\pi X_\mu = \mathcal{L}_{X_\mu}\pi = 0.$$

# Weinstein geometric approach

## Question

Given  $(M, \pi)$  is there a volume form  $\mu$  invariant under the flow of all hamiltonian vector fields, i.e, such that:

$$\mathcal{L}_{X_f}\mu = 0, \forall f \in C^\infty(M)?$$

The modular vector field relative to  $\mu$  gives:

$$X_\mu f = \mathcal{L}_{X_f}\mu / \mu = \operatorname{div}_\mu \pi^\sharp(df).$$

- $X_\mu$  is a 1-cocycle in Poisson cohomology

$$d_\pi X_\mu = \mathcal{L}_{X_\mu} \pi = 0.$$

- If  $\nu = f\mu$  then:  $X_\nu = X_\mu + \underbrace{\pi^\sharp(-d \ln |f|)}_{\text{Hamiltonian vector field}}$

Hamiltonian vector field

# Modular Class

## Definition

- $\text{mod}(\pi) = [X_\mu] \in H_\pi^1(M)$  is the **modular class of  $(M, \pi)$** .

# Modular Class

## Definition

- $\text{mod}(\pi) = [X_\mu] \in H_\pi^1(M)$  is the **modular class of  $(M, \pi)$** .
- $(M, \pi)$  is called **unimodular** if  $\text{mod}(\pi) = 0$  (i.e. if there exists a volume form invariant for all Hamiltonian vector fields).

# Modular Class

## Definition

- $\text{mod}(\pi) = [X_\mu] \in H_\pi^1(M)$  is the **modular class of  $(M, \pi)$** .
- $(M, \pi)$  is called **unimodular** if  $\text{mod}(\pi) = 0$  (i.e. if there exists a volume form invariant for all Hamiltonian vector fields).

## Remarks

- The Poisson cohomology groups are usually very hard to compute.

# Modular Class

## Definition

- $\text{mod}(\pi) = [X_\mu] \in H_\pi^1(M)$  is the **modular class of  $(M, \pi)$** .
- $(M, \pi)$  is called **unimodular** if  $\text{mod}(\pi) = 0$  (i.e. if there exists a volume form invariant for all Hamiltonian vector fields).

## Remarks

- The Poisson cohomology groups are usually very hard to compute.
- The modular class is a basic invariant, with geometric meaning, which is easy to determine in many examples.



# Examples

- Any **symplectic manifold** is unimodular: The Liouville form is invariant under all Hamiltonian flows.

## Examples

- Any **symplectic manifold** is unimodular: The Liouville form is invariant under all Hamiltonian flows.
- $M = \mathfrak{g}^*$  with Lie-Poisson structure  $\pi$ :  
For a translation invariant measure  $\mu$ , the modular vector field is the modular character of  $\mathfrak{g}$ :

$$\xi(\mathfrak{g}) = \text{Tr ad } \mathfrak{g}, \quad \mathfrak{g} \in \mathfrak{g}.$$

## Examples

- Any **symplectic manifold** is unimodular: The Liouville form is invariant under all Hamiltonian flows.
- $M = \mathfrak{g}^*$  with Lie-Poisson structure  $\pi$ :  
For a translation invariant measure  $\mu$ , the modular vector field is the modular character of  $\mathfrak{g}$ :

$$\xi(\mathfrak{g}) = \text{Tr ad } \mathfrak{g}, \quad \mathfrak{g} \in \mathfrak{g}.$$

- For a regular Poisson manifold, with transversally oriented symplectic foliation  $\mathcal{F}$ , one can show:

$$i^*(\text{mod}(\mathcal{F})) = \text{mod}(\pi), \quad i^* : H^1(\mathcal{F}) \rightarrow H^1_\pi(M).$$

# Cohomology and Homology

- Poisson cohomology  $H^\bullet(M, \pi): d_\pi = [\pi, -]$ ;

# Cohomology and Homology

- Poisson cohomology  $H^\bullet(M, \pi): d_\pi = [\pi, -]$ ;
- Poisson homology  $H_\bullet(M, \pi): \delta_\pi = [i_\pi, d]$ ;

# Cohomology and Homology

- Poisson cohomology  $H^\bullet(M, \pi)$ :  $d_\pi = [\pi, -]$ ;
- Poisson homology  $H_\bullet(M, \pi)$ :  $\delta_\pi = [i_\pi, d]$ ;

Theorem ((Evens, Lu and Weinstein, 1998), (Xu, 1999))

*If  $(M, \pi)$  is unimodular then*

$$H_\bullet(M, \pi) \cong H^{n-\bullet}(M, \pi).$$

# Holonomy and the modular class

Theorem (Ginzburg and Golubev, 2001)

For any cotangent loop  $a : I \rightarrow T^*M$ :

$$\det h(a) = \exp\left(\int_a \text{mod}(\pi)\right).$$

# Holonomy and the modular class

## Theorem (Ginzburg and Golubev, 2001)

For any cotangent loop  $a : I \rightarrow T^*M$ :

$$\det h(a) = \exp\left(\int_a \text{mod}(\pi)\right).$$

## Remarks

- $h$  denotes the **linear** Poisson holonomy (defined using parallel transport relative to contravariant version of the Bott connection).



# Holonomy and the modular class

Theorem (Ginzburg and Golubev, 2001)

For any cotangent loop  $a : I \rightarrow T^*M$ :

$$\det h(a) = \exp\left(\int_a \text{mod}(\pi)\right).$$

## Remarks

- $h$  denotes the **linear** Poisson holonomy (defined using parallel transport relative to contravariant version of the Bott connection).
- Integration of Poisson vector field over cotangent paths is invariant under **cotangent homotopies**.

## Following Evans, Lu and Weinstein (1998)

Substitute for volume form:  $\eta \otimes \mu$  section of the line bundle  
 $Q_A = \wedge^{\text{top}} A \otimes \wedge^{\text{top}}(T^*M).$

## Following Evans, Lu and Weinstein (1998)

Substitute for volume form:  $\eta \otimes \mu$  section of the line bundle  
 $Q_A = \wedge^{\text{top}} A \otimes \wedge^{\text{top}}(T^*M)$ .

### Definition

The **modular form** of the Lie algebroid  $A$  with respect to  $\eta \otimes \mu$  is the  $A$ -form  $\xi_A \in \Omega^1(A)$  defined by

$$\langle \xi_A, X \rangle \eta \otimes \mu = [X, \eta] \otimes \mu + \mu \otimes \mathcal{L}_{\rho(X)}\mu, \quad X \in \Gamma(A).$$

## Following Evans, Lu and Weinstein (1998)

Substitute for volume form:  $\eta \otimes \mu$  section of the line bundle  
 $Q_A = \wedge^{\text{top}} A \otimes \wedge^{\text{top}}(T^*M)$ .

### Definition

The **modular form** of the Lie algebroid  $A$  with respect to  $\eta \otimes \mu$  is the  $A$ -form  $\xi_A \in \Omega^1(A)$  defined by

$$\langle \xi_A, X \rangle \eta \otimes \mu = [X, \eta] \otimes \mu + \mu \otimes \mathcal{L}_{\rho(X)}\mu, \quad X \in \Gamma(A).$$

- $\nabla : \Gamma(A) \times \Gamma(Q_A) \rightarrow \Gamma(Q_A)$   
 $(X, \eta \otimes \mu) \rightarrow [X, \eta] \otimes \mu + \eta \otimes \mathcal{L}_{\rho(X)}\mu$   
 is a representation of the Lie algebroid  $A$  on  $Q_A$ .

## Following Evans, Lu and Weinstein (1998)

- $\xi_A$  is a 1-cocycle in the Lie algebroid cohomology

$$d_A \xi_A = 0$$

## Following Evans, Lu and Weinstein (1998)

- $\xi_A$  is a 1-cocycle in the Lie algebroid cohomology

$$d_A \xi_A = 0$$

- If  $\eta' \otimes \mu' = f\eta \otimes \mu$  is another section of  $Q_A$  then

$$\xi'_A = \xi_A - \underbrace{d_A \log f}_{\text{coboundary}} .$$

## Following Evans, Lu and Weinstein (1998)

- $\xi_A$  is a 1-cocycle in the Lie algebroid cohomology

$$d_A \xi_A = 0$$

- If  $\eta' \otimes \mu' = f\eta \otimes \mu$  is another section of  $Q_A$  then

$$\xi'_A = \xi_A - \underbrace{d_A \log f}_{\text{coboundary}} .$$

### Definition

$[\xi_A] \in H^1(A)$  is the **modular class** of the Lie algebroid  $A$ .

# Tangent Bundle: $A = TM$

Assume  $M$  is orientable.

- $\Gamma(Q_{TM}) = \mathfrak{X}^{\text{top}}(M) \otimes \Omega^{\text{top}}(M)$



# Tangent Bundle: $A = TM$

Assume  $M$  is orientable.

- $\Gamma(Q_{TM}) = \mathfrak{X}^{\text{top}}(M) \otimes \Omega^{\text{top}}(M)$
- $P \otimes \mu$  such that  $\langle \mu, P \rangle = 1$

# Tangent Bundle: $A = TM$

Assume  $M$  is orientable.

- $\Gamma(Q_{TM}) = \mathfrak{X}^{\text{top}}(M) \otimes \Omega^{\text{top}}(M)$
- $P \otimes \mu$  such that  $\langle \mu, P \rangle = 1$
- $\xi_{TM} = 0$

# Tangent Bundle: $A = TM$

Assume  $M$  is orientable.

- $\Gamma(Q_{TM}) = \mathfrak{X}^{\text{top}}(M) \otimes \Omega^{\text{top}}(M)$
- $P \otimes \mu$  such that  $\langle \mu, P \rangle = 1$
- $\xi_{TM} = 0$

If  $M$  is orientable then  $\text{mod } TM = 0$ .

# Lie algebra: $A = \mathfrak{g}$

A Lie algebra  $\mathfrak{g}$  is a Lie algebroid over one point.

# Lie algebra: $A = \mathfrak{g}$

A Lie algebra  $\mathfrak{g}$  is a Lie algebroid over one point.

- $Q_{\mathfrak{g}} = \wedge^{\text{top}} \mathfrak{g}$

# Lie algebra: $A = \mathfrak{g}$

A Lie algebra  $\mathfrak{g}$  is a Lie algebroid over one point.

- $Q_{\mathfrak{g}} = \wedge^{\text{top}} \mathfrak{g}$
- The modular form is the adjoint character

$$\langle \xi_{\mathfrak{g}}, g \rangle = \text{Tr ad}_g, \quad g \in \mathfrak{g}.$$

## Lie algebra: $A = \mathfrak{g}$

A Lie algebra  $\mathfrak{g}$  is a Lie algebroid over one point.

- $Q_{\mathfrak{g}} = \wedge^{\text{top}} \mathfrak{g}$
- The modular form is the adjoint character

$$\langle \xi_{\mathfrak{g}}, g \rangle = \text{Tr } \text{ad}_g, \quad g \in \mathfrak{g}.$$

$$\text{mod } \mathfrak{g} = \text{mod } (\mathfrak{g}^*, \pi_{\text{Lie}})$$

# Foliation: $A = T\mathcal{F}$

- $T\mathcal{F}$ : integrable subbundle of  $TM$  with orientable conormal bundle  $N$ .



## Foliation: $A = T\mathcal{F}$

- $T\mathcal{F}$ : integrable subbundle of  $TM$  with orientable conormal bundle  $N$ .
- $Q_A \cong \wedge^{\text{top}} N$

## Foliation: $A = T\mathcal{F}$

- $T\mathcal{F}$ : integrable subbundle of  $TM$  with orientable conormal bundle  $N$ .
- $Q_A \cong \wedge^{\text{top}} N$

$$\text{mod } T\mathcal{F} = \text{mod } \mathcal{F}.$$

# Poisson manifold: $A = T^*M$

- $(M, \pi)$  is a Poisson manifold

## Poisson manifold: $A = T^*M$

- $(M, \pi)$  is a Poisson manifold
- $T^*M$  is a Lie algebroid over  $M$  with anchor  $\rho = \pi^\sharp$  and Lie bracket

$$[\alpha, \beta] = \mathcal{L}_{\pi^\sharp \alpha} \beta - \mathcal{L}_{\pi^\sharp \beta} \alpha - d\pi(\alpha, \beta), \quad \alpha, \beta \in \Omega^1(M).$$

## Poisson manifold: $A = T^*M$

- $(M, \pi)$  is a Poisson manifold
- $T^*M$  is a Lie algebroid over  $M$  with anchor  $\rho = \pi^\sharp$  and Lie bracket

$$[\alpha, \beta] = \mathcal{L}_{\pi^\sharp \alpha} \beta - \mathcal{L}_{\pi^\sharp \beta} \alpha - d\pi(\alpha, \beta), \quad \alpha, \beta \in \Omega^1(M).$$

$$\text{mod } T^*M = 2\text{mod}(\pi)$$

# Cohomology and Homology

- Pairing between  $\underbrace{H^\bullet(A, d_A)}_{\text{trivial coefficients}}$  and  $\underbrace{H^{r-\bullet}(A, Q_A)}_{\text{coefficients on } Q_A}$ ,

# Cohomology and Homology

- Pairing between  $\underbrace{H^\bullet(A, d_A)}_{\text{trivial coefficients}}$  and  $\underbrace{H^{r-\bullet}(A, Q_A)}_{\text{coefficients on } Q_A}$ ,
- $(M, \pi)$  Poisson manifold.

Pairing  $H_\bullet(M)$  and  $H_{n-\bullet}(M, \pi)$

- P. Xu (1999) relates the study of modular vector fields on Poisson manifold with the notion of Gerstenhaber algebra.



- P. Xu (1999) relates the study of modular vector fields on Poisson manifold with the notion of Gerstenhaber algebra.
- Y. Kosmann-Schwarzbach (2000) use this to define a modular vector field for triangular Lie bialgebroids.

- P. Xu (1999) relates the study of modular vector fields on Poisson manifold with the notion of Gerstenhaber algebra.
- Y. Kosmann-Schwarzbach (2000) use this to define a modular vector field for triangular Lie bialgebroids.
- Extended to Lie-Rienart algebras by Huebschmann (1999).

# Generators of Gerstenhaber algebras

## Definition

An operator  $\partial$  of degree  $-1$  is a **generator of the Gerstenhaber algebra**  $\mathfrak{X}(A)$  if, for  $P \in \mathfrak{X}^p(A)$ ,  $Q \in \mathfrak{X}(A)$ ,

$$[P, Q] = (-1)^p (\partial(P \wedge Q) - \partial P \wedge Q - (-1)^p P \wedge \partial Q).$$

If  $\partial^2 = 0$  then  $\mathfrak{X}(A)$  is **exact** or a Batalin-Vilkosky algebra.

# Generators of Gerstenhaber algebras

## Definition

An operator  $\partial$  of degree  $-1$  is a **generator of the Gerstenhaber algebra**  $\mathfrak{X}(A)$  if, for  $P \in \mathfrak{X}^p(A)$ ,  $Q \in \mathfrak{X}(A)$ ,

$$[P, Q] = (-1)^p (\partial(P \wedge Q) - \partial P \wedge Q - (-1)^p P \wedge \partial Q).$$

If  $\partial^2 = 0$  then  $\mathfrak{X}(A)$  is **exact** or a Batalin-Vilkosky algebra.

## P. Xu (1999)

Let  $\partial$  and  $\partial'$  be two generators of the Gerstenhaber algebra  $\mathfrak{X}(A)$ , such that  $\partial^2 = \partial'^2 = 0$ . Then

$$\partial - \partial' = i_\alpha,$$

for a closed 1-form  $\alpha \in \Gamma(A^*)$ .

## What happens with triangular Lie bialgebroids?

- $P \in \mathfrak{X}^2(A)$  a Poisson bivector:  $[P, P] = 0$ ;

# What happens with triangular Lie bialgebroids?

- $P \in \mathfrak{X}^2(A)$  a Poisson bivector:  $[P, P] = 0$ ;
- $A^*$  is also a Lie algebroid;

## What happens with triangular Lie bialgebroids?

- $P \in \mathfrak{X}^2(A)$  a Poisson bivector:  $[P, P] = 0$ ;
- $A^*$  is also a Lie algebroid;
- $\partial_P = [i_P, d_A] = i_A d_A - d_A i_A$  is a generator of  $\Omega(A) = \mathfrak{X}(A^*)$ .

# What happens with triangular Lie bialgebroids?

- $P \in \mathfrak{X}^2(A)$  a Poisson bivector:  $[P, P] = 0$ ;
- $A^*$  is also a Lie algebroid;
- $\partial_P = [i_P, d_A] = i_A d_A - d_A i_A$  is a generator of  $\Omega(A) = \mathfrak{X}(A^*)$ .

Let  $\mu$  be a top-section of  $A^*$ . It defines another generator of  $\Omega(A)$ :

$$\partial_{P,\mu} = -\phi d_P \phi^{-1}$$

where  $\phi Q = i_Q \mu$ ,  $Q \in \mathfrak{X}(A)$ .



The **modular vector field** of a triangular Lie bialgebroid associated with  $\mu$  is the vector field  $X_\mu$  given by

$$\partial_{P,\mu} - \partial_P = i_{X_\mu}.$$

The **modular vector field** of a triangular Lie bialgebroid associated with  $\mu$  is the vector field  $X_\mu$  given by

$$\partial_{P,\mu} - \partial_P = i_{X_\mu}.$$

- $d_P X_\mu = 0$ ;

The **modular vector field** of a triangular Lie bialgebroid associated with  $\mu$  is the vector field  $X_\mu$  given by

$$\partial_{\mathcal{P},\mu} - \partial_{\mathcal{P}} = i_{X_\mu}.$$

- $d_{\mathcal{P}}X_\mu = 0$ ;
- $X_{f\mu} = X_\mu + d_{\mathcal{P}} \ln |f|$ ;

The **modular vector field** of a triangular Lie bialgebroid associated with  $\mu$  is the vector field  $X_\mu$  given by

$$\partial_{P,\mu} - \partial_P = i_{X_\mu}.$$

- $d_P X_\mu = 0$ ;
- $X_{f\mu} = X_\mu + d_P \ln |f|$ ;

$\text{mod}(A, P) = [X_\mu] \in H^1(A^*)$  is the **modular class** of  $(A, P)$ .

The **modular vector field** of a triangular Lie bialgebroid associated with  $\mu$  is the vector field  $X_\mu$  given by

$$\partial_{P,\mu} - \partial_P = i_{X_\mu}.$$

- $d_P X_\mu = 0$ ;
- $X_{f\mu} = X_\mu + d_P \ln |f|$ ;

$\text{mod}(A, P) = [X_\mu] \in H^1(A^*)$  is the **modular class** of  $(A, P)$ .

- If  $(A, P)$  unimodular then

$$H_\bullet(A, \delta_P) \cong H^{\text{top} - \bullet}(A, d_A).$$

## Relation between the two generalizations

- If  $(M, \pi)$  is Poisson manifold:

$$\text{mod}(TM, \pi) = \text{mod}(\pi) = \frac{1}{2} \text{mod } T^*M.$$

## Relation between the two generalizations

- If  $(M, \pi)$  is Poisson manifold:

$$\text{mod}(TM, \pi) = \text{mod}(\pi) = \frac{1}{2} \text{mod } T^*M.$$

- For general triangular bialgebroids

$$\text{mod}(A, P) \neq \frac{1}{2} \text{mod } A^*$$

## Y. Kosmann-Schwarzbach and A. Weinstein (2005)

The **relative modular class** of the Lie algebroid morphism  $\varphi : A \rightarrow B$  is

$$\text{mod } \varphi(A, B) = \text{mod } A - \varphi^*(\text{mod } B)$$



## Y. Kosmann-Schwarzbach and A. Weinstein (2005)

The **relative modular class** of the Lie algebroid morphism  $\varphi : A \rightarrow B$  is

$$\text{mod } \varphi(A, B) = \text{mod } A - \varphi^*(\text{mod } B)$$

- $\text{mod } \rho^A(A, TM) = \text{mod } A.$

## Y. Kosmann-Schwarzbach and A. Weinstein (2005)

The **relative modular class** of the Lie algebroid morphism  $\varphi : A \rightarrow B$  is

$$\text{mod}^\varphi(A, B) = \text{mod } A - \varphi^*(\text{mod } B)$$

- $\text{mod}^{\rho_A}(A, TM) = \text{mod } A$ .
- $\text{mod}^{P^\sharp}(A^*, A) = 2 \text{mod}(A, P)$ .

# Nijenhuis operators

- $N \equiv$  Nijenhuis operator of  $A$ ;

# Nijenhuis operators

- $N \equiv$  Nijenhuis operator of  $A$ ;
- $A_N = (A, \rho \circ N, [ , ]_N)$  is a Lie algebroid,

$$[X, Y]_N = [NX, Y] + [X, NY] - N[X, Y], \quad X, Y \in \mathfrak{X}^1(A).$$

# Nijenhuis operators

- $N \equiv$  Nijenhuis operator of  $A$ ;
- $A_N = (A, \rho \circ N, [ , ]_N)$  is a Lie algebroid,

$$[X, Y]_N = [NX, Y] + [X, NY] - N[X, Y], \quad X, Y \in \mathfrak{X}^1(A).$$

- $N : A_N \rightarrow A$  is a Lie algebroid morphism;

# Nijenhuis operators

- $N \equiv$  Nijenhuis operator of  $A$ ;
- $A_N = (A, \rho \circ N, [, ]_N)$  is a Lie algebroid,

$$[X, Y]_N = [NX, Y] + [X, NY] - N[X, Y], \quad X, Y \in \mathfrak{X}^1(A).$$

- $N : A_N \rightarrow A$  is a Lie algebroid morphism;

## Theorem

Fix  $\eta \otimes \mu$  section of  $Q_A \wedge \text{top}(A) \otimes \wedge^{\text{top}}(T^*M)$ . Then

$$\xi_{A_N} = d_A(\text{Tr } N) + N^*\xi_A.$$

- The 1-form

$$\xi_{A_N} - N^* \xi_A = d_A \text{Tr } N$$

is independent of the choice of section of  $Q_A$ .

- The 1-form

$$\xi_{A_N} - N^* \xi_A = d_A \text{Tr } N$$

is independent of the choice of section of  $Q_A$ .

$[d_A \text{Tr } N] \in H^1(A_N)$  is the relative modular class of the Lie algebroid morphism  $N : A_N \rightarrow A$ .



$(A, P, N)$ : Poisson-Nijenhuis Lie algebroid

$$\begin{array}{ccc}
 (A^*, [\cdot, \cdot]_{NP}) & \xrightarrow{N^*} & (A^*, [\cdot, \cdot]_P) \\
 \downarrow P^\sharp & \searrow NP^\sharp & \downarrow P^\sharp \\
 (A, [\cdot, \cdot]_N) & \xrightarrow{N} & (A, [\cdot, \cdot]_A)
 \end{array}$$

- $N^*$  is a Nijenhuis operator of the dual Lie algebroid

$$(A^*, [\ , \ ]_\rho, \rho \circ P^\sharp).$$

- $N^*$  is a Nijenhuis operator of the dual Lie algebroid

$$(A^*, [\cdot, \cdot]_{\rho}, \rho \circ P^{\sharp}).$$

- The relative modular class of  $N^*$  has the canonical representative  $d_{\rho}(\text{Tr } N^*)$ , so that:

$$\text{mod}^{N^*}(A_{N^*}^*, A^*) = [d_{\rho}(\text{Tr } N^*)] = [d_{\rho}(\text{Tr } N)].$$

- $N^*$  is a Nijenhuis operator of the dual Lie algebroid

$$(A^*, [\cdot, \cdot]_P, \rho \circ P^\sharp).$$

- The relative modular class of  $N^*$  has the canonical representative  $d_P(\text{Tr } N^*)$ , so that:

$$\text{mod}^{N^*}(A_{N^*}^*, A^*) = [d_P(\text{Tr } N^*)] = [d_P(\text{Tr } N)].$$

### Definition

The **modular vector field** of the Poisson-Nijenhuis Lie algebroid  $(A, P, N)$  is

$$X_{(N,P)} = \xi_{A_{N^*}^*} - N\xi_{A^*} = d_P(\text{Tr } N) = -P^\sharp(d_A \text{Tr } N) \in \Gamma(A).$$

## Theorem

*Suppose  $N$  is non-degenerated Nijenhuis operator.*

- $X_{(N,P)}$  is a  $d_{NP}$ -coboundary;

## Theorem

Suppose  $N$  is non-degenerated Nijenhuis operator.

- $X_{(N,P)}$  is a  $d_{NP}$ -coboundary;
- A hierarchy of vector fields

$$X_{(N,P)}^{i+j} = N^{i+j-1} X_{(N,P)} = d_{N^i P} h_j = d_{N^j P} h_i, \quad (i, j \in \mathbb{Z})$$

where  $h_0 = \ln(\det N)$  and  $h_i = \frac{1}{i} \text{Tr } N^i$ ,  $(i \neq 0)$ ;

## Theorem

Suppose  $N$  is non-degenerated Nijenhuis operator.

- $X_{(N,P)}$  is a  $d_{NP}$ -coboundary;
- A hierarchy of vector fields

$$X_{(N,P)}^{i+j} = N^{i+j-1} X_{(N,P)} = d_{N^i P} h_j = d_{N^j P} h_i, \quad (i, j \in \mathbb{Z})$$

where  $h_0 = \ln(\det N)$  and  $h_i = \frac{1}{i} \text{Tr } N^i$ ,  $(i \neq 0)$ ;

- A hierarchy of vector fields on  $M$  given by:

$$X_{i+j} = -\pi_i^\sharp dh_j = -\pi_j^\sharp dh_i \quad (i, j \in \mathbb{Z})$$

where  $\pi_j$  are Poisson structures on  $M$ .

The hierarchy of modular vector fields  $X_{(N^i, P)}$  is generated by the Nijenhuis operator  $N$ .



The hierarchy of modular vector fields  $X_{(N^i, P)}$  is generated by the Nijenhuis operator  $N$ .

BUT

The hierarchy of modular vector fields  $X_{(N^i, P)}$  is generated by the Nijenhuis operator  $N$ .

BUT

The covered hierarchy of bi-Hamiltonian vector fields on  $M$  may not be generated by any Nijenhuis operator.

## The two canonical examples

- Tangent bundle:  
P. Damianou and R. L. Fernandes [arXiv:math/0607784]  
Y. Kosmann and F. Magri [arXiv:math/0611202].
- Lie algebra  $(\mathfrak{g}, r, N)$ :  
 $N$  defines a sequence of modular forms on  $\mathfrak{g}^*$ ,  $\xi_{\mathfrak{g}_{N^k}^*}$ , which are associated with the higher brackets on  $\mathfrak{g}^*$ .

### Relation between modular forms of $\mathfrak{g}^*$

$$\xi_{\mathfrak{g}_{N^k}^*} = N^k \xi_{\mathfrak{g}_{N^*}^*}.$$

# Jacobi algebroid

A **Jacobi algebroid** is a Lie algebroid  $A$  equipped with a closed 1-form  $\phi$

# Jacobi algebroid

A **Jacobi algebroid** is a Lie algebroid  $A$  equipped with a closed 1-form  $\phi$

**Schouten-Jacobi bracket** on  $\mathfrak{X}(A)$ :

$$[P, Q]^\phi = [P, Q] + (p-1)P \wedge i_\phi Q - (-1)^{p-1}(q-1)i_\phi P \wedge Q,$$

for  $P \in \mathfrak{X}^p(A)$ ,  $Q \in \mathfrak{X}^q(A)$ .

# Poissonization

Consider the Lie algebroid  $\hat{A} = A \times \mathbb{R}$  over  $M \times \mathbb{R}$ , with Lie bracket

$$[X, Y]_{\hat{A}} = [X, Y], \quad X, Y \in \mathfrak{X}(A),$$

# Poissonization

Consider the Lie algebroid  $\hat{A} = A \times \mathbb{R}$  over  $M \times \mathbb{R}$ , with Lie bracket

$$[X, Y]_{\hat{A}} = [X, Y], \quad X, Y \in \mathfrak{X}(A),$$

and anchor

$$\hat{\rho}(X) = \rho(X) + \langle \phi, X \rangle \frac{\partial}{\partial t}, \quad X \in \mathfrak{X}^1(A). \quad (1)$$

# Poissonization

Consider the Lie algebroid  $\hat{A} = A \times \mathbb{R}$  over  $M \times \mathbb{R}$ , with Lie bracket

$$[X, Y]_{\hat{A}} = [X, Y], \quad X, Y \in \mathfrak{X}(A),$$

and anchor

$$\hat{\rho}(X) = \rho(X) + \langle \phi, X \rangle \frac{\partial}{\partial t}, \quad X \in \mathfrak{X}^1(A). \quad (1)$$

**induced Lie algebroid structure from  $A$  by  $\phi$ .**



# Poissonization

- $\phi = \hat{d}t$ ;

# Poissonization

- $\phi = \hat{d}t$ ;
- $\tilde{P} = e^{-(p-1)t}P, \quad P \in \mathfrak{X}^p(A)$ ;

# Poissonization

- $\phi = \hat{d}t$ ;
- $\tilde{P} = e^{-(\rho-1)t}P, \quad P \in \mathfrak{X}^\rho(A)$ ;
- $[\tilde{P}, \tilde{Q}]_{\hat{A}} = \widetilde{[P, Q]}^\phi$ ;

# Poissonization

- $\phi = \hat{d}t$ ;
- $\tilde{P} = e^{-(\rho-1)t}P, \quad P \in \mathfrak{X}^\rho(A)$ ;
- $[\tilde{P}, \tilde{Q}]_{\hat{A}} = \widetilde{[P, Q]}^\phi$ ;
- $[P, P]^\phi = 0 \iff \tilde{P} = e^{-t}P$  a Poisson bivector of  $\hat{A}$

# Poissonization

- $\phi = \hat{d}t$ ;
- $\tilde{P} = e^{-(\rho-1)t}P, \quad P \in \mathfrak{X}^\rho(A)$ ;
- $[\tilde{P}, \tilde{Q}]_{\hat{A}} = \widetilde{[P, Q]}^\phi$ ;
- $[P, P]^\phi = 0 \iff \tilde{P} = e^{-t}P$  a Poisson bivector of  $\hat{A}$   
 $(\hat{A}^*, [ , ]_{\tilde{P}}, \hat{\rho} \circ \tilde{P}^\sharp)$  is a Lie algebroid

# Poissonization

- $\hat{\alpha} = e^t \alpha, \quad \alpha \in \Omega^1(A);$

# Poissonization

- $\hat{\alpha} = e^t \alpha, \quad \alpha \in \Omega^1(A);$
- $$\left[ \hat{\alpha}, \hat{\beta} \right]_{\tilde{P}} = e^t \underbrace{(\mathcal{L}_{P\#\alpha}^\phi \beta - \mathcal{L}_{P\#\beta}^\phi \alpha - d^\phi P(\alpha, \beta))}_{[\alpha, \beta]_P}$$

# Poissonization

- $\hat{\alpha} = e^t \alpha, \quad \alpha \in \Omega^1(A);$
- $$[\hat{\alpha}, \hat{\beta}]_{\tilde{P}} = e^t \underbrace{(\mathcal{L}_{P\sharp\alpha}\beta - \mathcal{L}_{P\sharp\beta}\alpha - d^\phi P(\alpha, \beta))}_{[\alpha, \beta]_P}$$
- $(A^*, [ , ]_P, \rho \circ P\sharp)$  is a Lie algebroid;



# Poissonization

- $\hat{\alpha} = e^t \alpha, \quad \alpha \in \Omega^1(A);$
- $$\left[ \hat{\alpha}, \hat{\beta} \right]_{\hat{P}} = e^t \underbrace{(\mathcal{L}_{P\sharp\alpha}\beta - \mathcal{L}_{P\sharp\beta}\alpha - d^\phi P(\alpha, \beta))}_{[\alpha, \beta]_P}$$
- $(A^*, [ , ]_P, \rho \circ P\sharp)$  is a Lie algebroid;
- $N$  Nijenhuis operator of  $A \implies N$  Nijenhuis operator of  $\hat{A}$

# Poissonization

- $\hat{\alpha} = e^t \alpha, \quad \alpha \in \Omega^1(A);$
- $$\left[ \hat{\alpha}, \hat{\beta} \right]_{\tilde{P}} = e^t \underbrace{(\mathcal{L}_{P\sharp\alpha}\beta - \mathcal{L}_{P\sharp\beta}\alpha - d^\phi P(\alpha, \beta))}_{[\alpha, \beta]_P}$$
- $(A^*, [ , ]_P, \rho \circ P\sharp)$  is a Lie algebroid;
- $N$  Nijenhuis operator of  $A \implies N$  Nijenhuis operator of  $\hat{A}$

## Definition

$(A, \phi, P, N)$  is a **Jacobi-Nijenhuis algebroid** if  $(\hat{A}, \tilde{P}, N)$  is a Poisson-Nijenhuis algebroid.

$(\hat{A}, \tilde{P}, N)$  has a modular vector field

$$\hat{X}_{(N, \tilde{P})} = \xi_{\hat{A}_{N^*}} - N\xi_{\hat{A}^*} = e^{-t} d_P(\text{Tr } N).$$

$(\hat{A}, \tilde{P}, N)$  has a modular vector field

$$\hat{X}_{(N, \tilde{P})} = \xi_{\hat{A}_{N^*}} - N\xi_{\hat{A}^*} = e^{-t} d_P(\text{Tr } N).$$

### Definition

The **modular vector field** of the Jacobi-Nijenhuis algebroid  $(A, \phi, P, N)$  is

$$X_{(N, P)} = \xi_{A_{N^*}} - N\xi_{A^*}.$$

$(\hat{A}, \tilde{P}, N)$  has a modular vector field

$$\hat{X}_{(N, \tilde{P})} = \xi_{\hat{A}_{N^*}} - N\xi_{\hat{A}^*} = e^{-t} d_P(\text{Tr } N).$$

### Definition

The **modular vector field** of the Jacobi-Nijenhuis algebroid  $(A, \phi, P, N)$  is

$$X_{(N, P)} = \xi_{A_{N^*}} - N\xi_{A^*}.$$

$\text{mod}^{(N, P)} A = [X_{(N, P)}]$  the **modular class** of  $(A, \phi, P, N)$ .

## Also...

If  $N$  is non-degenerated, then

- $\hat{X}_{(N, \tilde{P})}^{i+j} = N^{i+j-1} \hat{X}_{(N, \tilde{P})} = d_{N^i \tilde{P}} h_j = d_{N^i \tilde{P}} h_i \in \Gamma \hat{A}$
- $X_{(N, P)}^{i+j} = N^{i+j-1} X_{(N, P)} = d_{N^i P} h_j = d_{N^i P} h_i \in \Gamma(A)$

$$h_0 = \ln(\det N) \quad \text{and} \quad h_i = \frac{1}{i} \text{Tr } N^i, \quad (i \neq 0, \quad i, j \in \mathbb{Z}).$$

- These hierarchies cover two hierarchies, one on  $M \times \mathbb{R}$  and another one on  $M$ .

## Homework

- A. Weinstein, The modular automorphism group of a Poisson manifold, *J. Geom. Phys.* **23** (1997), 379–394.
- S. Evens, J.-H. Lu and A. Weinstein, Transverse measures, the modular class and a cohomology pairing for Lie algebroid, *Quart. J. Math. Oxford (2)* **50** (1999), 417–436.
- P. Xu, Gerstanheber algebras and BV-algebras in Poisson geometry, *Comm. Math. Phys.* **200** (1999), 545–560.
- Y. Kosmann-Schwarzbach, Modular vector fields and Batalin-Vilkoviski algebras, *Poisson Geometry*, eds. J. Grabowski and P. Urbanski, Banach Center Publications, **51** (2000), 109–129.
- R. Caseiro, Modular classes of Poisson-Nijenhuis Lie algebroids, *Lett. Math. Phys.*, to appear. Preprint [math.DG/0701476](https://arxiv.org/abs/math/0701476).

# Homework

- I. Vaisman, The BV-algebra of a Jacobi algebroid, *Ann. Polon. Math.* **73** (2000), 275–290.
- D. Iglesias, B. López, J.C. Marrero and E. Padrón, Triangular generalized Lie bialgebroids: homology and cohomology theories, *Lie algebroids*, eds. J. Grabowski and P. Urbanski, Banach Center Publications, **54** (2001), 111–133.
- R. Caseiro and J. Nunes da Costa, Jacobi-Nijenhuis algebroids and their modular classes, *J. Phys. A*, to appear. *arXiv:0706.1475*